

# On The Consecutive Integers $n + i - 1 = (i + 1)P_i$

Chun-Xuan Jiang

P. O. Box 3924, Beijing 100854, China

and Institute for Basic Research

P.O .Box 1577, Palm Haror, FL 34682. U. S. A

Jiang chunxuan@vip.sohu.com

## Abstract

By using the Jiang's function  $J_2(\omega)$  we prove that there exist infinitely many integers  $n$  such that  $n = 2P_1$ ,  $n + 1 = 3P_2, \dots$ ,  $n + k - 1 = (k + 1)P_k$  are all composites for arbitrarily long  $k$ , where  $P_1, P_2, \dots, P_k$  are all primes. This result has no prior occurrence in the history of number theory.

AMS Mathematics subject classification: Primary 11N05, 11N32.

**Theorem 1.** There exist infinitely many integers  $n$  such that the consecutive integers  $n = 2P_1$ ,  $n + 1 = 3P_2, \dots$ ,  $n + k - 1 = (k + 1)P_k$  are all composites for arbitrarily long  $k$ , where  $P_1, P_2, \dots, P_k$  are all primes.

Proof. Suppose that  $m = \prod_{i=1}^k (i + 1)$ . We define the prime equations

$$P_i = \frac{m}{i + 1} x + 1, \quad (1)$$

Where  $i = 1, 2, \dots, k$ .

The Jiang's function [1] is

$$J_2(\omega) = \prod_{3 \leq P} (P - k - 1 - \chi(P)) \neq 0 \quad (2)$$

where  $\chi(P) = -k$  if  $P^2 | m$ ;  $\chi(P) = -k + 1$  if  $P | m$ ;  $\chi(P) = 0$  otherwise,

$$\omega = \prod_{2 \leq P} P.$$

Since  $J_2(\omega) \rightarrow \infty$  as  $\omega \rightarrow \infty$ , there exist infinitely many integers  $x$  such that  $P_1, P_2, \dots, P_k$  are all primes.

We have the asymptotic formula of the number of integers  $x \leq N$  [1]

$$\pi_{k+1}(N, 2) \sim \frac{J_2(\omega) \omega^k}{\phi^{k+1}(\omega)} \frac{N}{\log^{k+1} N}, \quad (3)$$

where  $\phi(\omega) = \prod_{2 \leq P} (P - 2)$ .

From (1) we have  $n = mx + 2 = 2 \left( \frac{mx}{2} + 1 \right) = 2P_1$ ,  $n + 1 = mx + 3 = 3 \left( \frac{m}{3} x + 1 \right)$

$$= 3P_2, \dots, n + k - 1 = mx + k + 1 = (k + 1) \left( \frac{m}{k + 1} x + 1 \right) = (k + 1)P_k.$$

**Example 1.** Let  $k = 5$ , we have  $n = 2 \times 53281$ ,  $n + 1 = 3 \times 35521$ ,  $n + 2 = 4 \times 26641$ ,  $n + 3 = 5 \times 21313$ ,  $n + 4 = 6 \times 17761$ .

**Theorem 2.** There exist infinitely many integers  $n$  such that the consecutive integers

$n = (1 + 2^b)P_1$ ,  $n + 1 = (2 + 2^b)P_2, \dots$ ,  $n + k - 1 = (k + 2^b)P_k$  are all composites for arbitrarily long  $k$ , where  $P_1, P_2, \dots, P_k$  are all primes.

Proof. Suppose that  $m = \prod_{i=1}^k (i + 2^b)$ . We define the prime equations

$$P_i = \frac{m}{i + 2^b} x + 1, \quad (4)$$

Where  $i = 1, 2, \dots, k$ .

The Jiang's function [1] is

$$J_2(\omega) = \prod_{3 \leq P} (P - k - 1 - \chi(P)) \neq 0 \quad (5)$$

where  $\chi(P) = -k$  if  $P^2 \mid m$ ;  $\chi(P) = -k + 1$  if  $P \mid m$ ;  $\chi(P) = 0$  otherwise.

Since  $J_2(\omega) \rightarrow \infty$  as  $\omega \rightarrow \infty$ , there exist infinitely many integers  $x$  such that  $P_1, P_2, \dots, P_k$  are all primes.

We have the asymptotic formula of the number of integers  $x \leq N$  [1]

$$\pi_{k+1}(N, 2) \sim \frac{J_2(\omega) \omega^k}{\phi^{k+1}(\omega)} \frac{N}{\log^{k+1} N}, \quad (6)$$

From (4) we have  $n = mx + 1 + 2^b = (1 + 2^b) \left( \frac{m}{1 + 2^b} x + 1 \right) = (1 + 2^b)P_1$ ,  $n + 1 =$

$mx + 2 + 2^b = (2 + 2^b) \left( \frac{m}{2 + 2^b} x + 1 \right) = (2 + 2^b)P_2, \dots$ ,  $n + k - 1 = mx + k + 2^b =$

$(k + 2^b) \left( \frac{m}{k + 2^b} x + 1 \right) = (k + 2^b)P_k$ .

**Example 2.** Let  $b = 1$  and  $k = 4$ , we have  $n = 3 \times 27361$ ,  $n + 1 = 4 \times 20521$ ,  $n + 2 = 5 \times 16417$ ,  $n + 3 = 6 \times 13681$ .

**Theorem 3.** There exist infinitely many integers  $n$  such that the consecutive integers  $n = 3P_1$ ,  $n + 2 = 5P_2, \dots, n + 2(k - 1) = (2k + 1)P_k$  are all composites for arbitrarily long  $k$ , where  $P_1, P_2, \dots, P_k$  are all primes.

Proof. Suppose that  $m = \prod_{i=1}^k (2i+1)$ . We define the prime equations

$$P_i = \frac{m}{2i+1}x + 1, \quad (7)$$

Where  $i = 1, 2, \dots, k$ .

The Jiang's function [1] is

$$J_2(\omega) = \prod_{3 \leq P} (P - k - 1 - \chi(P)) \neq 0 \quad (8)$$

where  $\chi(P) = -k$  if  $P^2 \mid m$ ;  $\chi(P) = -k + 1$  if  $P \mid m$ ;  $\chi(P) = 0$  otherwise.

Since  $J_2(\omega) \rightarrow \infty$  as  $\omega \rightarrow \infty$ , there exist infinitely many integers  $x$  such that  $P_1, P_2, \dots, P_k$  are all primes.

We have the asymptotic formula of the number of integers  $x \leq N$  [1]

$$\pi_{k+1}(N, 2) \sim \frac{J_2(\omega)\omega^k}{\phi^{k+1}(\omega)} \frac{N}{\log^{k+1} N}, \quad (9)$$

From (7) we have  $n = mx + 3 = 3\left(\frac{m}{3}x + 1\right) = 3P_1$ ,  $n + 2 = mx + 5 = 5\left(\frac{m}{5}x + 1\right) =$

$$5P_2, \dots, n + 2(k-1) = mx + 2k + 1 = (2k+1)\left(\frac{m}{2k+1}x + 1\right) = (2k+1)P_k.$$

**Example 3.** Let  $k = 4$ , we have  $n = 3 \times 631$ ,  $n + 2 = 5 \times 379$ ,  $n + 4 = 7 \times 271$ ,  $n + 6 = 9 \times 211$ .

**Theorem 4.** There exist infinitely many integers  $n$  such that the consecutive integers  $n = P_1$ ,  $n + 2 = 3P_2, \dots, n + 2(k-1) = (2k-1)P_k$  are all composites for arbitrarily long  $k$ , where  $P_1, P_2, \dots, P_k$  are all primes.

Proof. Suppose that  $m = \prod_{i=1}^k (2i-1)$ . We define the prime equations

$$P_i = \frac{m}{2i-1}x + 1 \quad (10)$$

where  $i = 1, 2, \dots, k$ .

The Jiang's function [1] is

$$J_2(\omega) = \prod_{3 \leq P} (P - k - 1 - \chi(P)) \neq 0 \quad (11)$$

where  $\chi(P) = -k$  if  $P^2 | m$ ;  $\chi(P) = -k + 1$  if  $P | m$ ;  $\chi(P) = 0$  otherwise.

Since  $J_2(\omega) \rightarrow \infty$  as  $\omega \rightarrow \infty$ , there exist infinitely many integers  $x$  such that  $P_1, P_2, \dots, P_k$  are all primes.

We have the asymptotic formula of the number of integers  $x \leq N$  [1]

$$\pi_{k+1}(N, 2) \sim \frac{J_2(\omega)\omega^k}{\phi^{k+1}(\omega)} \frac{N}{\log^{k+1} N}, \quad (12)$$

From (10) we have  $n = P_1 = mx + 1$ ,  $n + 2 = mx + 3 = 3\left(\frac{m}{3}x + 1\right) = 3P_2, \dots$ ,

$$n + 2(k - 1) = mx + 2(k - 1) = (2k - 1)\left(\frac{m}{2k - 1}x + 1\right) = (2k - 1)P_k.$$

**Example 4.** Let  $k = 4$ , we have  $n = 9661$ ,  $n + 2 = 3 \times 3221$ ,  $n + 4 = 5 \times 1933$ ,  $n + 6 = 7 \times 1381$ .

**Theorem 5.** There exist infinitely many integers  $n$  such that the consecutive integers  $n = 3P_1, n + 4 = 7P_2, \dots, n + 4(k - 1) = (4k - 1)P_k$  are all composites for arbitrarily long  $k$ , where  $P_1, P_2, \dots, P_k$  are all primes.

**Example 5.** Let  $k = 4$ , we have  $n = 3 \times 2311$ ,  $n + 4 = 7 \times 991$ ,  $n + 8 = 11 \times 631$ ,  $n + 12 = 15 \times 463$ .

**Theorem 6.** There exist infinitely many integers  $n$  such that the consecutive integers  $n = 5P_1, n + 4 = 9P_2, \dots, n + 4(k - 1) = (4k + 1)P_k$  are all composites for arbitrarily long  $k$ , where  $P_1, P_2, \dots, P_k$  are all primes.

#### Reference

[1] Chun-Xuan Jiang. Foundations of Santilli's isonumber theory with applications to new cryptograms, Fermat's theorem and Goldbach's conjecture. International Academic Press, 2002 MR 2004c: 11001. <http://www.i-b-r.org/docs/jiang/pdf>.