

How can GW/graviton generation has semi classical features, and what the implications are for cosmological re acceleration of the universe, and initial relic entropy generation in term of the importance of dimensions.

Andrew Beckwith

American institute of Beam energy propulsion, life member

beckwith@aibep.org, abeckwith@uh.edu

Abstract.

We argue in this document that initial vacuum state values possibly responsible for GW generation in relic conditions in the initial onset of inflation may have a temporary un squeezed , possibly even coherent initial value, which would permit in certain models classical coherent initial gravitational wave states. Secondly, if gravitational waves have semi classical features in their genesis, then the problem of how $q(Z)$ under goes through an inflection point and how the rate of acceleration increases a billion years ago needs to be re examined. Additional dimensions are not the problem. Determining if GW/Gravitons are semi classical in their initial formulation at the source of the big bang, is. Also, semi classical and quantum entropy generating processes are compared , with the role of looking for over lap between them..

Introduction

The author finds that the supposition as to the inevitability of low frequency GW from the big bang is supported only by the conclusion that large spatial dimensions above our four dimensions are conduits as to dumping cyclical universe matter-energy into. Numerous articles, i.e. one ‘dumping inflation’ “out of this world” , as well as the multiple universes popularized by Arkani-Hamid , among others, with most of the energy of relic inflation dumped into higher dimensions, than our present 4 dimensional space time. But notice the assumptions made to back up this claim. I.e. very large scale higher dimensions. The Calabi Yau manifold used as an initial embedding space for dimensions above our space time was for a long time thought of as having compact, almost undetectable small higher dimensions. The initial smallness of the higher dimensions was the reason why Gasperini and others as of (1995) wrote well received string theory articles predicting no favoring of low frequency GW as the primary relic GW signature from the big bang.

Invariance of photon equations of state to preserve black body radiation profiles, and the aftermath

Before 380 thousand years after the big bang, there was still photon related cosmological evolutions as defined by [J. A. S. Lima](#) (1996), which can be summarized, for temperature related behavior as photons

having number and energy densities specified as $n_r|_{Photons} \sim T^3, \rho_r|_{Photons} \sim T^4$, so that for an

instantaneous co moving number of photons, Lima write $N_r|_{Photons} = n_r|_{Photons} \cdot T$, where T is for

background temperature and states that this value $N_r|_{Photons}$ must be a constant. Lima quotes a researcher, Steigman in saying that “Unless the number of co moving photons in a co moving volume is constant, a blackbody distribution (of photons) is destroyed as the universe evolves”. In addition, Lima’s

key result which can be summarized as follows, that even if $N_r|_{Photons}$ has a changing time component that

there exists entropy associated with photons , $S_r|_{Photons}$ so that the following relationship holds for any

Friedman style cosmology, namely $\dot{S}_r|_{Photons} / S_r|_{Photons} = \dot{N}_r|_{Photons} / N_r|_{Photons}$, where the dot is time .

If what is suggested by Beckwith (2009), with respect to his revision of Y.J. Ng’s counting algorithm is correct, with respect to early universe conditions is correct, i.e.

$\dot{S}_r|_{gravitons} / S_r|_{gravitons} = \dot{N}_r|_{gravitons} / N_r|_{gravitons}$ is also equal to a ratio of the time derivative of the

number of gravitons, over the number of gravitons, and this in term is equal to the time derivative of entropy of graviton production, over entropy of graviton production at the onset of the universe, then in fact what one is working with is, de facto, one is looking at , then for initial conditions of

$$\dot{S}_r \Big|_{Photons} / S_r \Big|_{Photons} = \dot{N}_r \Big|_{Photons} / N_r \Big|_{Photons} \sim \dot{S}_r \Big|_{gravitons} / S_r \Big|_{gravitons} = \dot{N}_r \Big|_{gravitons} / N_r \Big|_{gravitons} \quad (0)$$

This should be a starting point to the analysis which proceeds in this paper, I.e. Eqn. (.0) as compared with the $S_{max} = \pi/H^2 \sim 10^5$ or larger at the origins of the big bang will be a starting point in information /data comparison.

So, what can be said about the Y.J. Ng paradigm of entropy generation, which Beckwith has modified and looked at ? For a start, consider if the counting algorithm , which is a string theory result , can have any common results with a quantum gas result, which comes from the WDW equation, whose solution is WKB, semi classical in nature?

The question of relative over lap of classical and quantum processes in terms of wave functions for the evolution of the universe will be crucially important in determining coherency issues as far as relic GW, and gravitons from relic conditions, which the author will return to repeatedly during this presentation.

Review of simple models as to gravitons as produced either by (Quantum gravity) strings , LQG,(or by processes which may not be Quantum Gravity based?)

We wish now to review what may be some of the counting algorithms appropriate for entropy generation, and which may contribute to answering if or not GW are mandated to be, from the beginning either a classical versus a quantum processes. IN part this next page is due to concepts A.Beckwith presented in Rencontres De Blois, 2009, and is a starting point for our inquiry as to the necessity, or lack of , of modeling Gravity as either classical / quantum based in relic conditions.

Introduction w.r.t. the NG paradigm

We wish to present two alternative routes to generation of entropy. The first, is a counting algorithm, is an adaptation of Y.J. Ng's infinite quantum (modified Boltzmann's) statistics; the second references A. Glinka's research presentation on "graviton gas" as a way to provide a perspective? as to how to get a partition function for gravitons that is congruent with the Wheeler De Witt equation. Here are a few questions which are posed for the reader.

1. Is each "particle count unit" as suggested by Ng equivalent to a brane-antibrane unit in brane treatments of entropy?
2. Is the change of entropy $\Delta S \approx \Delta N_{gravitons}$?
3. Is this graviton production scheme comparable to Glinka's quantum gas , from the Wheeler De Witt equation?

Entropy generation via Ng's infinite quantum statistics (short review)

This discussion is motivated to show a purely string theory approach and to see if its predictions may over lap with semi classical WDM (semi classical) treatments of cosmology.. The contention being advanced is that if there is an over lap between these two methods, that it may aid in obtaining experimentally falsifiable data sets for GW from relic conditions. We wish to understand the linkage between dark matter and gravitons. how relic gravitational waves relate to relic gravitons? To consider just that, we look at the "size" of the nucleation space, V for dark matter, DM. V for nucleation is HUGE. Graviton space V for nucleation is tiny , well inside inflation. Therefore, the log factor drops OUT of entropy S if V chosen properly for both eqn 1 and eqn 2. Ng's result begins with a modification of the entropy/ partition function Ng used the following approximation of temperature and its variation with respect to a spatial parameter, starting with temperature $T \approx R_H^{-1}$ (R_H can be thought of as a representation of the region of space where

we take statistics of the particles in question). Furthermore, assume that the volume of space to be analyzed is of the form $V \approx R_H^3$ and look at a preliminary numerical factor we shall call $N \sim (R_H/l_p)^2$, where the denominator is Planck's length (on the order of 10^{-35} centimeters). We also specify a "wavelength" parameter $\lambda \approx T^{-1}$. So the value of $\lambda \approx T^{-1}$ and of R_H are approximately the same order of magnitude. Now this is how Jack Ng changes conventional statistics: he outlines how to get $S \approx N$, which with additional arguments we refine to be $S \approx \langle n \rangle$ (where $\langle n \rangle$ is graviton density). Begin with a partition function

$$Z_N \sim \left(\frac{1}{N!}\right) \cdot \left(\frac{V}{\lambda^3}\right)^N \quad (1)$$

This, according to Ng, leads to entropy of the limiting value of, if $S = (\log[Z_N])$

$$S \approx N \cdot (\log[V/N\lambda^3] + 5/2) \xrightarrow{\text{Ng-inf inite-Quantum-Statistics}} N \cdot (\log[V/\lambda^3] + 5/2) \approx N \quad (2)$$

But $V \approx R_H^3 \approx \lambda^3$, so unless N in Eqn (2) above is about 1, S (entropy) would be < 0 , which is a contradiction. Now this is where Jack Ng introduces removing the $N!$ term in Eqn (1) above, i.e., inside the Log expression we remove the expression of N in Eqn. (0.2) above. The modification of Ng's entropy expression is in the region of space time for which the general temperature dependent entropy Kolb and Turner expression breaks down. In particular, the evaluation of entropy we do via the modified Ng argument above is in regions of space time where g before re heat is an unknown, unmeasurable number of degrees of freedom. The Kolb and Turner entropy expression (1991) has a temperature T related entropy density which leads to that we are able to state total entropy as the entropy density time's space time volume V_4 with $g_{re-heat} \approx 1000$, according to De Vega, while dropping to $g_{electro-weak} \approx 100$ in the electro weak era. This value of the space time degrees of freedom, according to de Vega has reached a low of $g_{today} \approx 2-3$ today. We assert that Eqn (2) above occurs in a region of space time before $g_{re-heat} \approx 1000$, so after re heating Eqn (2) no longer holds, and we instead can look at

$$S_{total} \equiv s_{Density} \cdot V_4 = \frac{2\pi^2}{45} \cdot g \cdot T^3 \cdot V_4 \quad (3)$$

Where $T < 10^{32} K$. We can compare eqn (1) and (2), as how they stack up with Glinka's (2007) quantum gas, if we

identify $\Omega = \frac{1}{2|u|^2 - 1}$ as a partition function (with u part of a Bogoliubov transformation) due to a

graviton-quintessence gas, to get information theory based entropy

$$S \equiv \ln \Omega \quad (4)$$

Such a linkage would open up the possibility that the density of primordial gravitational waves could be examined, and linked to modeling gravity as an effective theory. The details of linking what is done with (0.2) and bridging it to (0.3) await additional theoretical development, and are probably conceptually understandable if the following is used to link the two regimes. I.e. we can use the number of space time operations used to create (0.2), via Seth Lloyds

$$I = S_{total} / k_B \ln 2 = [\#operations]^{3/4} = [\rho \cdot c^5 \cdot t^4 / \hbar]^{3/4} \quad (5)$$

Essentially, what will be done is to use 5 to show linkage between a largely thermally based production of entropy, as implied by (3) and a particle counting algorithm, as given by (0.2). This due to the problems inherent in making connections between a particle count generation of entropy, and thermal contributions. I.e two different processes are involved. The big news is though that the WKB is semi classical, whereas anything from string theory is, well, QFT, plus.

Where there is an overlap between a classical wave function, and its quantum mechanical analog, that means there is a minimization of spreading of a wave functional. i.e. see Roy Glauber (1963) One can say the following. That if there is an overlap between the Wheeler De Witt equation derived quantum gas which was brought up by Glinka (2007), where the WDW can have WKB semi classical solutions, and the string theory counting algorithm, Then, if the end results are similar, the fact is that the quantum procedure, i.e. brane theory, is overlapping with WKB, means that there is a minimization of uncertainty. Note that the supposition of how classical and quantum processes can give similar answers is presented in rich detail by Roy Glauber(1963) and the example talked about here is its GW analog. Gravitons are stated conceptually to be akin to photons in light waves. In simple physics analogies. But this simple quantum generalization breaks down, since gravitons are spin two particles with a complex set of interactions not only with themselves, but with evolving space time geometry. We mention that gravitons may be important to initial entropy generation. Entropy generation and entropy perturbations affect the gaussianity of evolving wave functions of matter and energy evolving in space time. If there is a large deviation of the initially Gaussian states of space time wave functions, there is likely a break from classical physics due to the complexity of evolving wave function states influenced increasingly by non Gaussian perturbations. This non Gaussian process is reflected by marked deviation from planar wave state approximations used in the evolution of wave functions Hence the issue of apparently combined sources of planar wave generation of gravitational waves is a precursor to what would happen if squeezed states occurred at the onset of the big bang. I.e., what would happen with multiple superpositions of different coherent states?. A good reference as to coherent states in cosmology, as in this example, Bianchi I universes, was given by Brett Bolen, Luca Bombelli, Alejandro Corichi (2004) In particular, look at their equation 3.1. If states are largely coherent, such a small variation/ smoothness of observables will have observational consequences as to relic gravitational wave signals seen in the onset of inflation. The basic reason for making such an examination of the relative importance of squeezing/ lack of squeezing is to determine if or not relic GW are due to classical versus quantum Gravitational processes. The answer to if or not relic GW are due to classical versus quantum processes has huge consequences as to the dominant GW harmonics in terms of what are the most important frequencies researchers need to look for, for relic GW identification, with instrumentation. The problem facing GW researchers is how to find dominant sub harmonics, in GW signals, i.e. how to use pattern recognition, and updated advanced fourier analysis in order to identify dominant frequency ranges of GW signals which are of interest and which carry the most relevant physics information for cosmologists to review and learn from. How does one actually know about first or second order phase transitions, due to GW. Does one see, as an example classically based non linear superposition of GW, which have consequences as to admissible spectrum of GW frequencies to detect? Since it has been brought up, let us now review, briefly the issue of coherence, versus de coherence of initial vacuum states, and its relevance as to classical versus quantum factors as to generation of GWs

Issues about Coherent state of Gravitons (linking gravitons with GW)

In the quantum theory of light (quantum electrodynamics) and other bosonic quantum field theories, coherent states were introduced by the work of Roy J. Glauber in 1963 Now, it is well appreciated that Gravitons are NOT similar to light. So what is appropriate for presenting gravitons as coherent states? Coherent states, to first approximation are retrievable as minimum uncertainty states. If one takes string theory as a reference, the minimum value of uncertainty becomes part of a minimum uncertainty which can be written as given by Veneziano (1993), where $l_s \cong 10^\alpha \cdot l_{Planck}$, with $\alpha > 0$, and $l_{Planck} \approx 10^{-33}$ centimeters

$$\Delta x > \frac{\hbar}{\Delta p} + \frac{l_s^2}{\hbar} \cdot [\Delta p] \quad (6)$$

To put it mildly, if we are looking at a solution to minimize graviton position uncertainty, we will likely be out of luck if string theory is the only tool we have for early universe conditions. Mainly, the momentum will not be small, and uncertainty in momentum will not be small either. Either way, most likely, $\Delta x > l_s \cong 10^\alpha \cdot l_{Planck}$ In addition, it is likely, as Klaus Kieffer in the book “Quantum Gravity” on page 290 of that book that if gravitons are excitations of closed strings, then one will have to look for conditions for which a coherent state of gravitons, as stated by Mohaupt (2003) occurs. What Mohaupt is referring to is a string theory way to reproduce what Ford gave in 1995, i.e. conditions for how Gravitons in a

squeezed vacuum state, the natural result of quantum creation in the early universe will introduce metric fluctuations. Ford's (1995) treatment is to have a metric averaged retarded Green's function for a massless field becoming a Gaussian. The condition of Gaussianity is how to obtain semi classical , minimal uncertainty wave states, in this case de rigor for coherent wave function states to form. Ford uses gravitons in a so called 'squeezed vacuum state' as a natural template for relic gravitons. I.e. the squeezed vacuum state (a **squeezed coherent state**) is any state such that the uncertainty principle is saturated.: In QM coherence would be when $\Delta x \Delta p = \hbar/2$. In the case of string theory it would have to be

$$\Delta x \Delta p = \frac{\hbar}{2} + \frac{l_s^2}{2 \cdot \hbar} \cdot [\Delta p]^2 \quad (7)$$

Begin with noting that if one is not using string theory, we merely set the term $l_s \xrightarrow{\text{non-string}} 0$, but that we are still considering Roy Glauber(1963) with string theory replacing his stated example. However, what one sees in string theory, is a situation where a vacuum state as a template for graviton nucleation is built out of an initial vacuum state, $|0\rangle$. To do this though, as Venkatartnam, and Suresh did, involved using a

squeezing operator $Z[r, \vartheta]$ defining via use of a squeezing parameter r as a strength of squeezing interaction term , with $0 \leq r \leq \infty$, and also an angle of squeezing, $-\pi \leq \vartheta \leq \pi$ as used in $Z[r, \vartheta] = \exp\left[\frac{r}{2} \cdot \left([\exp(-i\vartheta)] \cdot a^2 - [\exp(i\vartheta)] \cdot a^{+2}\right)\right]$, where combining the $Z[r, \vartheta]$ with $|\alpha\rangle = D(\alpha) \cdot |0\rangle$ (8)

(8) leads to a single mode squeezed coherent state, as they define it via

$$|\zeta\rangle = Z[r, \vartheta] |\alpha\rangle = Z[r, \vartheta] D(\alpha) \cdot |0\rangle \xrightarrow{\alpha \rightarrow 0} Z[r, \vartheta] \cdot |0\rangle \quad (9)$$

The right hand side . of eqn. (1.16) given above becomes a highly non classical operator, i.e. in the limit that the super position of states $|\zeta\rangle \xrightarrow{\alpha \rightarrow 0} Z[r, \vartheta] \cdot |0\rangle$ occurs, there is a many particle version of a 'vacuum state' which has highly non classical properties. Squeezed states are thought to occur at the onset of vacuum nucleation , but what is noted for $|\zeta\rangle \xrightarrow{\alpha \rightarrow 0} Z[r, \vartheta] \cdot |0\rangle$ being a super position of vacuum states, means that classical analog is extremely difficult to recover in the case of squeezing, and general non

classical behavior of squeezed states. Can one, in any case, faced with $|\alpha\rangle = D(\alpha) \cdot |0\rangle \neq Z[r, \vartheta] \cdot |0\rangle$ do a better job of constructing coherent graviton states, in relic conditions, which may not involve squeezing ?. Note L. Grishchuk wrote in (1989) in "On the quantum state of relic gravitons", where he claimed in his abstract that 'It is shown that relic gravitons created from zero-point quantum fluctuations in the course of cosmological expansion should now exist in the squeezed quantum state. The authors have determined the parameters of the squeezed state generated in a simple cosmological model which includes a stage of inflationary expansion.'. Grishchuk , et al, (1989) reference their version of a cosmological

perturbation h_{nlm} via the following argument. How we work with the argument will affect what is said about the necessity, or lack of, of squeezed states in early universe cosmology. From Class. Quantum Gravity: 6 (1989), L 161-L165, where h_{nlm} has a component $\mu_{nlm}(\eta)$ obeying a parametric oscillator equation, where K is a measure of curvature which is $= \pm 1, 0$, $a(\eta)$ is a scale factor of a FRW metric, and $n = 2\pi \cdot [a(\eta)/\lambda]$ is a way to scale a wavelength, λ , with n , and with $a(\eta)$

$$h_{nlm} \equiv \frac{l_{plank}}{a(\eta)} \cdot \mu_{nlm}(\eta) \cdot G_{nlm}(x) \quad (10)$$

$$\mu_{nlm}''(\eta) + \left(n^2 - K - \frac{a''}{a}\right) \cdot \mu_{nlm}(\eta) \equiv 0 \quad (11)$$

If $y(\eta) = \frac{\mu(\eta)}{a(\eta)}$ is picked, and a Schrodinger equation is made out of the Lagrangian used to formulate

(11) above, with $\hat{P}_y = \frac{-i}{\partial y}$, and $M = a^3(\eta)$, $\Omega = \frac{\sqrt{n^2 - K^2}}{a(\eta)}$, $\vec{a} = [a(\eta)/l_{planck}] \cdot \sigma$, and $F(\eta)$ an

arbitrary function. $y' = \partial y / \partial \eta$. Also, we have a finite volume $V_{finite} = \int \sqrt{(3)} g d^3 x$

Then the Lagrangian for deriving (11) is (and leads to a Hamiltonian which can be **also** derived from the Wheeler De Witt equation), with $\zeta = 1$ for zero point subtraction of energy

$$L = \frac{M \cdot y'^2}{2a(\eta)} - \frac{M^2 \cdot \Omega^2 a \cdot y^2}{2} + a \cdot F(\eta) \quad (12)$$

$$\frac{-1}{i} \cdot \frac{\partial \psi}{a \cdot \partial \eta} \equiv \hat{H} \psi \equiv \left[\frac{\hat{P}_y^2}{2M} + \frac{1}{2} \cdot M \Omega^2 \hat{y}^2 - \frac{1}{2} \cdot \zeta \cdot \Omega \right] \cdot \psi \quad (13)$$

then there are two possible solutions to the S.E. Grushchuk created in 1989, one a non squeezed state, and another a squeezed state. So in general we work with

$$y(\eta) = \frac{\mu(\eta)}{a(\eta)} \equiv C(\eta) \cdot \exp(-B \cdot y) \quad (14)$$

The **non squeezed state** has a parameter $B|_{\eta} \xrightarrow{\eta \rightarrow \eta_b} B(\eta_b) \equiv \omega_b/2$ where η_b is an initial time, for which the Hamiltonian given in (14) in terms of raising/ lowering operators is ‘diagonal’, and then the rest of the time for $\eta \neq \eta_b$, the **squeezed state** for $y(\eta)$ is given via a parameter B for squeezing which when looking at a squeeze parameter r, for which $0 \leq r \leq \infty$, then (14) has, instead of $B(\eta_b) \equiv \omega_b/2$

$$B|_{\eta} \xrightarrow{\eta \neq \eta_b} B(\omega, \eta \neq \eta_b) \equiv \frac{i}{2} \cdot \frac{(\mu/a(\eta))'}{(\mu/a(\eta))} \equiv \frac{\omega}{2} \cdot \frac{\cosh r + [\exp(2i\mathcal{G})] \cdot \sinh r}{\cosh r - [\exp(2i\mathcal{G})] \cdot \sinh r} \quad (15)$$

Taking Grishchuck’s formalism literally, a state for a graviton/ GW is not affected by squeezing when we are looking at an initial frequency, so that $\omega \equiv \omega_b$ initially corresponds to a non squeezed state which may have coherence, but then right afterwards, if $\omega \neq \omega_b$ which appears to occur whenever the time evolution,

$$\eta \neq \eta_b \Rightarrow \omega \neq \omega_b \Rightarrow B(\omega, \eta \neq \eta_b) \equiv \frac{i}{2} \cdot \frac{(\mu/a(\eta))'}{(\mu/a(\eta))} \neq \frac{\omega_b}{2} \quad \text{A reasonable research task would be to}$$

determine, whether or not $B(\omega, \eta \neq \eta_b) \neq \frac{\omega_b}{2}$ would correspond to a vacuum state being initially formed

right after the point of nucleation, with $\omega \equiv \omega_b$ at time $\eta \equiv \eta_b$ with an initial cosmological time some order of magnitude of a Planck interval of time $t \approx t_{planck} \propto 10^{-44}$ seconds

Open questions. Turbulence in initial GW production and how to model it ? Either classically or quantum mechanically

What happens if there is a switch over from an initially uncompressed state, to one which has compression? Several things could happen. First of all, one may be able to see colliding plane wave representations of GW, i.e the geometry of the colliding wave space time becomes amendable to analysis, as was presented by Vladimir Belinski, and Enric Venrauger (2001) in their book on Gravitational solitons, starting on page 202. In particular, their equation (7.60) has parameters which represent gravitational shock waves in collision, followed by trailing gravitational radiation. If one believes that relic GW processes can be largely preserved in the onset of the big bang in a ‘frozen’ profile then the interactive region for generation of GW

signals from GW shock waves in collision could account for the datum represented by Fangyu Li et al (2009) as far as the alleged random back ground as far as GW processes. Secondly is the issue which Bojowald (2008) talked to the author, Beckwith, about in the 12 Marcel Grossman conference , mainly what is known, and what is not know about the geometry of space time , presumably in the aftermath of the big bounce (LQG). Bojowald’s (2008) paper leaves the relative degree of squeezing mandated by the big bounce as a ‘to be solved’ datum .

For the sake of comparison, Furthermore, Abhay Ashtekar (2006) wrote a simple treatment of the Bounce causing Wheeler De Witt equation along the lines of , for $\rho_* \approx const \cdot (1/8\pi G\Delta)$ as a critical density, and Δ the eignvalue of a minimum area operator. Small values of Δ imply that gravity is a repulsive force, leading to a bounce effect.

$$\left(\frac{\dot{a}}{a}\right)^2 \equiv \frac{8\pi G}{3} \cdot \rho \cdot (1 - (\rho/\rho_*)) + H.O.T. \quad (16)$$

Furthermore, Bojowald (2008) specified a criteria as to how to use an updated version of Δ and $\rho_* \approx const \cdot (1/8\pi G\Delta)$ in his GRG manuscript on what could constitute grounds for the existence of generalized squeezed initial (graviton ?) states. Bojowald (2008) was referring to the existence of squeezed states, as either being necessarily, or NOT necessarily a consequence of the quantum bounce. As Bojowald (2008) wrote it up, in both his equation (26) which has a quantum Hamiltonian $\langle \hat{V} \rangle \approx H$,with

$$\left. \frac{d\langle \hat{V} \rangle}{d\phi} \right|_{\phi \approx 0} \xrightarrow{\text{existence-of-un-squeezed-states} \Leftrightarrow \phi \approx 0} 0 \quad (17)$$

, and \hat{V} is a ‘volume’ operator where the ‘ volume’ is set as V , Note also, that Bojowald has , in his initial Friedman equation, density values $\rho \equiv \frac{H_{matter}(a)}{a^3}$, so that when the Friedman equation is quantized, with an initial internal time given by ϕ , with ϕ becoming a more general evolution of state variable than ‘internal time’. If so, Bojowald (2008) writes, when there are squeezed states

$$\left. \frac{d\langle \hat{V} \rangle}{d\phi} \right|_{\phi \neq 0} \xrightarrow{\text{existence-of-squeezed-states}} N(\text{value}) \neq 0 \quad (18)$$

for his equation (26), which is incidently when links to classical behavior break down , and when the bounce from a universe contracting goes to an expanding present universe,. Bojowald also writes that if one is looking at an isotropic universe, that as the large matter ‘H’ increases, that in certain cases, one observes more classical behavior, and a reduction in the strength of a quantum bounce.. Bojowalds states that “Especially the role of squeezed states is highlighted. The presence of a bounce is proven for uncorrelated states, but as squeezing is a dynamical property and may change in time”I claim that what Bojowald (2008) is leading up to, is specifying a parameter space in initial conditions which one may be able to do a semi classical analysis of the sort referenced by Vladimir Belinski, and Enric Venrauger (2001) in their book on Gravitational solitons, starting on page 202 of their text.. As stated earlier, their equation (7.60) has parameters which represent gravitational shock waves in collision, followed by trailing gravitational radiation. Not only that, but initial un squeezed states may be, in part represented/ presentable as due to the worm hole analysis of initially introduced from a prior universe, to today’s universe by the WdM pseudo time representation of an initial vacuum state, as has been brought up by Beckwith, in 2008, 2009,.Last, but not least, would be to also examine, from first principles, what Christian Corda raised as a distinct possibility Namely using “ investigation of the transverse effect of gravitational waves (GW's) could constitute a further tool to discriminate among several relativistic theories of gravity on the ground.

After a review of the TT gauge, the transverse effect of GW's arising by standard general relativity (called Einstein's GW's in this paper) is reanalyzed with a different choice of coordinates.” . I.e. using transverse effects as another further tool to distinguish on the foundations of what Li et al (2009) listed as random background for the processes in which relic GW are generate in early space time conditions

Table 1: magnitude, sources, and top frequency values for HFGW (from Li et al. 2009)

Sources	Amplitude	frequency	Characteristics
HFGW in Quintessence inflationary models	$h_{rms} \sim 10^{-30} - 10^{-32} / \sqrt{Hz}$	$\nu \sim 10^9 - 10^{10} Hz$	Random background
HFGW in some string theory scenarios	$h_{rms} \sim 10^{-30} - 10^{-34} / \sqrt{Hz}$	$\nu \sim 10^8 - 10^{11} Hz$	Random background
HFGW in some string theory scenarios	$h_{rms} \sim 10^{-30} - 10^{-34} / \sqrt{Hz}$	$\nu \sim 10^8 - 10^{11} Hz$	Random background
Solar Plasma	$h_{rms} \sim 10^{-39} / \sqrt{Hz}$	$\nu \sim 10^{15} Hz$	On the Earth

How would DM be influenced by gravitons, in 4 dimensions

We will also discuss the inter relationship of structure of DM, with challenges to Gaussianity. The formula as given by

$$\delta \equiv - \left[\frac{3}{2} \cdot \Omega_m \cdot H^2 \right]^{-1} \cdot \nabla^2 \Phi \quad (19)$$

will be gone into. The variation, so alluded to which we will link to a statement about the relative contribution of Gaussianity, via looking at the gravitational potential

$$\Phi \equiv \Phi_L + f_{NL} \cdot \left[\Phi_L^2 - \langle \Phi_L^2 \rangle \right] + g_{NL} \cdot \Phi_L^3 \quad (20)$$

Here the expression $f_{NL} =$ variations from Gaussianity, while the statements as to what contributes, or does not contribute will be stated in our presentation. Furthermore, Φ_L is a linear Gaussian potential, and the over all gravitational potential is altered by inputs from the term, presented, f_{NL} . The author discussed inputs into variations from Gaussianity, which were admittedly done from a highly theoretical perspective with Sabino Matarre, on July 10, with his contributions to non Guassianity being constricted to a reported range of $-4 < f_{NL} < 80$, as given to Matarre, by Senatore, et al, 2009.. . Needless to state, though, dealing with what we can and cannot measure, what is ascertained as far as DM, via a density profile variation needs to have it reconciled with DM detection values

$$\sigma_{DM-detection} \leq 3 \times 10^{-8} \text{ pb (pico barns)} \quad (21)$$

It is note worthy to note that the question of DM/ KK gravitons, and also the mass of the graviton not only has relevance to whether or not, higher dimensions are necessary/ advisable in space time models, but also may be relevant to if massive gravitons may solve / partly fulfill the DE puzzle. To whit, \ KK gravitons would have a combined sum of Bessel equations as a wave functional representation. In fact V. A Rubasov (2009) writes that KK graviton representation as, after using the following

normalization $\int \frac{dz}{a(z)} \cdot [h_m(z) \cdot h_{\tilde{m}}(z)] \equiv \delta(m - \tilde{m})$, where J_1, J_2, N_1, N_2 are different forms of

Bessel functions, to obtain the KK graviton/ DM candidate representation along RS dS brane world

$$h_m(z) = \sqrt{m/k} \cdot \frac{J_1(m/k) \cdot N_2([m/k] \cdot \exp(k \cdot z)) - N_1(m/k) \cdot J_2([m/k] \cdot \exp(k \cdot z))}{\sqrt{[J_1(m/k)]^2 + [N_1(m/k)]^2}} \quad (22)$$

This can be conflated with Marcio E. S. Alves, Oswaldo D. Miranda, Jose C. N. de Araujo's results arguing that non zero graviton mass may lead to acceleration of our present universe, in a manner usually conflated with DE, i.e. their graviton mass would be about

$m_{graviton} (4 - Dim GR) \sim 10^{-48} \times 10^{-5} eV \sim 10^{-65}$ grams, leading to a possible explanation for when the universe accelerated, i.e. the de-acceleration parameter, due to changes in the scale factor, written as

$$q = -\frac{\ddot{a}a}{\dot{a}^2} \quad (23)$$

In the case of working with a simpler version of the Friedman equation with no graviton mass, but with pressure and density factored in, we can obtain

$$\frac{\ddot{a}}{a} = \frac{4\pi G}{3} \cdot [(-3p/c^2) - \rho] \quad (24)$$

This will lead to a very simple de celebration parameter value of

$$q = -\frac{\ddot{a}a}{\dot{a}^2} = \left(\frac{4\pi G}{3c^2 H^2} \right) \cdot [3p + \rho] \quad (25)$$

Needless to say, if one has a graviton mass $m_{graviton} \neq 0$, then (15) changes. Using a modification of GR, with scale factor evolution of , with non zero graviton mass terms added in to obtain

$$\left(\frac{\dot{a}}{a} \right)^2 + \frac{m_g^2 c^4}{2\hbar^2} \cdot (1 - a^2) \equiv \frac{8\pi G}{3c^2} \cdot \rho \quad (26)$$

and

$$\frac{\ddot{a}}{a} + .5 \cdot \left(\frac{\dot{a}^2}{a^2} \right) + \frac{m_g^2 c^4}{4\hbar^2} a^2 \cdot (a^2 - 1) = \frac{8\pi G}{3c^2} \cdot p . \quad (27)$$

For the matter dominated era, it is important to note that the R.H.S. of (27) is zero. This leads to (23) and (25) having increasingly positive acceleration values as would be definitely be given for masses of

$m_{graviton} (4 - Dim GR) \sim 10^{-48} \times 10^{-5} eV \sim 10^{65}$ grams for red shift values $z \sim .3$ for (25) just becoming > 0 to maximum values of (25) today, with $z = 0$, all at mass of the order of 10^{65} grams.

$r_c = \frac{M_P^2}{2M_{(5)}^3}$ and , often $\epsilon = 1$ and r_c is usually though of as the separation between branes. I.e. if

$r_c \rightarrow \infty$, we recover the usual first Friedman equation. For now we write the first Friedman equation for a brane system as.

$$\frac{\dot{a}}{a} \equiv H = \frac{\epsilon}{2r_c} + \sqrt{\frac{8\pi G\rho}{3} + \frac{1}{4r_c^2}} \quad (27a)$$

As can be related to, if we wish to look at string theory versions of the FRW equation , in Friedman-Roberson – Walker metric space, we can do the following de composition , with different limiting values of the mass, and other expressions, e.g. as a function of an existing cosmological constant

$$\left(\frac{\dot{a}}{a} \right)^2 = \frac{\rho_{Total}}{3M_{Planck}^2} - \frac{k}{a^2} + \frac{\Lambda}{3} \quad (28)$$

As well as

$$\left(\frac{\ddot{a}}{a} \right) = -\frac{(\rho_{Total} + 3p_{Total})}{6M_{Planck}^2} + \frac{\Lambda}{3} \quad (29)$$

Not only this, if looking at the brane theory Friedman equations as presented by / for Randall Sundrum theory, it would be prudent working with

$$\dot{a}^2 = \left[\left(\frac{\rho}{3M_4^2} + \frac{\Lambda_4}{3} + \frac{\rho^2}{36M_{Planck}^2} \right) a^2 - \kappa + \frac{C}{a^2} \right] \quad (30)$$

For the purpose of Randal Sundrum brane worlds, (25) is what will be differentiated with respect to $d/d\tau$, and then terms from (1.5) will be used, and put into a derivable equation which will be for a RS

brane world version of $q = -\frac{\ddot{a}a}{\dot{a}^2}$.. Note that Roy Maartens has written as of 2004 that KK modes (graviton) satisfy a 4 Dimensional Klein – Gordon equation, with an effective 4 dim mass,

$m_n(Graviton) = \frac{n}{L}$, with $m_0(Graviton) = 0$, and L as the stated ‘dimensional value’ of higher

dimensions. The value $m_0(Graviton) \sim 10^{-65} - 10^{-60}$ gram in value picked is very small, but

ALMOST zero. Grossing has shown how the Schrodinger and Klein Gordon equations can be derived from classical lagrangians, i.e. using a version of the relativistic Hamilton-Jacobi- Bohm equation, with a wave functional $\psi \sim \exp(-iS/\hbar)$, with S the action, so as to obtain working values of for a tier of

purported masses of a graviton from the equation , for 4 D of $\left[g^{\alpha\beta} \partial_\alpha \partial_\beta \xrightarrow{FLAT-SPACE} \nabla^2 - \partial_\tau^2 \right]$, and

$\left[\nabla^2 - \partial_\tau^2 \right] \cdot \psi_n = m_n^2(graviton) \cdot \psi_n$ If one is adding , instead the small mass of

$m_n(Graviton) = \frac{n}{L} + 10^{-65}$ grams, with $m_0(Graviton) \approx 10^{-65}$ grams, then the problem being

worked with is a source term problem of the form given by Peskins as of the type

$$\psi_n(x) \equiv \int d^3 p \cdot \frac{1}{(2\pi)^3} \cdot \frac{1}{\sqrt{2E_p}} \cdot \left\{ \left(a_p + \frac{i}{\sqrt{2E_p}} \cdot FT(m_0(graviton)) \right) \exp(-ipx) + H.C. \right\} \quad (31)$$

This is, using the language V.A. Rubakob (2009) put up equivalent to writing, using (1.3)

$$\psi_m(x) \approx h_m(x) + \int d^3 p \cdot \frac{1}{(2\pi)^3} \cdot \left(\frac{1}{\sqrt{2E_p}} \right)^2 \cdot \{ (i \cdot FT(m_0(graviton))) \exp(-ipx) + H.C. \} \quad (32)$$

I.e. how to intepret the quantity $FT(m_0(graviton))$ being the issue which will be covered in this document. If $m_0(graviton)$ is a constant, then the expression (32) has delta functions. This datum may especially show up about modification of the typical galaxy models, as follows

Controversies of DM/ DE applications to cosmology. How HFGW may resolve them.

The following is meant as a travelogue as to current problems in cosmology which will require significant revision of our models. Exhibit A as to what to consider is The cosmic void hypothesis?. See Timothy Clifton, Pedro G. Ferreira and Kate Land . I.e. Clifton raises the following question- can HFGW and detectors permit cosmologist to get to the bottom of this ? “Solving Einstein’s equations for an averaged matter distribution is NOT the same as solving for the real matter distribution and then averaging the resultant geometry”(“We average, then solve when in effect we should solve, then average”) .Next, let us look at a recently emerging conundrum of DM feeding into the structure of new galaxies and their far earlier than expected development, i.e. 5 billion years after the big bang. Galaxy formation issues.... Hierarchical Galaxy Formation theory at a glance usually proceeds as follows. I.e. what happens when the following diagram of simple addition of new structure no longer holds ? This is very significant, since when the significant formation of galaxies occurs, as of about $z \sim .2$ is before the turn up in the expansion rate for the universe, which will be referenced as of occurring about $z \sim .5 - .55$. What do we do if, as an example, find that the initial start of galaxy formation occurred five billion years ago, at, say $z \sim .5$. What could cause the earlier clumping? Several scenarios which will be investigated. First of all, note the formula of variation of DM density which exists has, among other things a Hubble parameter H , and also the 2nd derivative of the gravitational potential $\nabla^2 \Phi$, where ρ_0, a_0 are today’s values for density and

‘distance’ .Note that if $H^2 = \left(\frac{\dot{a}}{a} \right)^2 \xrightarrow[\Lambda_4 \rightarrow 0]{\kappa \rightarrow 0} \left[\frac{\rho}{3M_4^2} + \frac{\rho^2}{36M_{Planck}^2} + \frac{C}{a^4} \right]$, and $1+z = a_0/a$ and

$\rho \rightarrow \rho(z) \equiv \rho_0 \cdot (1+z)^3 - \left[\frac{m_g}{8\pi G} \right] \cdot \left(\frac{a_0^4}{14 \cdot (1+z)^4} + \frac{2a_0^2}{5 \cdot (1+z)^2} - \frac{1}{2} \right)$, , then the contribution of large z , i.e. large contributions from red shift, that a significant early contributions will be for non zero contributions from $1/\rho^\beta$ terms , for [**large number**] $> \beta \geq 1$ in the DM density variation , and the situations in which this is the case.




Figure 1. A schematic representation of a halo merging history 'tree'.

Figure 1. I.e. how we obtain from the ‘bottom up’ development of galactic super structure.

What is actually observed, contradicts this halo emerging history ‘tree’, i.e. Although this ‘story’ for DM seems to be well established. i.e. Just ONE little problem: DM appears to be fattening up young galaxies, allowing for far-earlier-than-expected creation of early galaxies. “A clutch of massive galaxies that seem to be almost fully-formed just 5 billion years after the big bang challenge models that suggest galaxies can only form slowly. Tendrils of dark matter that fed the young galaxies on gas could be to blame (NASA/CXC/ESO/P Rosati et al)”

<http://www.newscientist.com/article/dn16912-overweight-galaxies-forcefed-by-dark-matter-tendrils.html>

I.e. considering the following. The relative imprecision of graviton measurement, can be given as follows. This is a measurement in particle physics, and if the KK graviton is linkable to DM, it means that we will have to have very good ways to test for production rates, as will be argued later. The following below is a representation of the KK tower model for gravitons, with the zeroth KK mode being approximately the 4 dimensional graviton.

Kaluza Klein modes in detector simulations for / as a DM candidate.




Figure 2a:

Figure 2a. **Example:** Number of Events in $e^+e^- \rightarrow \mu^+\mu^-$ For a conventional braneworld model with a single curved extra dimension of size $\sim 10^{-17}$ cm. Numbers range from 10^4 to about 10^8 for the number of events in scattering. First peak is for KK zero mode, a.k.a. the standard Z. boson, ending with the 4th peak for the 3rd KK mode,

Production rate for $e^+e^- \rightarrow \gamma + \text{Graviton}$




Figure 2b: Example: Production of Graviton Kaluza-Klein modes in flat extra dimensions, probes gravity at distances of ~ 10 to the -18 power cm. The LHS of the graph have production rates ranging from a low point of 10^4 at about 600 GeV, to values of about 10^6 at about 1000 GeV. Understanding the KK gravitons as a DM candidate may permit us to understand how DM and DE are inter related. See as given below. The discussion of such will involve coherent state of gravitons as contributors to GW.

2. Creating an analysis of how graviton mass, assuming branes, can influence expansion of the universe

Following development of *(1.13) as mentioned above, with inputs from Friedman eqns. To do this,, the following normalizations will be used, i.e. $\hbar = c = 1$, so then

$$q = A1 + A2 + A3 + A4 \quad (33)$$

Where

$$A1 = \frac{C}{a^3} \cdot \left[\frac{1}{\sqrt{\frac{C}{a^4} - \frac{\kappa}{a^2} + \left(\frac{\rho}{3M_4^2} + \frac{\Lambda_4}{3} + \frac{1}{36} \cdot \frac{\rho^2}{M_p^6} \right)}} \right] \quad (34)$$

$$A2 = -\left(\frac{\rho}{3M_4^2} + \frac{\Lambda_4}{3} + \frac{1}{36} \cdot \frac{\rho^2}{M_p^6}\right) \left/ \left[\frac{C}{a^4} - \frac{\kappa}{a^2} + \left(\frac{\rho}{3M_4^2} + \frac{\Lambda_4}{3} + \frac{1}{36} \cdot \frac{\rho^2}{M_p^6}\right) \right] \right. \quad (35)$$

$$A3 = -\frac{1}{2} \cdot \left[\frac{(d\rho/d\tau)}{3M_4^2} + \frac{(d\Lambda_4/d\tau)}{3} + \frac{1}{18} \cdot \frac{\rho \cdot (d\rho/d\tau)}{M_p^6} \right] \left/ \left[\frac{C}{a^4} - \frac{\kappa}{a^2} + \left(\frac{\rho}{3M_4^2} + \frac{\Lambda_4}{3} + \frac{1}{36} \cdot \frac{\rho^2}{M_p^6}\right) \right]^{3/2} \right. \quad (36)$$

$$A4 = \frac{\kappa}{a^3} \cdot \left[\frac{(da/d\tau)}{3} \right] \left/ \left[\frac{C}{a^4} - \frac{\kappa}{a^2} + \left(\frac{\rho}{3M_4^2} + \frac{\Lambda_4}{3} + \frac{1}{36} \cdot \frac{\rho^2}{M_p^6}\right) \right]^{3/2} \right. \quad (37)$$

Furthermore, if we are using density according to whether or not 4 dimensional graviton mass is used, then

$$\rho \equiv \rho_0 \cdot \left(\frac{a_0}{a}\right)^3 - \left[\frac{m_g c^6}{8\pi G \hbar^2} \right] \cdot \left(\frac{a^4}{14} + \frac{2a^2}{5} - \frac{1}{2}\right) \quad (38)$$

So, then one can look at $d\rho/d\tau$ obtaining

$$d\rho/d\tau = -\left(\frac{\dot{a}}{a}\right) \cdot \left[3 \cdot \rho_0 \cdot \left(\frac{a_0}{a}\right)^3 + 4 \cdot \left(\frac{a^4}{14} + \frac{a^2}{5}\right) \cdot \left(\frac{m_g c^6}{8\pi G \hbar^2}\right) \right] \quad (39)$$

Also, then , set the curvature equal to zero. i.e. $\kappa = 0$, so then $A4 = 0$, and

Pick, here, $\rho \equiv \rho_0 \cdot \left(\frac{a_0}{a}\right)^3 - \left[\frac{m_g}{8\pi G} \right] \cdot \left(\frac{a^4}{14} + \frac{2a^2}{5} - \frac{1}{2}\right)$, after $\hbar = c = 1$, and also set

$$\Phi(\rho, a, C) = \frac{C}{a^4} + \left(\frac{\rho}{3M_4^2} + \frac{1}{36} \cdot \frac{\rho^2}{M_p^6}\right) \quad (40)$$

For what it is worth, the above can have the shift to red shift put in by the following substitution. I.e. use $1+z = a_0/a$.. Assume also that C is the dark radiation term which in the brane version of the Friedman equation scales as a^{-4} and has no relationship to the speed of light. a_0 is the value of the scale factor in the present era, when red shift $z=0$, and $a \equiv a(\tau)$ in the past era, where τ is an interval of time after the onset of the big bang. $(a_0/a)^3 = (1+z)^3$, and $a \equiv a_0/(1+z)$, Then

$$\rho(z) \equiv \rho_0 \cdot (1+z)^3 - \left[\frac{m_g}{8\pi G} \right] \cdot \left(\frac{a_0^4}{14 \cdot (1+z)^4} + \frac{2a_0^2}{5 \cdot (1+z)^2} - \frac{1}{2}\right) \quad (41)$$

$$A1(z) \cong \frac{C \cdot (1+z)^3}{a_0^3} \cdot \left[1/\sqrt{\Phi(\rho(z), a_0/(1+z), C)} \right] \quad (42)$$

$$A2(z) \cong -\left(\frac{\rho(z)}{3M_4^2} + \frac{1}{36} \cdot \frac{\rho(z)^2}{M_p^6}\right) \left/ \left[\Phi(\rho(z), a_0/(1+z), C) \right] \right. \quad (43)$$

$$A3(z) \cong \frac{1}{2} \left(\left[\frac{1}{3M_4^2} + \frac{1}{18} \cdot \frac{\rho(z)}{M_p^6} \right] \left/ \left[\Phi(\rho(z), a_0/(1+z), C) \right]^{1/2} \right. \right) \cdot \quad (44)$$

$$\left[3 \cdot \rho_0 \cdot (1+z)^3 + 4 \cdot \left(\frac{a_0^4/(1+z)^4}{14} + \frac{a_0^2/(1+z)^2}{5}\right) \cdot \left(\frac{m_g}{8\pi G}\right) \right]$$

$$\Phi(\rho(z), a_0/(1+z), C) = \frac{C \cdot (1+z)^4}{a_0^4} + \left(\frac{\rho(z)}{3M_4^2} + \frac{1}{36} \cdot \frac{\rho(z)^2}{M_p^6} \right) \quad (45)$$

So, for $4 < z \leq 0$, i.e. not for the range, say $z \sim 1100$ 380 thousand years after the big bang, it would be possible to model, here

$$q(z) = A1(z) + A2(z) + A3(z) \quad (46)$$

Easy to see though, that to first order, $q(z) = A1(z) + A2(z) + A3(z)$ would be enormous when $z \sim 1100$, and also that for $Z=0$, $q(0) = A1(0) + A2(0) + A3(0) > 0$. Negative values for Eqn. (46) appear probable at about $z \sim 1.5$, The positive value conditions rely upon, the C dark radiation term,

And here are the results! Assume X is red shift, Z . $q(X)$ is De - Celebration . Here De celebration parameter due to small $m_{graviton} \propto 10^{-65}$ grams, with one additional dimension added




Figure 4 a: re duplication of basic results of Marcio E. S. Alves, Oswaldo D. Miranda, Jose C. N. de Araujo, 2009, using their parameter values, with an additional term of C for Dark flow added, corresponding to one KK additional dimensions




Figure 4 b: re duplication of basic results of Marcio E. S. Alves, Oswaldo D. Miranda, Jose C. N. de Araujo, 2009, using their parameter values, with an additional term of C for 'Dark flow' added, corresponding to one KK additional dimensions. Results show asymptotic 'collapse' of de celebration parameter as one comes away from the red shift $Z=1100$ of the CMBR 'turn on' regime for de coupling of photons from 'matter', in end of 'dark ages' Figures 4a, and 4b suggest that additional dimensions are permissible. They do not state that the initial states of GW/ initial vacuum states have to form explicitly due to either quantum or semi classical processes.

CONCLUSION: Investigating fundamentals of generation of GW in relic conditions. Time to re set GW physics to empirical foundations.

The author, Beckwith, is fully aware of how unpopular his conclusions will be with respect to current string theory proponents, who have managed to move string theory from its initial Calabi Yau compact higher

dimensions focus, as Giovannini, and others used successfully (1995) to argue for almost unlimited higher frequencies as to relic GW, to the unlimited higher dimensions specified by Arkani-Hamid, and others. It is time for the GW and the GR community to cease being intimidated by some of these practitioners and to re focus and re set the discussion on relic conditions to becoming once again an empirical science, one whose focus should be data analysis, and not post modern physics. The author, also views as potentially revolutionary the implications as argued by t'Hooft, Corda, and others that Gravity is essentially classical in its origins. A datum which can be investigated by determining if or not Vladimir Belunski , and Enric Vergaguer are right about their interaction region for shock waves, as could be modeled for initial conditions. I.e. this modeling of Vladimir Belunski , and Enric Vergaguer 's modeling of the collision of GW is under way right now by the author, and the results will be mapped onto possible relic GW spectra, once numerical protocol for doing so are fully developed by the author, Beckwith. The final pay off, of moving beyond post modern physics, and re-setting the discussion back to laboratory science, will be in investigating a supposition t'Hooft advanced as to Quantum mechanics, which has never been satisfactorily investigated. The reconstruction of generation of GW in initial conditions may be allowing us **to illustrate 't Hooft's proposal to reconstruct quantum mechanics** as an emergent theory. It does not get any better than this, in terms of learning reality as we know it. The author, Beckwith, will in a subsequent publication, elaborate upon why early generation of GW could be the **perfect template** as to investigating T'Hooft's supposition in proper detail, and what that could mean with respect to physics.

BIBLIOGRAPHY

- B. Allen, E. Flanagan, and M.A. Papa, Is the squeezing of relic gravitational waves produced by inflation detectable? Phys.Rev.D61:024024,2000., http://arxiv.org/PS_cache/gr-qc/pdf/9906/9906054v2.pdf ...
- Marcia E. S. Alves, Oswaldo D. Miranda and Jose C. N. de Araujo," Can Massive Gravitons be an Alternative to Dark Energy?", arXiv:0907.5190 (July 2009)
- Andrić, I, Jonke , L. and Jurman , D."Solitons and giants in matrix models", JHEP12(2006)006
- Abhay Ashtekar (2006), solvaynet.pdf
- Daniel Babich, Paolo Creminelli, Matias Zaldarriaga,"The shape of non – Gaussianities", CAP 0408:009,2004; <http://arxiv.org/abs/astro-ph/0405356>
- Baker, R.M.L, Stephenson, G. V. , and Li Fangyu (2008), "Proposed Ultra-High Sensitivity HFGW Detector,"in the proceedings of Space Technology and Applications International Forum (STAIF-2008), edited by M.S. El-Genk, American Institute of Physics Conference Proceedings, Melville , NY 969, pp. 1045-54 <http://www.gravwave.com/docs/ProposedUltraHighSensitivityHFGWDetector2005-15-08.pdf>
- Carlos Barcelo, Luis J. Garay,"Quantum fields in Anti-De Sitter Wordholes", http://arxiv.org/PS_cache/gr-qc/pdf/9703/9703028v2.pdf
- James M. Bardeen * Gauge-invariant cosmological perturbations', Phys. Rev. D 22, 1882 - 1905 (1980); Nicola Bartolo;
- Barvinsky, A. "Quantum geometrodynamics: The Wheeler-DeWitt equations for the wave function of the universe"Physics letters B, Volume 175, issue 4, August 14, 1986, pp 401-
- Baumann, Daniel, Ichiki, Kiyotomo, Steinhardt, Paul J. Takahashi , Keitaro ,," Gravitational Wave Spectrum Induced by Primordial Scalar Perturbations", arXiv:hep-th/0703290
- Beckwith,A.W. "Entropy growth in the early universe and confirmation of initial big bang conditions (Wheeler De Witt eqn. results vs. String theory ?)", <http://vixra.org/abs/0908.0109>
- Beckwith, A.W. "Instanton formation of vacuum energy via the Reissner-Nordstrom geometry of a wormhole bridge between a prior to our present universe.", arXiv:0710.3788, October 2007
- Beckwith, A.W., "Implications for the Cosmological Landscape: Can Thermal Inputs from a Prior Universe Account for Relic Graviton Production?" AIP Conference Proceedings 969, (2008), pp. 1091-1102
- Beckwith, A. W. "Relic High Frequency Gravitational waves from the Big Bang, and How to Detect Them", <http://arxiv.org/abs/0809.1454> (2009), AIP Conf.Proc.1103:571-581,2009 (SPESIF)
- Martin Bojowald,(2008), "Quantum nature of cosmological bounces "General Relativity and Gravitation, pp 2659-2683, Volume 40, Number 12, December, 2008; <http://arxiv.org/abs/0801.4001>
- Vladimir Belunski, and Enric Verdaguer ,," Gravitational Solitons", Cambridge Monographs on Mathematical physics, Cambridge, UK, (2001)
- Brett Bolen, Luca Bombelli, Alejandro Corichi "Semiclassical States in Quantum Cosmology: Bianchi I Coherent States", Class.Quant.Grav. 21 (2004) 4087-4106; <http://arxiv.org/abs/gr-qc/0404004>
- R. Brustein, M. Gasperini, M. Giovannini, and G. Veneziano,"Relic Gravitational waves from String Cosmology", <http://cdsweb.cern.ch/record/284281/files/9507017.pdf>; Physics Letters B 361(1995) 45-51
- Buonanno, A.; Ungarelli, C."Gravitons and String Pre Big Bang cosmology", String Theory and Fundamental Interactions, Lecture Notes in Physics, Volume 737. ISBN 978-3-540-74232-6. Springer-Verlag Berlin Heidelberg, 2008, p. 845-851

Camp, J. , and Cornish, N., “Gravitational wave Astronomy”,pp 525-577, as part of “ Annual Review of Nuclear and Particle Science”, Volume 54, with Kayser, B., Holstein, B., and Jawahery, A. as editors. Menlo Park, California, 2004

Capozziello, S., Feoli, A., Lambiase, G., Papini, G., ‘ Thin shell quantization in Weyl Space – time’, Physics Letters A, 273 (2000), pp 25-30

Marco Cavaglià, “Quantum Electromagnetic Wormholes and Geometrical Description of the Electric Charge”, Phys.Rev. D50 (1994) 5087-5092; <http://arxiv.org/abs/gr-qc/9407030>

Ceresole, A., D’ Auria, R., Ferrara, S., and van Poeyen, A., “ Duality transformations in super symmetric Yang-Mills theories, coupled to supergravity,” Nucl. Phys. B444 (1995), pp 92-124, hep-th/ 9502072

P. Chingabam, C. Park,”Statistical nature of Non Gaussianity from cubic order Primordial perturbations : CMB map simulations and genus statistics” http://arxiv.org/PS_cache/arxiv/pdf/0908/0908.1696v1.pdf

Dah-Wei Chiou, Li-Fang Li,” Loop quantum cosmology with higher order holonomy corrections”, Phys. Rev. D 80, 043512 (2009); <http://arxiv.org/abs/0907.0640>

Christian Corda,”Analysis of the transverse effects of Einsteins Gravitational waves”, International Journal of Modern Physics A, Volume 22, Issue 26, pp. 4859-4881 (2007); <http://adsabs.harvard.edu/abs/2007arXiv0707.1943C>

Christian Corda,” Primordial production of massive relic gravitational waves from a weak modification of General Relativity”, Astropart.Phys.30:209-215,2008, <http://arxiv.org/abs/0812.0483>

Crowell, L., “Quantum Fluctuations of Space Time “, World Scientific Series in Contemporary Chemical Physics, Vol 25, World Scientific, PTE, LTD, 2005, Singapore

Dadhich, N., “Derivation of the Raychaudhuri Equation”, <http://arxiv.org/abs/gr-qc/0511123>,

De Laurentis, Mariafelicia, and Capozziello, Salvatore,,” Stochastic Background of Relic Scalar Gravitational Waves tuned by Extended Gravity”, <http://arxiv.org/abs/0906.3689>

Rainer Dick,” Heterotic string dark matter from the graviton multiplet”, Physics Letters B, Volume 510, Issues 1-4, 21 June 2001, Pages 10-16

Dowker, H.F. “Causal sets and the deep structure of spacetime”, arXIV gr-qc/0508109v1 26 Aug 2005; in 100 Years of Relativity Space-Time Structure: Einstein and Beyond, edited by Ashtekar, A., World Press Scientific, Singapore, 2005

Durrer, R., “ Cosmological Perturbation theory”, From the book “Physics of the Early Universe”, in Lecture Notes in Physics, 653”, with Papantonopoulos, E., as editor and printed by Springer- Verlag 2004

Ruth Durrer, Massimiliano Rinaldi , “Graviton production in non-inflationary cosmology”, Phys.Rev.D79:063507,2009, <http://arxiv.org/abs/0901.0650>

Enqvist, K., Mazumdar, A., Perez--Lorenzana, A., “ Dumping inflaton energy out of this world”,Phys Rev D, Volume 70, 103508

Feng, Jonathan L. and Kumar, Jason “Dark-Matter Particles without Weak-Scale Masses or Weak Interactions.” Physical Review Letters (2008). Available online: <http://link.aps.org/doi/10.1103/PhysRevLett.101.231301>.

Feinstein,A , Vazquez-Mozo, M.A, “M-theory resolution of four-dimensional cosmological singularities via U-duality” , <http://arxiv.org/abs/hep-th/9906006> , Nucl.Phys. B568 (2000) 405-420

L.H. Ford,” Gravitons and Lightcone Fluctuations”, 10.1103/PhysRevD.51.1692, <http://arxiv.org/abs/gr-qc/9410047>

Luis J. Garay ,” Quantum state of wormholes and path integral” Phys. Rev. D 44, 1059 – 1066;

Giovannini, Massimo, ” A Primer on the Physics of the Cosmic Microwave Background”, World Press Scientific, 2008, Singapore, Republic of Singapore, (2008)

Glinka, L. “Quantum Information from Graviton-Matter Gas ”, Sigma 3, (2007), 087, 13 pages

Roy Glauber, “Coherent and Incoherent States of the Radiation Field”, Phys. Rev. 131, 2766 - 2788 (1963)

Leonid P Grishchuk”The problem of relic gravitational radiation”, Sov. Phys. Usp. 23 705-706

L P Grishchuk and Yu Sidorov,” On the quantum state of relic gravitons”, 1989 Class. Quantum Grav. 6 L161-1

L.P. Grishchuk,” Quantum Effects In Cosmology” Class.Quant.Grav. 10 (1993) 2449-2478, <http://arxiv.org/abs/gr-qc/9302036>

Lenoid P, Grushchuk,”The Detectability of Relic (Squeezed) Gravitational Waves by Laser Interferometers”, <http://arxiv.org/abs/gr-qc/9810055> , 1998

Grishchuk,L. “Discovering Relic Gravitational Waves in Cosmic Microwave Background Radiation”, <http://arxiv.org/abs/0707.3319> (2008)

Hartle, J., and Hawking, S.W., "Wave function of the universe", Phys. Rev. D28, 2960 (1983)

Bert Janssen, Yolanda Lozano, “A Microscopical Description of Giant Gravitons” Nucl.Phys. B658 (2003) 281-299, , <http://arxiv.org/abs/hep-th/0207199>

Kahniashvili ,Tina”Relic Gravitational Waves as a Test of the Early Universe”,arXiv:0705.1733 [astro-ph], (2007)

Claus Kiefer,” Quantum gravity”, (International Series of Monographs on Physics) (Hardcover),Oxford Science Publications, Oxford University Press, 2007

Kolb, E., and Turner, S. “The Early Universe”, Westview Press, Chicago, USA, 1994

Lawrence Krauss (1996); “Old Galaxies at high Red shifts, and the Cosmological constant”, THE ASTROPHYSICAL JOURNAL, 480:466–469, 1997 May 10 <http://www.iop.org/EJ/article/0004-637X/480/2/466/35523.text.html>

Kuchiev, M. Yu, “Can gravity appear due to polarization of instantons in SO(4)gauge theory?”, Class. Quantum Grav. 15 1895-1913 doi: 10.1088/0264-9381/15/7/008

Dawood Kothawala, T. Padmanabhan, Sudipta Sarkar, "Is gravitational entropy quantized?" *Phys. Rev.*, D78, 104018 (2008) [arXiv:0807.1481]

Kuchiev, M. Yu, "Can gravity appear due to polarization of instantons in SO(4)gauge theory?", *Class. Quantum Grav.* 15 1895-1913 doi: 10.1088/0264-9381/15/7/008

B Lamine, A Lambrecht, M T Jaekel and S Reynaud, "Relic gravitational wave background and the isotropy of space", *Class. Quantum Grav.* 21 S1209-S1214

Jean-Luc Lehners, Paul J. Steinhardt, "Non-Gaussian Density Fluctuations from Entropically Generated Curvature Perturbations in Ekpyrotic Models", *Phys.Rev.D77:063533,2008*; <http://arxiv.org/abs/0712.3779>

Li, Fangyu, Baker, Robert. M.L., and Chen, Zhenya (2006), "Perturbative photon flux generated by high frequency relic gravitational waves and utilization of them for their detection," <http://arxiv.org/abs/gr-qc/0604109>

Li, Fangyu, Baker, Robert M L., Fang, Zhenyun, Stephenson, Gary V., and Chen, Zhenya, "Perturbative Photon Fluxes Generated by High-Frequency Gravitational Waves and Their Physical Effects," [Li-Baker Chinese HFGW Detector], *European Physical Journal C.*, 56, 407- 423 (2008), <http://www.gravwave.com/docs/Li-Baker%206-22-08.pdf>

J. A. S. Lima, "Cosmologies with Photon Creation and the 3K Relic Radiation Spectrum", *Gen.Rel.Grav.* 29 (1997) 805-813; <http://arxiv.org/abs/gr-qc/9605056>

A.D. Linde, "The New inflationary Universe Scenario", pp 205-249, from the volume, "The Very Early Universe", edited by G.W.Gibbons, S.W. Hawking, S.T.C. Siklos, Cambridge University press, Cambridge, UK, 1982.

Eric Linder, "Exploring the Expansion History of the Universe", *Phys. Rev. Lett.* 90, 091301 (2003)

Lifschytz, G., "Black Hole thermalization rate from brane antibrane model", arXIV hep-th/0406203 v1 23 june, 2004

Linde in article in *Physics 738, Inflationary cosmology*, editors M. Lemoine, J. Martin, P. Peter, as Editors, Springer Verlag, 2008 Berlin, Federal Republic of Germany

Lloyd, Seth "Computational capacity of the universe", *Phys. Rev. Lett.* 88, 237901 (2002)

David H. Lyth, "Generating curvature perturbations at the end of inflation", *JCAP* 0511 (2005) 006; <http://arxiv.org/abs/astro-ph/0510443>

Roy Maartens, "Brane-World Gravity (2004)", <http://www.livingreviews.org/lrr-2004-7>

J. Martin, "Inflationary Perturbations: The cosmological Schwinger effect", pages 193-241, *Lecture notes in Physics, 738*, editors M. Lemoine, J. Martin, P. Peters, Springer - Verlag (2008), Berlin

Mathur, S., and Chowdhury, B., "Fractional Brane States in the Early Universe", *Classical and Quantum Gravity*, 24 (2007), pp. 2689-2720

Mina, A., Takuya, O., and Hiroshi, O., "Quantum entanglement of baby universes", *Nuclear Physics B, Volume 778, Issue 1-2*, p. 36-68. (2007)

Prado Martin-Moruno, Pedro F. Gonzalez-Diaz, "Thermal radiation from Lorentzian traversable wormholes", *Phys. Rev. D* 80, 024007 (2009); <http://arxiv.org/abs/0907.4055>

Ng, Y. Jack, "Article: Spacetime Foam: From Entropy and Holography to Infinite Statistics and Nonlocality" *Entropy* 2008, 10(4), 441-461; DOI: 10.3390/e10040441

Ng, Y. Jack, "Quantum Foam and Dark Energy", *International work shop on the Dark Side of the Universe*, <http://ctp.bue.edu.tw/workshops/Talks/Monday/QuantumFoamAndDarkEnergy.pdf>

Kouros Nozari and Tahereh Azizi, "Quantum Mechanical Coherent States of the Harmonic Oscillator and the Generalized Uncertainty principle", *International Journal of Quantum Computing*, Vol 3, No 4, (2005), pp623-632

Peskins, M., Schroeder, D. "An introduction to Quantum Field Theory", Westview Press, (1995)

Carlo Rovelli, "Graviton Propagator from Background-Independent Quantum Gravity", *Phys. Rev. Lett.* 97, 151301 (2006); <http://arxiv.org/abs/gr-qc/0508124>

Leszek M. Sokolowski, Andrzej Staruszkiewicz, "On the issue of Gravitons", *Class.Quant.Grav.* 23 (2006) 5907-5918, <http://arxiv.org/abs/gr-qc/0606111>.

Steinhardt, Paul J., Turok, Neil, "Endless Universe, Beyond the Big Bang" Doubleday Broadway Publishing Group, a division of Random House, Inc., New York (2007); <http://map.gsfc.nasa.gov/media/060915/index.html>

Gary Stephenson, (Private discussions, August 20, 2009), w.r.t. Li-Baker detector sensitivity.

Ming-Lei Tong and Yang Zhang (2007), "Detecting very-high-frequency relic gravitational waves by electromagnetic wave polarizations in a waveguide", *Chin. J. Astron. Astrophys.* 8 (2008) 314-328, <http://arxiv.org/abs/0711.4909>

K.K.Venkataratnam, P.K.Suresh, "Density Fluctuations in the Oscillatory Phase of Nonclassical Inflaton in FRW Universe", *Int.J.Mod.Phys.D17:1991-2005,2008*, <http://arxiv.org/abs/0712.3129>

Licia Verde, "Non Gaussianity versus non linearity of cosmological perturbations"; *proc. of the 15th Florida workshop in Nonlinear Astronomy and Physics. Annals of The New York Academy of Sciences* as well as Clifford Will, "Bounding the mass of the graviton using gravitational-wave observations of inspiralling compact binaries", <http://arxiv.org/abs/gr-qc/9709011>