

Applications of Euclidian Snyder geometry to the foundations of space-time physics

Dr. Andrew Beckwith

American Institute of Beam Energy Propulsion, life member (I will put in the address later!)

The following document is to answer if higher dimensions add value to answering fundamental cosmology questions. The results are mixed, **1st** with higher dimensions apparently helping in reconstructing and preserving the value of Planck's constant, and the fine structure constant from a prior to a present universe, while **2nd** failing to add anything different from four dimensional cosmological models to the question of what would cause an increase in the expansion rate of the universe, as of a billion years ago. Finally **3rd**, higher dimensions may allow creation of a joint DM and DE model. A choice between LQG and brane world geometry is introduced by Snyder geometry, where Snyder geometry's minimum uncertainty length calculations Δx may help determine to what extent gravity is an emergent field that is classical. Independent of the choice of LQG and branes (four dimensions versus higher dimensional cosmology models) is the following question: If gravity is largely classical, how much nonlinearity is involved? Gravitons and their structure as information carriers may help answer these questions. The main point of this document: DM and DE may be unified in terms of cosmological dynamics if the higher dimensional models of DM, as seen by KK towers of gravitons are seen to be pertinent to increasing acceleration of the universe a billion years ago via a 4th dimensional small graviton mass term added to the KK tower DM representation of gravitons (a model of DM).

PACS: 89.70.Cf, 95.35.+d, 95.36.+x

I. INTRODUCTION

Recently, a big bounce has been proposed¹ as an alternative to singularity conditions that Hawkings, Ellis [1], and others use. The **1st problem** is though that there appears to be no fundamental argument presented in either traditional Friedman metric GR or LQG for preservation of the same value for Planck's constant or the fine structure constant from prior universes (before ours) and the present universe. If \hbar and α (fine structure) constant were very different prior to our present universe, then cosmological evolution would be a highly random process. Ashtekar [2] ,in the inaugural opening of the Penn State gravity center (2007) told the author that the universe preserves most of its " memory" in cosmological cycles, but the proof of this assertion does not show up in Rovelli's [3] r . This present document tries to ascertain what could be a way to preserve some of prior universe "memory" through the use of gravitons as information carriers. . I.e. as a way about the singularity theorems of classical GR [1].The author is aware of Goswami et. al's. proof [4] of how cosmological singularities can be resolved in four dimensions, but no evidence of a structure to preserve \hbar and α constant appears to emerge from a prior universe to today's cosmos. Furthermore, both higher dimensional cosmologies such as brane theory , and more classical 4 dimensional cosmologies can be employed to (if gravitons have a small rest mass) account for the rate of expansion of the universe increasing one billion years ago, without invoking DE. The **2nd problem** is that, as demonstrated by Alves et. al. [5] four dimensional geometries **also** can account for the same increase in acceleration of the cosmos, one billion years ago. In particular, Batistini [6] uses Snyder geometry to find a limiting approximation to determine either brane world or LQG conditions for cosmological evolution. The motivation for a choice between either brane world (5 or higher dimensions) and LQG (four dimensions) comes from Alves et. al. [5] with regards to their following argument : "It is worth stressing that Gabadadze and Gruzinov [7] have analyzed the instabilities of the background and ghosts produced by massive gravitons in 4D Minkowski space-time. They conclude that a natural way to account for a massive graviton on a(t) space is to invoke theories with extra dimensions. However, Visser's model [8] is truly continuous with general relativity (GR) in the limit of vanishing graviton mass. Together with Pauli-Fierz (PF) massive term, Visser's theory is the simplest attempt to incorporate mass for the graviton in a ghost-free manner. Moreover, the van Dam-Veltman-Zakharov discontinuity [9] (vDVZ) present in the PF term can be circumvented in Visser's model by introducing a non-dynamical at-background metric [10] " While

¹ Papers on LCQ at the 12th Marcell Grossman Meeting in 2009 (<http://www.icra.it/MG/mg12/en/>)

this paragraph is excellent physics, it does not answer if preserving the values of \hbar and α (fine structure) constant between a prior to present universe is possible. The heart of [6] is a deformed Euclidian Snyder space, using $\hbar = c = 1$ units, obtaining $[x, p] = i \cdot \sqrt{1 - \alpha \cdot p^2} \Leftrightarrow \Delta x \Delta p \geq \frac{1}{2} \cdot \left\langle \left\langle \sqrt{1 - \alpha \cdot p^2} \right\rangle \right\rangle$. The LQG condition implies $\alpha > 0$, while brane worlds have instead, $\alpha < 0$. As suggested in [6], it is possible to obtain a string-theory limit of $\Delta x \geq \left[(1/\Delta p) + l_s^2 \cdot \Delta p \right] \equiv (1/\Delta p) - \alpha \cdot \Delta p$. The Δx lower bound will be set depending upon either using LQG condition $\alpha > 0$, or brane worlds condition. $\alpha < 0$ Finally, if higher dimensions of DM models, projected to 4 dimensions yield a KK type model of DM, with the lowest order mode of DM (4 dimensions) mimicking DE, then a new phenomenological model of partial unity between DM and DE is presented. Which may be better than the Chapygin Gas model, as stated by Roos [11]

A. What role do gravitons play in distinguishing between $\alpha < 0$ and $\alpha > 0$ in terms of Snyder geometry?

The author finds no difference in either $\alpha < 0$ and $\alpha > 0$ situations as far as determining if a speed up of the universe's rate of expansion occurred a billion years ago. Alves [5] shows that an accelerating expansion at the present time is related to the value of the graviton mass, while not using contributions from higher dimensions. If $m_{graviton} \cong 205 \times 10^{-65}$ grams in four dimensions yields few difference in behavior in the LQG and Brane world situations in a speed up of the rate of acceleration of the cosmos one billion years ago, then what worth are higher dimensions to cosmology? The next paragraph presents how higher dimensions may preserve the values of \hbar and α (fine structure) constant from prior to present universes. To do that, the author tries to establish the graviton as a carrier of information placed into both initial values of \hbar and α at the big bang. Normalized energy density of gravitational waves, as given by Maggiore [12] is

$$\Omega_{gw} \equiv \frac{\rho_{gw}}{\rho_c} \equiv \int_{\nu=0}^{\nu=\infty} d(\log \nu) \cdot \Omega_{gw}(\nu) \Rightarrow h_0^2 \Omega_{gw}(\nu) \cong 3.6 \cdot \left[\frac{n_\nu}{10^{37}} \right] \cdot \left(\frac{\nu}{1kHz} \right)^4 \quad (1.1)$$

Where n_ν is a frequency-based count of gravitons per unit cell of phase space. In terms of early universe nucleation, the choice of n_ν may depend upon interaction of gravitons with neutrinos [13]. The supposition is that eventually, (1.1) could be modified with a change of

$$n_\nu \propto n_\nu[\text{graviton}] + n_\nu[\text{neutrinos}] \quad (1.2)$$

Eq. (1.2) is part of a weighted average of neutrino-graviton coupled frequency $\langle \nu \rangle$, so that for detectors,

$$h_0^2 \Omega_{gw}(\nu) \cong \frac{3.6}{2} \cdot \left[\frac{n_\nu[\text{graviton}] + n_\nu[\text{neutrino}]}{10^{37}} \right] \cdot \left(\frac{\langle \nu \rangle}{1kHz} \right)^4 \quad (1.3)$$

Having confirmed a linkage between gravitons and GW energy density the next step will be to confirm if possible that higher dimensional cosmology models do not invalidate or alter the Alves [5] results

II. Proof that gravitons with a small mass in four dimensions, $m_{graviton} \cong 205 \times 10^{-65}$ grams, contribute to a brane world speed up of the acceleration of the universe

As suggested by Beckwith [14], gravitons may contribute to the re-acceleration of the universe one billion years ago, when both LQG and brane world models showed increased acceleration of the rate of expansion. Is there direct proof of the above assertion? Yes, if the value of $q(z)$ as appearing in Fig 1 would be almost the same in LQG and brane world geometry. . The results for Beckwith [14] show the same speed up of acceleration which Alves [5] obtained for cosmological expansion one billion years ago, but with higher dimensions. Generally, as specified by [5, 11, 14], the dimensionless decelerating parameter, Eq. (1.4),

models the rate of expansion of the universe, q , as a function of the scale factor $a \sim 1/(1 + \text{redshift})$. When q becomes negative, the rate of acceleration of the universe is increasing [5], [11], [14]

$$q = -\frac{\ddot{a}a}{\dot{a}^2} \quad (1.4)$$

Eq. (1.10) follows what Beckwith [14] did (to duplicate [5] for a brane world) in order to plot an analysis of the deceleration, $q(z)$, with z set = X in the calculation for Eq. (1.4). Eq. (1.10) is the input for for Figure 1 of page 6. The derivation of Eq. (1.16) as explained by **Appendix A** assumes setting a small mass for the graviton, for a brane world treatment of the Friedman equation, with the density of a brane world given in Eq. (1.5), as used by Alves, et al. [6] As, in **Appendix A**

$$\rho \equiv \rho_0 \cdot \left(\frac{a_0}{a}\right)^3 - \left[\frac{m_g c^6}{8\pi G \hbar^2}\right] \cdot \left(\frac{a^4}{14} + \frac{2a^2}{5} - \frac{1}{2}\right) \quad (1.5)$$

Eq. (1.5) assumes use of the following inequality for a change in the HUP, [5]

$$\Delta x \geq \left[(1/\Delta p) + l_s^2 \cdot \Delta p\right] \equiv (1/\Delta p) - \alpha \cdot \Delta p \quad (1.6)$$

and that the mass of the graviton is partly due to the stretching alluded to by Fuller and Kishimoto [13], a supposition the author is investigating for a modification of a joint KK tower of gravitons, as given by Maartens [16] for DM. Assume that the stretching of early relic neutrinos that would eventually lead to the KK tower of gravitons--for when $\alpha < 0$, is [14], which can be understood as given by **Appendix B**

$$m_n(\text{Graviton}) = \frac{n}{L} + 10^{-65} \text{ grams} \quad (1.7)$$

Also assume that to calculate the deceleration, the following modification of the HUP is used: [2] $\Delta x \geq \left[(1/\Delta p) + l_s^2 \cdot \Delta p\right] \equiv (1/\Delta p) - \alpha \cdot \Delta p$, where the LQG condition is $\alpha > 0$, and brane worlds have, instead, $\alpha < 0$. The deceleration parameter in (1.4) will have either higher-dimensional contributions, in the brane theory case, or no higher-dimensional contributions, in the LQG case. Also Eq. (1.8) will be the starting point used for a KK tower version of Eq. (1.4) [14] So from Maarten's 2005 paper, [16]

$$\dot{a}^2 = \left[\left(\frac{\tilde{\kappa}^2}{3} \left[\rho + \frac{\rho^2}{2\lambda} \right] \right) a^2 + \frac{\Lambda \cdot a^2}{3} + \frac{m}{a^2} - K \right] \quad (1.8)$$

Maarten also gives a 2nd Friedman equation, as

$$\dot{H}^2 = \left[- \left(\frac{\tilde{\kappa}^2}{2} \cdot [p + \rho] \cdot \left[1 + \frac{\rho^2}{\lambda} \right] \right) + \frac{\Lambda \cdot a^2}{3} - 2 \frac{m}{a^4} + \frac{K}{a^2} \right] \quad (1.9)$$

Also, if we are in the regime for which $\rho \cong -P$, for red shift values z between zero to 1.0-1.5 with exact inequality for z between zero to .5. The net effect will be to obtain, due to Eq. (1.10)

$$q = -\frac{\ddot{a}a}{\dot{a}^2} \equiv -1 - \frac{\dot{H}}{H^2} = -1 + \frac{2}{\tilde{\kappa}^2 [m/a^4] \cdot (\rho + \rho^2/2\lambda) + 1} \approx -1 + \frac{2}{2 + \delta(z)} \quad (1.10)$$

Eq. (1.10) assumes $\Lambda = 0 = K$, and the net effect is to obtain, a substitute for DE, by presenting how gravitons with a small mass in four dimensions can account for reacceleration of the universe, a datum which is usually done with $\Lambda \neq 0$, even if curvature \mathbf{K} is set equal to zero.

Note Maartens [16] states that KK modes (graviton) satisfy a 4-dimensional Klein-Gordon equation, with an effective 4-dim mass, $m_n(\text{Graviton}) = \frac{n}{L}$ with $m_0(\text{Graviton}) = 0$, and L as the stated "dimensional value" of higher dimensions. The value $m_0(\text{Graviton}) \sim 10^{-65} - 10^{-60}$ gram is almost zero. A non-zero

mass for the graviton is a violation of the correspondence principle for spin two objects, in quantum mechanical reasoning. Grossing [19] and Baker-Jarvis, and Kabos [20] have shown how the Schrodinger and Klein Gordon equations can be derived from classical Lagrangians, i.e., using a version of the relativistic Hamilton-Jacobi -Bohm equation, with a wave functional $\psi \sim \exp(-iS/\hbar)$, with S the action, so as to obtain working values for a tier of purported masses of a graviton from the equation, for 4-D of making the flat space approximation

$$\left[g^{\alpha\beta} \partial_\alpha \partial_\beta \xrightarrow{FLAT-SPACE} \nabla^2 - \partial_\tau^2 \right], \text{ and } \left[\nabla^2 - \partial_\tau^2 \right] \cdot \psi_n = m_n^2(\text{graviton}) \cdot \psi_n. \text{ This involves a small graviton mass, while assuming that a small added mass, } m_0(\text{Graviton}) \approx 10^{-65} \text{ grams, is a result of a semi-classical superstructure which containing the usual field theory/brane world treatment of quantum theory gravitons. All this discussion will show up in Appendix B below to derive Eq. (1.10). For book keeping, the author uses the substitutions, } (a_0/a)^3 = (1+z)^3, \text{ and } a \equiv a_0/(1+z), \text{. Then the density value}$$

$$\rho(z) \equiv \rho_0 \cdot (1+z)^3 - \left[\frac{m_g}{8\pi G} \right] \cdot \left(\frac{a_0^4}{14 \cdot (1+z)^4} + \frac{2a_0^2}{5 \cdot (1+z)^2} - \frac{1}{2} \right) \quad (1.11)$$

Where

Negative values for Eq. (1.10) appear probable at about $z \sim 1.5$, when (1.14) would dominate, leading to $q(z \sim 1.5)$ with a negative value. The positive value conditions rely upon contributions from the C dark radiation term in Eq. (1.10). The final result is that the deceleration parameter calculation can be done for the brane world case and KK gravitons. Now what can one expect with LQG condition with respect to the HUP, with $\alpha > 0$? The claim is made by Beckwith that LQG, due to the starting point of the LQG Friedmann equations, similar to what Alves [5] used, duplicates deceleration results by Beckwith [13]




Fig. 1: Reacceleration of the universe based on [13] (note that $q < 0$ if $z < .42$): Motivated by a similar diagram in Alves [5], where $q(z)$ is the rate of deceleration of the universe due to $m_{\text{graviton}} \cong 205 \times 10^{-65}$ grams. Limits of applicability are in line with where $p = -\rho$, i.e. $w = -1$,

Note $z = 1.5$ is 4.5 billion years after the big bang. Figure 1 is to be compared with Figure 4, page 11, of Alves [5] The significance of figure 1 lies in that higher dimensions in themselves do not change, if a four dimensional representation of the graviton has a low value mass of $m_{\text{graviton}} \cong 205 \times 10^{-65}$ gram what is seen in four dimensions. Notice what the author claims is a break down of applicability of the same figure, i.e. for $z = 3.0$, the plot of $q(z)$ asymptotically approaches, +1, but flattens as to approaching +1. This is also where $p = -\rho$ no longer holds. **Also, $z = 3$ corresponds to 2.2 billion years after the big bang**

B. Using the LQG condition $\alpha > 0$, in Snyder geometry modified HUP

The claim is that almost all the complexity is removed with $\alpha > 0$, and what is left is a set of equations similar to the tried and true $\left(\frac{\dot{a}}{a}\right)^2 = \frac{\rho_{Total}}{3M_{Planck}^2} - \frac{k}{a^2} + \frac{\Lambda}{3}$. To get an idea of what happens with LQG versions of the Friedman equation, one can look at the Taveras [21] treatment of the Friedmann equations, where he obtains, to first order, if ρ is a scalar field density,

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{\kappa}{3} \cdot \rho \quad (1.12)$$

and

$$\left(\frac{\ddot{a}}{a}\right) = -\frac{2 \cdot \kappa}{3} \cdot \rho \quad (1.13)$$

The interpretation of ρ as a scalar field DENSITY, and if one does as Alves et al [5] .did, i.e., work with

flat space, then with $k=0$, in $\left(\frac{\dot{a}}{a}\right)^2 = \frac{\rho_{Total}}{3M_{Planck}^2} - \frac{k}{a^2} + \frac{\Lambda}{3}$, $\rho \equiv \rho_0 \cdot \left(\frac{a_0=1}{a}\right)^3 - \left[\frac{m_g c^6}{8\pi G \hbar^2}\right] \cdot \left(\frac{a^4}{14} + \frac{2a^2}{5} - \frac{1}{2}\right)$

The sticking point in all of this is to interpret the role of ρ . In the LQG version by [21], $\left(\frac{\dot{a}}{a}\right)^2 = \frac{\kappa}{3} \cdot \rho$

may be rewritten as follows: If conjugate momentum is in many cases, "almost" or actually a constant,

$$\left(\frac{\dot{a}}{a}\right)^2 \equiv \frac{\kappa}{6} \cdot \frac{p_\phi^2}{a^6} \quad (1.14)$$

This assumes that the conjugate dimension in this case has a quantum connection specified via an effective scalar field, ϕ obeying the relationship

$$\dot{\phi} = -\frac{\hbar}{i} \cdot \frac{\partial}{\partial \cdot p_\phi} \quad (1.15)$$

Alves research results [5] for both LQG and brane worlds means using the Friedmann equations directly as a way to computer deceleration of the universe, from the Friedmann equations, especially if the graviton has nearly zero rest mass. To do that, we delve into a LQG version of deceleration parameter behavior.

C. What 4-dimensional representations of the Graviton are in common with both models for both $\alpha < 0$ and $\alpha > 0$?

There are two hypotheses to consider. The first is that there is an interaction between neutrinos and gravitons. Bashinsky [22] suggests an alleged modification of density fluctuations via neutrino-graviton interactions. A second far more radical hypothesis is that there are a few "stretched neutrinos" [4] that may span many light years, and these may affect gravitons, as suggested by Bashinsky [22]. What is being considered is that there are graviton-neutrino interactions, as proposed by [22], and [13] asks, what are the natures of the neutrino-graviton interactions if a few of the neutrinos "stretch" by many light years? If there is a coupling between gravitons and neutrinos, as by [18], stretched neutrinos leading to stretched gravitons brings into question the correspondence principle, that gravitons be spin 2, with zero mass.

1. The probable effect of stretching of the neutrino on graviton wavelengths

Assume that with stretching of the neutrino, and a graviton neutrino coupling with zeroth order value of $m_0(Graviton) \approx 10^{-65}$ grams as a consequence of at least a few of the neutrino-gravitons obeying density

fluctuation modified , according to [22], $\left[1 - 5 \cdot (\rho_{\text{neutrino}} / \rho) + \mathcal{G}[(\rho_{\text{neutrino}} / \rho)^2]\right]$. Note that according to [14], the overall density of the evolving space-time continuum has neutrino-graviton interactions that effectively shrink the magnitude of overall space-time density. In addition, neutrino and graviton wavelengths then have the same order of magnitude as the matter wavelength values of neutrinos, with, initially

$$\begin{aligned} m_{\text{graviton}} \Big|_{\text{RELATIVISTIC}} &< 4.4 \times 10^{-22} h^{-1} eV / c^2 \\ \Leftrightarrow \lambda_{\text{graviton}} &\equiv \frac{\hbar}{m_{\text{graviton}} \cdot c} < 2.8 \times 10^{-8} \text{ meters} \end{aligned} \quad (1.16)$$

A few select gravitons, coupled to stretched neutrinos with almost infinite wavelengths, would lead to (1.17), if the graviton wavelengths were, according to an argument ventured by Valev [23],

$$\lambda_{\text{graviton}} \equiv \frac{\hbar}{m_{\text{graviton}} \cdot c} < 10^4 \text{ meters or larger} \quad (1.17)$$

III. THE CORRESPONDENCE PRINCIPLE AND 'T HOOFT'S SUPPOSITION OF 'DETERMINISTIC QM' AS APPLIED TO GRAVITONS

The author suggests that the stretch out of the graviton implied by (1.39) may be a sign that the correspondence principle, used by string theorists and others as a way to insist that the graviton be of zero mass, may have to be amended. After presenting why the author states that the correspondence principle needs revision, the author will suggest a mechanism for replacement of the correspondence principle, which the author suggests is consistent with 't Hooft's [24,25] deterministic quantum mechanics. The final part to this paper suggests what "information" a particle like the graviton may carry. What can be stated about the "correspondence principle" and its connections to gravitons? Rothman and Boughn [26] suggest that it is unrealistic to envision gravitons ever being measured. However, it assumes current detector technology. Finally, the author will look into what a graviton "construction" with a tiny mass may entail regarding instanton-anti-instantons, and its relationship to 't Hooft's [24, 25] deterministic quantum mechanics. To recap what they are suggesting as a way to quantify the scattering needed to observe a graviton in a detector, Rothman, and Bohn [26] suggest

$$\tilde{n} \cdot \sigma \cdot \tilde{\lambda} \geq 1 \quad (1.18)$$

Where \tilde{n} is the purported numerical density of "detector particles," σ is the detector cross-sectional area, and $\tilde{\lambda}$ is the mean "distance" a graviton would have to travel. However, [26] assumes the cross-sectional area for a graviton would have to be larger "than the diameter of Jupiter." Note that the variable \tilde{n} is given by [26] to be $\tilde{n} \equiv M_{\text{det}} / [m_{\text{proton}} \cdot V_{\text{det}}]$. I.e., this is for a detector with gravitons interacting with some version of hydrogen, with M_{det} the "mass" of the detector, and with V_{det} the purported volume of the detector. Also, m_{proton} is the mass of protons in the detector that the gravitons may interact with. If so, the volume V_{det} being Jupiter-sized may look reasonable. However, this assumes gas-based graviton detector technology, which uses collision cross sections. Electromagnetic and graviton interactions may allow for a far smaller V_{det} , according to [27], and [28], using a different numerical count procedure for gravitons in a unit of phase space. Beckwith [27] uses a very explicit numerical count, which can be interpreted as a phase space count from fundamental principles. Li, et al, [28] uses electromagnetic fields as affected by gravitons in a containment vessel, as a way to "count" incident gravitons. These two alternatives raise the question of the utility of Eq. (1.23) as the optimal way to measure gravitons in experimental devices. Eq. (1.23) implies a numerical count of gravitons detected during the lifetime of an experiment, where $L_{\text{graviton-production}}$ is the luminosity of graviton production, R as the purported distance the graviton would

travel, while setting up the right-hand side with $\frac{A_{\text{det}} \cdot \tau_B}{\in_{\text{graviton}}} \equiv (\text{detector cross sectional area} \cdot \text{time of process})$

for the graviton source to be operating) / graviton energy . Also, the time limit is $\tau_B \leq \frac{M_{\text{graviton-generating-source}}}{L}$. Here $M_{\text{graviton-generating-source}}$ is the relative mass of the graviton producing source, and L the luminosity of the source. The bounds for τ_B are effectively exceeded by kinematic considerations if the graviton production "site" is relic early universe gravitons, instead of what is cited for non-zero graviton energies, $\epsilon_{\text{graviton}}$

$$N_{\text{graviton=exp-lifetime}} \equiv \left[\frac{L_{\text{graviton-production}}}{4\pi R^2} \right] \cdot \left[\frac{A_{\text{det}} \cdot \tau_B}{\epsilon_{\text{graviton}}} \right] \quad (1.19)$$

Rothman and Boughn [26] give a coherent argument that for neutron stars, black holes and the like, Eq. (1.19) has an upper bound of $N_{\text{graviton=exp-lifetime}} \approx 10^{-5}$. The author suggests that the total source luminosity L versus luminosity of graviton production process of the source $L_{\text{graviton-production}}$ may be different from the ratio values given by Rothman, and Bohn, of $L_{\text{graviton-production}}/L = f_{\text{graviton}}$ as $\sim .01-.02$. If the f_{graviton} is ten times larger, plus the life time $\tau_B \leq \frac{M_{\text{graviton-generating-source}}}{L}$ of graviton production from

black holes with a larger time due to having a value of $M_{\text{graviton-generating-source}} \gg 10^{15}$ grams, with 10^{15} grams \sim mass of a black hole, then $N_{\text{graviton=exp-lifetime}} \approx 10^{-5}$ may be way too small. Furthermore, if the stretched neutrino hypothesis, with coupling to the graviton, occurs, then, assuming that there is at a minimum $\lambda_{\text{graviton}} \equiv \frac{\hbar}{m_{\text{graviton}} \cdot c} < 10^4 \text{ meters}$, instead of $\lambda_{\text{graviton}} \equiv \frac{\hbar}{m_{\text{graviton}} \cdot c} < 2.8 \times 10^{-8} \text{ meters}$. Even

with a non-giant planet sized detector, one would see a count of data that $N_{\text{graviton=exp-lifetime}} \gg N_{\text{Rothman-calculated-graviton-exper-lifetime}} \approx 10^{-5}$, perhaps with $N_{\text{graviton=exp-lifetime}}$ as nearly unity. And this is due to recalibration of the different input coefficients. The inequality given counting $N_{\text{graviton=exp-lifetime}} \gg N_{\text{Rothman-calculated-graviton-exper-lifetime}}$ raises a question regarding the use of standard correspondence principle to characterize gravitons, and suggests an alternative: gravitons with spin 2, but perhaps masses slightly larger than zero. Eventually, this will lead -to considering the correspondence principle, as well as "deterministic" quantum mechanics as't Hooft wrote in [25] as a way to consider the nature of gravitons

A. Can the graviton have a small mass? Embedding the laws of QM regarding gravitons in a nonlinear theory.

Recently, an alternative to usual space-time gravitation theories was proposed by Mukohyama.. Which is called HoYYava gravity [29] Brandenberger [30] also modeled this new theory in terms of the early universe, with the claim that there was a matter bounce instead of standard inflation. This theory, ironically, depends upon a chaotic inflationary potential $V(\phi) = (1/2) \cdot m^2 \phi^2$ for its pre-bounce conditions, and uses "dark radiation" for obtaining a "bounce." [30] has also presented "scale-invariant, super-horizon curvature perturbations." Both [29] and [30] accept scale-free "perturbations" as long as the contraction phase does use "quantum vacuum fluctuations." So it will be necessary to wait to see if HoYYava gravity develops further, or is provided with a mechanism to transfer energy to the standard model of cosmology predictions for the radiation and matter eras. Visser presented [31] HoYYava gravity in terms of the "benefits of Lorentz symmetry breaking" with a nonphysical graviscalar spin zero "graviton" contribution [27], which, if not removed, yields mathematical artifacts with no physical content in them. Spin zero gravitons (gravit scalars) completely eliminate string theory as a consideration in gravitons. Note that at that the lowest vibration of a cosmological string, a spin two particle with zero mass, can be interpreted as a graviton, Graviton modes with spin zero would sever the connection to quantum mechanics, and complicate making semi classical analogies to quantum mechanics. Visser viewed the removal of the gravi scalar as crucial to

the proof of the supposition about semi classical gravity models. Gravi scalars would require a complete break with the correspondence principle of quantum mechanics for spin two quantum spin particles [33]. By way of contrast, what the author will attempt to do is to with gravitons is far more modest: referencing the construction of a graviton in terms of instanton-anti-instantons, and asking if a composition of a graviton as an "object" comprising such kink-ant kinks can be tied in with 't Hooft's "deterministic quantum mechanics" [24,25] . Beginning the analysis, the author will review what he did with CDW in $1 + \varepsilon^+$ dimensions, and then reference the chances for doing the same for four dimensions for gravitons, closing with a discussion of the ability of the graviton to carry information .

B. Brief review of S-S' in CDW and its relevance to higher dimensional 'objects'

The following is a presentation of Beckwith's density-wave instanton-anti-instanton construction for CDW, which has classical analogies. Beckwith's kink-anti-kink models [32, 34] have a classical analogy with

$$\phi_{\pm}(z, \tau) = 4 \cdot \arctan \left(\exp \left\{ \pm \frac{z + \beta \cdot \tau}{\sqrt{1 - \beta^2}} \right\} \right) \quad (1.20)$$

which is a solution to

$$\frac{\partial^2 \phi(z, \tau)}{\partial \tau^2} - \frac{\partial^2 \phi(z, \tau)}{\partial z^2} + \sin \phi(z, \tau) = 0 \quad (1.21)$$

A tunneling Hamiltonian version of such solutions uses a Gaussian wave functional formalism, with

$$\Psi_{i,f} [\phi(\mathbf{x})]_{\phi=\phi_{ci,cf}} = c_{i,f} \cdot \exp \left\{ - \int d\mathbf{x} \alpha \left[\phi_{Ci,f}(\mathbf{x}) - \phi_0(\mathbf{x}) \right]^2 \right\}, \quad (1.22)$$

This procedure in Eq. (1.22) allowed for deriving a confirmation [32,34] of the fit between the false vacuum hypothesis and data obtained for current vs. applied electrical field (I-E) curves of experiments initiated in the mid-1980s by [35]. Further research by Beckwith led to the finding that the modulus of the tunneling Hamiltonian was proportional to current, with E_T a threshold pinning field [32, 34]. The threshold pinning field, as presented by Beckwith [32] is an absolute magnitude of applied electric field for which above a magnitude for an applied electric field there is current flow in CDW 'information' in a laboratory measurement of current versus electric field strength in NbSe₃--. This phenomenon was presented experimentally in Miller, et al, [35] in the mid 1980s..Pinning is an artifact of historical terminology and is less important than the term threshold. Below this applied electric field threshold, no CDW transport occurred. The use of instantons and judicious use of Eq. (1.23) is necessary for modeling a current of CDW waves, which cannot be obtained by conventional CDW models {32, 34}. I.e. the use of instanton-anti-instanton structure is necessary for modeling I-E behavior in quasi-one- dimensional systems. A similar situation arises in cosmology, once evolution equations are defined.

$$I \propto \tilde{C}_1 \cdot \left[\cosh \left[\sqrt{\frac{2 \cdot E}{E_T \cdot c_V}} - \sqrt{\frac{E_T \cdot c_V}{E}} \right] \right] \cdot \exp \left(- \frac{E_T \cdot c_V}{E} \right) \quad (1.23).$$

The phase used in Eq. (1.22) is indicative of what instanton physics can be used for, i.e., this is not a substitute for a well thought out treatment of instantons, which will be connected via cited derivations which link instanton models with appropriate metrics, as is discussed explicitly with respect to GR space time metrics in **Appendix C**. Fig. 4 in particular shows how the author will model a pop up effect of a S-S' pair, in a quantum mode, using S and S' pairs. Note in the case of CDW the author [32] found a current

derived as being of the form $J \propto T_{if}$. I.e. a current proportional to a density wave tunneling Hamiltonian matrix element. This argument actually became a modulus argument due to considering a current density proportional to $|T|$ rather than $|T|^2$ since tunneling, in this case, would involve coherent transfer of individual (first-order) bosons rather than pairs of fermions. The use of $J \propto T_{if}$ instead of having current proportional to $|T|^2$ permitted the author to come up with current versus applied electric field graphs which matched experimental Zenier plots of CDW obtained by Miller in the mid 1980s [35]. The author rules out time independent Wheeler De Witt equations with Hamiltonians [33] for this sort of detailed analysis of graviton particle creation. In particular, there is no counter part to a tunneling Hamiltonian in cosmology which can take the place of what was done in the authors CDW studies [32], [34]. However, the overlap in the case of cosmology is in the use of solution- anti solution pairs, which allow for abrupt





Figure 2, creation of instanton – anti instanton pairs. From Beckwith, [30]

movement of gravitons, a topic which will be dealt with extensively later. The author argues that instanton- anti instanton (kink- anti kink) construction in both CDW [32,34] and its counter part in cosmology will give, as indicated in Figure 4 below, vacuum nucleation pop up, in the beginning of inflation, of gravitons. As can be quoted by Mukhanov [37] for times $10^{-12} - 10^{-43}$ seconds after the big bang, and at temperatures 10 TeV to 10^{19} GeV, those topological defects may play a role in the production of new particles is relevant to the author's presentation of a kink anti kink model of gravitons.



Electric field

Fig 3: Results of applying Eq. (1.23) as opposed to J, H. Miller’s Zenier plot [35]

$$I \propto G_p \cdot (E - E_T) \cdot \exp\left(-\frac{E_T}{E}\right) \quad \text{If } E > E_T ,$$

$$I = 0 \quad \text{if } E \leq E_T . \quad [35]$$

The blue dots represent Eq. (1.23) whereas the black dots represent uniformly applying the non zero plot for electric fields as given by the Zenier plot [35] approximation (as taken from the authors [32], [34]).




Fig. 4. The pop up effects of an instanton-anti-instanton in Euclidian space from [32, 34]

In order to connect with GR, one needs to have a higher dimensional analog of

$$\phi_{\pm}(z, \tau) = 4 \cdot \arctan\left(\exp\left\{\pm \frac{z + \beta \cdot \tau}{\sqrt{1 - \beta^2}}\right\}\right)$$

that is consistent with space-time metrics, a topic which will be presented in the next section. We need to associate Instanton structure with typical GW topologies, and the way to do for a five dimensional cosmology is indicated in Appendix A. The five-dimensional "line element" involves an instanton that is almost identical in structure to Eq. (1.22) and (1.23). Having mentioned this similarity in form for the instanton, it is appropriate to note that the fifth dimension of the line element may form a semi-classical conduit of information for vacuum nucleation of the instanton itself.

C. Dropping in ‘information’ to form an instanton-anti-instanton pair, and avoiding the cosmological singularity via the 5th dimension

As suggested by Beckwith [38], there is no reliable way to reconcile the formation of an instanton-anti-instanton pair, and, for example, to avoid having an instanton disrupted by a cosmological singularity. The

graphical example in [38] was suggested to consider the question: What if there were, in higher than four dimensions, a region of **space** about the four dimensional singularity at the beginning of expansion of the universe via the use of the fifth dimension in which an instanton-anti-instanton could have information transferred from a prior universe and re materialize in the onset of the big bang after the point of nucleation of a new universe, thereby avoiding the cosmic singularity?. This question lead to, in Chongqing the author led to defining in Chouguing [38] the region about the singularity as a ring of space-time about the origin, but not overlapping it, with a time dimension defined [38] as

$$\Delta t \equiv 10^\beta \cdot t_{Planck} \quad (1.24)$$

The exact uncertainty principle in five dimensions is open to discussion, but the author in his presentation in Chongqing [38] suggested, as an example, a five-dimensional version of $\Delta E \Delta t \geq \hbar$. For the tiny mass specified via the $m_{graviton} \propto 10^{-65}$ grams, if energy is equivalent to mass, then small mass times the speed of light, squared, in the case of instanton-anti-instanton (kink-anti-kink) is the S-S' pair for the instanton nucleated about the cosmic singularity. The classical treatment of this problem is to assume that the transfer of information from a prior universe to our own went through a 5th dimension with the cosmic singularity, a 4th dimensional artifact The initial 4 dimensional 'graviton' would have , $m_{graviton} \propto 10^{-65}$ grams, with, a top value for the graviton mass after acceleration of $m_{graviton} \propto 10^{-61}$ grams. This is abrupt acceleration, making the graviton mass at least 10^4 times heavier than initially. To understand the rationale for such a supposition, a brief review of typical field theories involving "massive" gravitons and the limit $m_{graviton} \rightarrow 0$ will be presented, with a description of why these effects may lead to semi-classical approximations.

D. Massive Graviton field theories, and the limit $m_{graviton} \rightarrow 0$

As given by M. Maggiore [12], the massless equation of the graviton evolution equation takes the form

$$\partial_\mu \partial^\sigma h_{\mu\nu} = \sqrt{32\pi G} \cdot \left(T_{\mu\nu} - \frac{1}{2} \eta_{\mu\nu} T^\mu_\mu \right) \quad (1.25)$$

When $m_{graviton} \neq 0$, the above becomes

$$\left(\partial_\mu \partial^\sigma - m_{graviton} \right) \cdot h_{\mu\nu} = \left[\sqrt{32\pi G} + \delta^+ \right] \cdot \left(T_{\mu\nu} - \frac{1}{3} \eta_{\mu\nu} T^\mu_\mu + \frac{\partial_\mu \partial_\nu T^\mu_\mu}{3m_{graviton}} \right) \quad (1.26)$$

The mismatch between these two equations, when $m_{graviton} \rightarrow 0$, is due to $m_{graviton} h^\mu_\mu \neq 0$ as $m_{graviton} \rightarrow 0$, which in turn is due to setting $m_{graviton} \cdot h^\mu_\mu = -\left[\sqrt{32\pi G} + \delta^+ \right] \cdot T^\mu_\mu$. The mismatch between these two expressions is one of several reasons for exploring what happens for semi-classical models when $m_{graviton} \neq 0$, $m_{graviton} \sim 10^{-65}$ grams, noting that in QM, a spin 2 $m_{graviton} \neq 0$ has five degrees of freedom, whereas the $m_{graviton} \rightarrow 0$ gram case has only two helicity states. Note that string theory treats gravitons as "excitations" of a closed string, as given by Keifer [39], with a term added to a space-time metric, \bar{g}_{uv} , such that $g_{uv} \equiv \bar{g}_{uv} + \sqrt{32\pi G} f_{\mu\nu}$ with $f_{\mu\nu}$ a linkage to coherent states of gravitons. This is partly in relation to the Venziano [40] expression of $\Delta x \geq \frac{\hbar}{\Delta p} + \frac{l_s^2}{\hbar} \Delta p$, where $G \sim g^2 l_s^2$.

Kieffer [39] gives a correction due to quantum gravity in page 179 of the order of $\left(\frac{m}{M_{Planck}} \right)^2$ If the mass, $m_{graviton} \sim 10^{-65}$ g, it will be hard to measure as an individual "particle." But, if $m_{graviton} \sim 10^{-65}$ grams exist, as a macro effect, it may play a role, as indicated by Fig. 1.

1. So, what about representing a graviton as a kink-anti-kink?

How does this fit in with 't Hooft's deterministic QM?

In 2006, 't Hooft [24] used an equivalence-class argument as an embedding space for simple harmonic oscillators, as given in figure 2 in [24]. It is also noteworthy to consider that in 2002, 't Hooft [25] also wrote in his introduction, that "Beneath Quantum Mechanics, there may be a deterministic theory with (local) information loss. This may lead to a sufficiently complex vacuum state." In addition, the embedding equivalence class structure may be a consequence of a family of wave functionals which can be written as

$$\Psi_{i,f} \left[\phi(\mathbf{x}) \right]_{\phi=\phi_{ci,cf}} = c_{i,f} \cdot \exp \left\{ - \int d\mathbf{x} \alpha \left[\phi_{Ci,f}(\mathbf{x}) - \phi_0(\mathbf{x}) \right]^2 \right\}, \text{ solutions to a graviton state, if one is}$$

taking the $\phi(x)$ as a kink-anti-kink combination. I.e., looking at a history plot of equivalent solutions to the graviton problem, in a 5 dimensional metric space, as outlined by Belunski and Verdager [41] and Giovannini [42]. This "functional" (if one assumes equivalence classes of solutions) may be part of a deterministic embedding space for the vacuum space of space-time where the graviton exists. Reformulating the above solution in terms of different values of $\phi_0(x)$ in a wave-functional representation of a graviton is consistent with for equivalence-class structures. This would mean, for example, a considerable refinement of the metric in 5 dimensions, as stated in Appendix A, [29], $dS^2 = a(w) \cdot [\eta_{uv} dx^u dx^v - dw^2]$. The closeness of $m_{graviton} \sim 10^{-65}$ to a zero mass should not be seen as a failure of quantum physics, but a success story. However, it is suggested that establishing equivalence classes as part of a procedure to embed gravitons in space-time will require generalizing 't Hooft's equations 4.3 and 4.4 of [24] to the following wave functional,.

$$c_{i,f} \cdot \exp \left\{ - \int d\mathbf{x} \alpha \left[\phi_{Ci,f}(\mathbf{x}) - \phi_0(\mathbf{x}) \right]^2 \right\}, \text{ in terms of a solution using } \phi_0 \text{ as similar to the equivalence}$$

class 't Hooft [24, 25] is working with harmonic oscillators, then a connection is made to equivalence classes showing up in 't Hooft's 2006 manuscripts figure 2. The overlap with 't Hooft [24] comes from the fact that the functional given by $\Psi_{i,f} \left[\phi(\mathbf{x}) \right]_{\phi=\phi_{ci,cf}}$ is to a good approximation using a $\phi_{Ci,f}(\mathbf{x})$ with an evolution equation similar to what happens with damped harmonic oscillators with additional energy/information place into the damped harmonic oscillator. Notions of equivalence classes come in choices for $\phi_0(\mathbf{x})$ above. Whether the graviton can actually carry information will be considered in the next section..

IV. HOW MUCH INFORMATION NEEDS TO BE MAINTAINED TO PRESERVE THE COSMOLOGICAL CONSTANTS FROM COSMOLOGICAL CYCLE TO CYCLE?

No answer to this question has yet emerged. It is useful however to note that de La Peña [43] in 1997 proposed an order-of-magnitude estimate of information to derive a relation between Planck's constant (as a measure of the strength of the field fluctuations) and cosmological constants. If, for example, the fine structure constant has input parameter variance, as was explored by Livio et al, [44], with an explanation of why the fine structure constant has $\Delta\tilde{\alpha}/\tilde{\alpha} \leq 10^{-5} - 10^{-6}$ when redshift values are $z \sim 1.5$ compared to the present era and there is, for example, from QED, a proportional argument that $\tilde{\alpha} \equiv e^2/\hbar \cdot c$, is

$$\tilde{\alpha} \equiv e^2/\hbar \cdot c \equiv \frac{e^2}{d} \times \frac{\lambda}{hc} \tag{1.27}$$

However, extrapolating Eq. (1.27) after the turn on of the CMBR at $z \sim 1100$ 380 thousand years after the big bang is inappropriate, so instead, one should look for a similar statement for what $\tilde{\alpha} \equiv e^2/\hbar \cdot c$ would be at the onset of the big bang. The minimum length can be specified via use of the Snyder geometry

inequality [6] $\Delta x \geq [(1/\Delta p) + l_s^2 \cdot \Delta p] \equiv (1/\Delta p) - \alpha \cdot \Delta p$, to how know much ‘information’ is stored by $\tilde{\alpha} \equiv e^2/\hbar \cdot c$. Note that Δx is bounded by the Veneziano [40] specifications of minimum length.

$$\Delta x \equiv 10^\beta \cdot l_p \sim [(1/\Delta p) + l_s^2 \cdot \Delta p] \equiv (1/\Delta p) - \alpha \cdot \Delta p \quad (1.28)$$

Veneziano [40] wrote, $10^\beta \cdot l_p \equiv l_{string}$, where Planck length $l_p \equiv \sqrt{\hbar G/c^3}$, and this minimum length would be a way of specifying ‘transferrable information’ from the past to the present universe.

V. Conclusions

A. A reliable algorithm is needed in order to store information in a graviton

A way to obtain traces of information exchange, from prior to present universe cycles is finding a linkage between information and entropy. If such a parameterization can be found and analyzed, then Seth Lloyd’s [45] shorthand for entropy,

$$I = S_{total} / k_B \ln 2 = [\#operations]^{3/4} = [\rho \cdot c^5 \cdot t^4 / \hbar]^{3/4} \quad (1.29)$$

could be utilized as a way to represent information from a prior to the present universe. Eq. (1.3) is basic, but there could be different initial values of information placed into, \hbar based upon arguments at Eq. (1.29). If one views gravitons according to the idea refined by Beckwith [27] (from Ng [46]) -- that a counting algorithm for entropy is required -- then if the total number of gravitons in inflation is $n \sim 10^{20}$ gravitons $\approx 10^{20}$ entropy counts, Eq. (1.34) implies up to $\approx 10^{27}$ operations. If so, there is at least a one-to-one relationship between an operation and a bit of information, so a graviton has at least one bit of information.

B. Sensitivity limits for graviton detectors need to be improved

Note that the initial question posed in the beginning of the paper was: does the graviton have a mass? The stretch-out of a graviton wave, perhaps greater than the size of the solar system, gives, according to [4], an upper limit of a graviton mass of $\lambda_{graviton} > 300 \cdot h_0 kpc \Leftrightarrow m_{graviton} < 2 \times 10^{-29} h_0^{-1} eV$. I. e. a massively stretched graviton wave, at ultra-low frequency, may lead to a low mass limit. However, more careful and actually artificially narrow limits due to experimental searches, as presented by Buonanno [47] have narrowed the upper limit to $10^{-20} h_0^{-1} eV$. Giovanni’s [42] letter is important due to its embedding of an instanton structure in a 5 dimensional GR line element. The author chose a 5th dimensional conduit for information in a more classical model as a way to get about the singularity existing in four dimensions. One which Dr. Li and the author discussed at length in Chongqing that no instanton or instanton- anti instanton structure would survive a cosmic singularity.[48] However, it is possible that gravitons have higher frequencies [27], which would lead to smaller regions of space needed for high frequency gravitons nucleating in the beginning of the big bang. [27]. A counting algorithm, discussed in **Appendix D**, may be a way for information to be transferred from a prior universe to our present universe. Note that

in $\tilde{\alpha} \equiv e^2/\hbar \cdot c \equiv \frac{e^2}{d} \times \frac{\lambda}{hc}$, most of the information probably will be packed in the wavelength given as λ

which is part of how, $\tilde{\alpha}$ is defined and that the amount of information packed into this wavelength λ is packed into gravitons, as given in **Appendix C**. However, in order to obtain information put in information/

data for $\tilde{\alpha} \equiv \frac{e^2}{d} \times \frac{\lambda}{hc}$, experiments may require better sensitivity limits than what was assumed with

advanced LIGO. Advanced LIGO chooses [48], [49] $h \sim 10^{-23}$ as the maximum sensitivities for higher frequency GW. Eq. (1.35) is a formula for strain values for 1000 Hz GW.

$$h \sim \frac{\lambda}{Lb\sqrt{N\tau}} \sim 10^{-23} \quad (1.30)$$

Where L is interferometer length, N is the number of quanta per second at a beam splitter, and τ is the integration time. For LIGO systems the strain value given by Eq. (1.54) appears to be $h \sim 10^{-23}$. If one wishes to measure $\tilde{\alpha} \equiv e^2/\hbar \cdot c \equiv \frac{e^2}{d} \times \frac{\lambda}{hc}$, one may find that $h \sim 10^{-23}$ is inadequate [28, 48]

C. Further research questions for investigative inquiry and how to link our inquiry to the overall geometry of the universe

The problem of reconciling the existence of a graviton mass with quantum mechanics, in spin two particles usually having zero mass appears to be resolvable, and may show linkage between DE and DM in ways richer than the Chapygin gas models [11] By a problem to be solved, the author means the correspondence principle of quantum mechanics; i.e., that spin 2 particles have no mass.. Furthermore, the radius of the universe problem, given by Roos [11], will yield applications of the Friedmann equations used, once there are experimental criteria for determining the Hubble Parameter, and $\Omega \equiv \rho(t)/\rho_{critical}$.

$$r_U \equiv \frac{1}{H \cdot \sqrt{|\Omega - 1|}} \quad (1.31)$$

Combining experimental confirmation of Eq. (1.36) with observations and use of different choices for $H = \frac{\dot{a}}{a}$ and $\Omega \equiv \rho(t)/\rho_{critical}$ will be tied in, with analysis of the diagram of Figure 5 below




Figure 5. This figure from B. P. Abbott et al. [50] (2009) shows the relation between Ω_g and frequency.

The relation between Ω_g and the spectrum $h(v_g, \tau)$ is written by Grishchuk, [51], as

$$\Omega_g \approx \frac{\pi^2}{3} \left(\frac{v}{v_H} \right)^2 h^2(v, \tau), \quad (1.32)$$

The author looks to an interplay between Eq. (1.31) and Eq. (1.32) as a way to resolve questions as to if the universe is open or closed and shed some light as to the existence of relic GW. Answering questions about the interplay between Eq. (1.31) and Eq. (1.32) will use **Appendix D's** view of entropy to modify Eq. (1.32) inputs. Also, try to make refinements to Chapygin gas [52] joint DM-DE models. Furthermore, the ideas of this document are equivalent to finding experimental confirmation of the following relationship between non standard cosmology treatments of gravitons and the standard model [53]

$$\mathfrak{S} = -\frac{\mathcal{K}}{2} \cdot \sum_n \int d^4x h_{uv}^n \cdot T^{uv} \sim 1/M_{PL}^2 \quad (1.33)$$

The value of h_{uv}^n would come from further analysis of the sort started by **Appendix B** below, which would give the significance of Eq. (1.7) in terms of a projection of a modified version of a tower of KK

gravitons into the standard model T^{uv} stress energy tensor expression. Confirmation of Eq. (1.33) and also giving definite values as to Eq. (1.31) experimentally would go a long way toward turning GW astronomy into an exciting experimental science. Understanding Eq. (1.33) would be equivalent to, among other matters, getting some view of what gravity is, in terms of extra dimensions. Maartens [16] has related the scale of purported dimensions L , to potential behavior of gravity as of the form, when $d=1$, as can be interpreted by the model used for Eq. (1.7) above.

$$V \sim L^{-d} r^{-1} \approx L^{-1} r^{-1} \Big|_{d=1} \quad (1.34)$$

The key datum is to consider how Planck mass factors in with respect to dimensions, and extra dimensions. i.e. if extra dimensions, should they exist, mean $L \sim M_p^{-1}$ which appears to be possibly true, according to Maartens, who stated that collider and table top experiments tend toward $L \leq 0.1 \text{ mm}$. Should this be confirmed, then it would tie in with a re statement of Eq. (1.33) as

$$\mathfrak{S} = -\frac{\kappa}{2} \cdot \sum_n \int d^4 x h_{uv}^n \cdot T^{uv} \sim L^2 \quad (1.35)$$

What we are concerned with would be examining what can be said about the zeroth order interpretation of the KK graviton state, h_{uv}^0 Which according to traditional KK graviton theory is for a massless 4 dimensional 'graviton'. If there is no additional dimensions, and L is zero, then the only way to interpret the above is to set, when n is zero, that Eq. (1.35) is zero. If true, then examining the h_{uv}^0 mode needs to be interpreted along the lines of what [Kisselev, and Petrov](#) wrote [54], i.e. setting the mass of the graviton as $m_n = \tilde{\kappa} \cdot x_n$, with x_n the n th zero of the $J_1(\mathbf{x})$ Bessel function., with the zeroth mode of a graviton (corresponding to purely 4 dimensional gravitons) having no mass. If this is not true, then, if it has a slight mass, h_{uv}^0 will then have a different history corresponding to having a more complicated structure

$$(1.36)$$

Among other things, Eq. (1.36) would mean re doing the calculations done by Garriga, and Tanaka [55], and finding different values for perturbation on RS brane structure. Finally is the datum

$$w_{0de} = -1.08 < -1 \quad (1.37)$$

As brought up with respect to Eq. (1.7) and **Appendix A**, if one really has $w_{DE} \neq -1$ instead of $w_{DE} = -1$, the deceleration parameter as given by Eq. (A.1) of **Appendix A** would have to be revised, and made far more complicated. Aside from determining if $w_{DE} \neq -1$ for redshift $z < 1/2$ should be made a priority, there should be also investigations as to the implications of Eq. (1.36) as well as determining if the slope of the line, in Fig. 1, for $1 < z < 1.5$, which turns downward is due to $w_{DE} \neq -1$. Doing so would inevitably envision re confirming t'Hoofts [24,25] vision of embedding quantum mechanics in a larger theoretical construction, as mentioned at the end of **Appendix D**. This with respect to gravitons and kink-anti kink structures.

Appendix A: An analysis of how graviton mass assuming branes can influence expansion of the universe

Based on (1.10), with inputs from Friedmann equations, $\hbar = c = 1$ so then [14], the author claims that the following is appropriate to use

$$q = -\frac{\ddot{a}a}{\dot{a}^2} \equiv -1 - \frac{\dot{H}}{H^2} = -1 + \frac{2}{1 + \tilde{\kappa}^2 [\rho/m] \cdot (1+z)^4 \cdot (1 + \rho/2\lambda)} \quad (A.1)$$

The author found, in the course of numerical approximations, that $\tilde{\kappa}^2[\rho/m] \sim 2.03$, and $\rho/2\lambda \sim .0012$. The role of the ρ which if red shift is put in, becomes, if ρ_0 is today's density, which is low which does not vary much even for $z \sim 1.5$

$$\rho \equiv \rho_0 \cdot (1+z)^3 - \left[\frac{m_g \cdot (c=1)^6}{8\pi G(\hbar=1)^2} \right] \cdot \left(\frac{1}{14 \cdot (1+z)^3} + \frac{2}{5 \cdot (1+z)^2} - \frac{1}{2} \right) \quad (\text{A.2})$$

The upshot is for $z \sim .42$, or so, that a graviton mass is important in obtaining the value used for the brane world, i.e. $\rho/2\lambda \sim .0012$, which is needed for Fig. 1. Note that add on effect for ρ becomes useful in delineating when $q(z)$ goes to zero at $z \sim .42$, which is when data from galaxies one billion years ago indicate the rate of expansion of the universe is no longer slowing, but speeding up. Without $m_g \neq 0$, $q(z)$ goes to zero faster later on, in the simulation reported in Fig 1 on page 4

Apoendix B: Understanding modification of Maarten's higher dimensional DM candidtate, to incorporate operational 4 Dim DE

Starting with $m_{\tilde{n}}(\text{Graviton}) = \frac{\tilde{n}}{L}$, as given by Maarten [16], the author will build a case as to $m_{\tilde{n}}(\text{Graviton}) = \frac{\tilde{n}}{L} + 10^{-65}$ grams, and suggest how to confirm the 2nd equation. As given by Maarten, perturbative five dimensional graviton amplitudes can be Fourier expanded as:

$$f(x^a, y) = \sum_{\tilde{n}} [\exp(i\tilde{n}y/L)] \cdot f_{\tilde{n}}(x^a) \quad (\text{B.1})$$

leading to with regards to flat space geometry a d'Alembertain operator rendition of

$$\left[\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right] f_{\tilde{n}} = m_{\tilde{n}}^2 f_{\tilde{n}}, \quad m_{\tilde{n}} = \frac{\tilde{n}}{L} \quad (\text{B.2})$$

Here, f_0 is a four dimensional graviton mode, whereas the non zero \tilde{n} values correspond to effects of the five dimensional gravitational field upon a four dimensional brane.. In doing so, the massive modes are, of the form $L^{-1}, 2L^{-1}, \dots$ so if one has an infinite fifth dimension, the effect is to remove the separation between modes. $m_{\tilde{n}}(\text{Graviton}) = \frac{\tilde{n}}{L} + 10^{-65}$ grams $\approx m_{\tilde{n}} = \frac{\tilde{n}}{L} + \delta^+$ It is note worthy to note that the question of DM/ KK gravitons, and also the mass of the graviton not only has relevance to whether, higher dimensions are necessary in space time models, but also may be relevant to massive gravitons solving the DE puzzle. Rubakov [54] wrote that KK graviton representation as, after using the following normalization $\int \frac{dz}{a(z)} [h_m(z) \cdot h_{\tilde{m}}(z)] \equiv \delta(m - \tilde{m})$ where J_1, J_2, N_1, N_2 are different forms of Bessel

$$h_m(z) = \sqrt{m/k} \cdot \frac{J_1(m/k) \cdot N_2([m/k] \cdot \exp(k \cdot z)) - N_1(m/k) \cdot J_2([m/k] \cdot \exp(k \cdot z))}{\sqrt{[J_1(m/k)]^2 + [N_1(m/k)]^2}} \quad (\text{B.3})$$

This allegedly is for KK gravitons having an order of TeV magnitude mass $M_z \sim k$ (i.e. for mass values at .5 TeV to above a TeV in value) on a negative tension RS brane. What would be useful would be managing to relate this KK graviton, which is moving with a speed proportional to H^{-1} with regards to the negative tension brane with $h \equiv h_m(z \rightarrow 0) = \text{const} \cdot \sqrt{\frac{m}{k}}$ as an initial starting value for the KK graviton mass, before the KK graviton, as a 'massive' graviton moves with velocity H^{-1} along the RS dS

brane. If so, and if $h \equiv h_m(z \rightarrow 0) = \text{const} \cdot \sqrt{\frac{m}{k}}$ represents an initial state, then one may relate the mass of the KK graviton, moving at high speed, with the initial rest mass of the graviton, which in four space in a rest mass configuration would have a mass lower in value, i.e. of $m_{\text{graviton}}(4 - \text{Dim GR}) \sim 10^{-48} \text{ eV}$, as opposed to $M_X \sim M_{\text{KK-Graviton}} \sim .5 \times 10^9 \text{ eV}$. Whatever the range of the graviton mass, it may be a way to make sense of what was presented by Dubovsky, Flauger, Starobinsky, and Thackev [57] who argue for graviton mass using CMBR measurements, of $m_{\text{graviton}}(4 - \text{Dim GR}) \sim 10^{-20} \text{ eV}$. This can be conflated with Alves et. al. [5] arguing that non zero graviton mass may lead to acceleration of our present universe, in a manner usually conflated with DE, i.e. their graviton mass would be about $m_{\text{graviton}}(4 - \text{Dim GR}) \sim 10^{-48} \times 10^{-5} \text{ eV} \sim 10^{65} \text{ grams}$. Several different versions of what q should be will be offered as far as what the time dependence of terms in Eq. (1.10) actually is. Note that Roy Maartens has written as of 2004 and 2005 [16] that KK modes (graviton) satisfy a 4 Dimensional Klein – Gordon equation, with an effective 4 dim mass, $m_n(\text{Graviton}) = \frac{n}{L}$, with $m_0(\text{Graviton}) = 0$, and L as the stated ‘dimensional value’ of higher dimensions. The value $m_0(\text{Graviton}) \sim 10^{-65} - 10^{-60} \text{ gram}$ in value picked is very small, but ALMOST zero. If one is adding the small mass of $m_n(\text{Graviton}) = \frac{n}{L} + 10^{-65} \text{ grams}$, with $m_0(\text{Graviton}) \approx 10^{-65} \text{ grams}$, then the problem being worked with is a source term problem of the form given by Peskins [58] as of the type

$$\psi_n(x) \equiv \int d^3 p \cdot \frac{1}{(2\pi)^3} \cdot \frac{1}{\sqrt{2E_p}} \cdot \left\{ \left(a_p + \frac{i}{\sqrt{2E_p}} \cdot FT(m_0(\text{graviton})) \right) \exp(-ipx) + H.C. \right\} \quad (\text{B.4})$$

This is, using the language V.A. Rubakov [56] put up equivalent to writing,

$$\psi_m(x) \approx h_m(x) + \int d^3 p \cdot \frac{1}{(2\pi)^3} \cdot \left(\frac{1}{\sqrt{2E_p}} \right)^2 \cdot \{ (i \cdot FT(m_0(\text{graviton}))) \exp(-ipx) + H.C. \} \quad (\text{B.5})$$

I.e. how to interpret the quantity $FT(m_0(\text{graviton}))$ being the issue which will be covered in this document. If $m_0(\text{graviton})$ is a constant, then the expression (B.5) has delta functions.

APPENDIX C: INSTANTONS IN GENERAL RELATIVITY WITH REGARDS TO SPACE-TIME METRICS

The best, physically consistent models of GR-admissible solitons appears to be given by Belunski and Verdager [41], in work that ties in the instanton formulation for gravitation with specific metrics in space-time physics. In addition, Givannini [42] gives a kink- Ibanez and Verdager [59] suggest. Instantons only reach speeds up to light speed in nearly infinite distance travel. This instanton- anti instanton nearly fits with QM, and correspondence mass zero values for the graviton The author suggests that a kink-anti-kink construction for the graviton to avoid trouble with instantons traveling initially slowly as noted by Ibanez and Verdager [59]. Having a kink-anti kink construction for the graviton would avoid having low velocities for relic gravitons at their moment of creation. Updating Fig. 4, [41] gives an example of how to generaliz instantons from metric g, with $g \equiv \text{diag}\{t \cdot \exp(\phi), t \cdot \exp(-\phi)\}$ with

$$\phi \equiv d \cdot \ln t + \sum_{k=1}^s h_k \ln(\mu_k / t) \quad (\text{C.1})$$

The 2nd part of this equation corresponds to $\phi_+(z, \tau) = 4 \cdot \arctan \left(\exp \left\{ \frac{z + \beta \cdot \tau}{\sqrt{1 - \beta^2}} \right\} \right)$. Further work by

Belunski, and Verdager [41] yields instanton-anti-instanton solutions that are elaborations of Eq. (C1)

above. The tie in with Belunski and Verdauer [41] will be in similarities between Beckwith [30] use of

$$\arctan\left(\exp\left\{\frac{z+\beta\cdot\tau}{\sqrt{1-\beta^2}}\right\}\right)$$

and Belunski and Verdauer [41] using the formula $\arctan((bw)^\nu)$, as part of their solution containing instantons, as explained in Eq. (C.2). The justification [41] for instantons in cosmology is that a kink (anti kink) can be embedded in metrics which are used in General Relativity. As to the warning given by Ibanez, and Verdauer [59] that instantons travel at speeds very much smaller at the speed of light, in cosmology and reach peak velocities at ‘infinite’ distance from a source. It is important now to reference a kink (instanton) solution presented by Giovannini, [42] from a least action version of the Einstein – Hilbert action for ‘quadratic’ theories of gravity involving Euler- Gauss-Bonnet. Then Giovannini’s [42] equation 6 corresponds to

$$\phi = \tilde{v} + \arctan((bw)^\nu) \quad (C.2)$$

Givannini [42] represents of Eq. (C.2) as a kink, and makes references to an anti-kink solution, in Fig. 1 in

$$[42]. \text{ Furthermore the similarity between Eq. (C.2) and } \phi_+(z, \tau) = 4 \cdot \arctan\left(\exp\left\{\frac{z+\beta\cdot\tau}{\sqrt{1-\beta^2}}\right\}\right) \text{ in [30] is}$$

obvious. If $\arctan((bw)^\nu)$ overlaps in behavior with $\sum_{k=1}^s h_k \ln(\mu_k/t)$ the problem is amendable to analysis.

If a graviton is a kink-anti-kink combination, arising from a 5 dimensional line element, [37]

$$dS^2 = a(w) \cdot [\eta_{uv} dx^u dx^v - dw^2] \quad (C.3)$$

then how the graviton may be nucleated in the geometry specified by Eq. (C.3) is important. The stated ideas in **Appendix C** are presented to give equivalence between the degree for correspondence between QM, if assuming instanton- anti instanton pairs have thin walls. Box like shapes, as opposed to slopes, as seen in Figure 2 and Figure 4 indicate a fit with QM, and support a non linear theory involving instantons as being similar to results obtained with QM. All this is a better model than elaborate brane models of the graviton [60]. **Final** elaboration of this model will be similar to what was done by Ruutu et al. [61]. The hope is that work in instantons will lead to a refinement of DE/DM models, offering as big an advance as presented by Debnath once a kink-anti kink structure is linked to observations [62]

APPENDIX D : ENTROPY GENERATION VIA NG’S INFINITE QUANTUM STATISTICS

The author brings up entropy development due to the convergence of the instanton structure brought up in **Appendix C**, which is also in common with string theory. Furthermore, information counting ties in with information packing as brought up in the use of small graviton creation volume, V; for relic gravitons of a high frequency (short wave length) right after the big bang would be consistent Graviton volume V for nucleation is tiny, well inside inflation. So the log factor drops out of entropy S if V is chosen properly for both Eq. (D.1) and Eq. (D.2). Ng’s [46] result begins with modification of the entropy/partition function Ng used in an approximation of temperature, starting with early temperature $T \approx R_H^{-1}$ (R_H can be thought of as a representation of the region of space of the particles in question). Furthermore, assume that the volume of space is of the form $V \approx R_H^3$ and look at a numerical factor $N \sim (R_H/l_P)^2$, where the denominator is Planck’s length (on the order of 10^{-35} centimeters). We also specify a “wavelength” $\lambda \approx T^{-1}$. So the value of $\lambda \approx T^{-1}$ and of R_H are the same order of magnitude. Note Ng [46] changed conventional statistics: he outlined how to get $S \approx N$, or $S \approx \langle n \rangle$ (where $\langle n \rangle$ is graviton density). Begin with a partition function

$$Z_N \sim \left(\frac{1}{N!}\right) \cdot \left(\frac{V}{\lambda^3}\right)^N \quad (D.1)$$

This, according to Ng, leads to entropy of the limiting value of, if $S = (\log[Z_N])$ will be modified by

$$S \approx N \cdot (\log[V/N\lambda^3] + 5/2) \xrightarrow{\text{Ng-inf inite-Quantum-Statistics}} N \cdot (\log[V/\lambda^3] + 5/2) \approx N \quad (D.2)$$

But $V \approx R_H^3 \approx \lambda^3$, so unless N in Eq. (D.2) above is > 1 , S (entropy) would be < 0 , which is a contradiction. Now **Eq. (D.2)** is where [46] introduces removing the $N!$ term in Eq. (D.1) above, removing the expression of N inside the Log expression in Eq. (D.2). Since string theory involves instanton like structures, i.e. branes; the quantum infinite statistics, which Beckwith used [27] for gravitons is a limiting case of **Appendix C**. I.e. the two structures compliment each other. Further work in comparing Appendix C and Appendix D may refine Garriga's pioneering work done in 1994 [63] which treats instantons as classical solutions of the Euclidian equations of motion, in order to reconcile instanton treatment of gravitons as kink-anti kink components, i.e. non linear classical physics embedding a quantum spin two object, the spin two graviton. Eventually, the author hopes to put on a sound foundation what 'tHooft [24, 25] is doing with respect to his deterministic quantum mechanics and equivalence classes embedding quantum particle structures.. Doing so will hopefully answer some of the questions Kay raised about particle creation, and the limitations of the particle concept in curved and flat space, i.e. the global hyperbolic space time which is flat everywhere expect in a localized "bump" of curvature [64]

Bibliography-----

- [1] S. Hawking and G. . Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, 1973)
- [2] A. Ashtekar, private communications with the author, at the Penn State Cosmology center, 2007
- [3] C. Rovelli, *Quantum Gravity* (Cambridge University Press, 2005)
- [4] R. Goswami, P. S. Joshi, P. Singh, *Phys. Rev. Lett.* 96 (2006) 031302; <http://arxiv.org/abs/gr-qc/0506129>
- [5] E. Alves, O. Miranda. and J. de Araujo, arXiv: 0907.5190 (July 2009)
- [6] V. M. Battisti, *Phys. Rev.D* **79**, 083506 (2009)
- [7] G. Gabadadze and A. Gruzinov *Phys. Rev. D* **72**, 124007 (2005),
- [8] M. Visser, 1998, *General Relativ. Gravit.*, 30, 1717
- [9] H. Van Dam and M. Veltman , *Nuc Phys B* , **22**, 397 (1970)
- [10] C. M. Will, 2006, <http://relativity.livingreviews.org/Articles/lrr-2006-3/>
- [11] M. Roos, *Introduction to Cosmology, 3rd edition* (Wiley Interscience , 2003)
- [12] M. Maggiore, *Gravitational Waves, Volume 1: Theory and Experiment*, (Oxford University press,2008)
- [13] G. Fuller and C. Kishimoto ,*Phys. Rev. Lett.* **102**, 201303 (2009)
- [14] A.Beckwith,,<http://vixra.org/abs/0910.0057>, version 2 (2009)
- [15] A. Friedmann , *Class. Quantum. Grav* **31**, Number 12 / December, 1999, pages 2001-2008
- [16] R. Maartens , (2004), <http://www.livingreviews.org/lrr-2004-7> ; from the conference *The physics of the Early Universe* , editor Papantronopoulos, (Lect. notes in phys., Vol **653**, Springer Verlag, **2005**), R, Maartens *Brane world cosmology*, pp **213-247**
- [17] V. Mukhanov, *Physical Foundations of Cosmology* (Cambridge University Press, 2005)
- [18] K. Becker, M. Becker, J. H. Schwartz ,*String theory and M Theory , a Modern Introduction*, (Cambridge University Press, 2007)
- [19] J. Martin, *Inflationary Perturbations:The cosmological Schwinger effect* ,pages 193-241, Lecture notes in Physics, 738, editors M. Lemonine, J. Martin, P. Peters, (Springer - Verlag 2008)
- [20]G. Grossing, *Phys Lett A*, **296** , 1, 2002
- [21] Victor Taveras, *Phys.Rev.D***78**:064072,2008; <http://arxiv.org/abs/0807.3325>
- [22] S.Bashinsky, <http://arxiv.org/abs/astro-ph/0505502>, 4 May 2005
- [23] Dim. Valev., *Aerospace Res. Bulg.* 22:68-82, 2008; <http://arxiv.org/abs/hep-ph/0507255>
- [24]] G. 't Hooft., in *Beyond the Quantum*, edited by Th. M. Nieuwenhuizen et al. (World Press Scientific 2006), http://arxiv.org/PS_cache/quant-ph/pdf/0604/0604008v2.pdf, (2006)
- [25] G. 't Hooft, http://arxiv.org/PS_cache/quant-ph/pdf/0212/0212095v1.pdf (2002)
- [26]T. Rothman, S. Boughn., *Found.Phys.* 36 (2006) 1801-1825, <http://arxiv.org/abs/gr-qc/0601043>
- [27] A.Beckwith, **accepted contribution** to the Rencontres de Blois conference, 2009, <http://vixra.org/abs/0908.0109>
- [28] F. Li, N. Yang, Z. Fang, R. M. L. Baker Jr., G. V. Stephenson, and H. Wen, *Phys. Rev. D* **80**, 064013 (2009)
- [29] S. Mukohyama, *JCAP* 0906:001,2009; <http://arxiv.org/abs/0904.2190> (2009)
- [30] R. Brandenberger, *Phys. Rev. D* **80**, 043516 (2009); <http://arxiv.org/abs/0904.2835> (2009)
- [31] M. Visser, *Who's Afraid of Lorentz Symmetry Breaking* as invited Plenary talk at the ACGRC 5 meeting, December 16, 2009
- [32] A..Beckwith, *Classical and Quantum Models of Density Wave Transport, a comparative study*, PhD dissertation University of Houston , December 2001
- [33] M. Visser, private discussions with the author December 17 at ACGRG 5, in Christchurch, New Zealand,
- [34] A. Beckwith, *Mod.Phys.Lett.* **B20** (2006) 849-861, <http://arxiv.org/abs/math-ph/0411045>
- [35] J.H. Miller, J. Richards, R.E. Thorne, W.G.Lyons and J.R. Tucker, *Phys.Rev. Lett.* 55, 1006 (1985)
- [36] L. Crowell, *Quantum Fluctuations of Space Time* (World Press Scientific, , 2005)
- [37] V. Mukhanov, *Physical elements of Cosmology* (Cambridge University press, 2005)
- [38] Lecture by A. Beckwith, at Chongqing University, November 10, 2009

- [39] C. Kiefer, *Quantum Gravity*, (2nd edition, Oxford Science publications 2007)
- [40] G. Veneziano pp 134-180“Classical and Quantum Gravity from String Theory”, from the volume: ‘*Classical and Quantum Gravity*’, editors M.C. Bento, O. Bertolami, J.M. Mourao, and R.F. Picken, World press Scientific, Singapore, 1993
- [41] V. Belunski, and E. Verdaguer, *Gravitational Solitons*: (Cambridge University press 2001)
- [42] M. Giovannini, *Class. Quantum. Grav.* 23, 2006, L73-80
- [43] L. de La Peña and A. M. Cetto, *Foundations of Physics Letters*,_issue_Volume 10, Number 6 / December, 1997, pages 591-598
- [44] M. Livio and M. Stiavelli, (1999), *The Astrophysical Journal*, 507:L13–L15, 1998 November 1, <http://www.iop.org/EJ/article/1538-4357/507/1/L13/985313.text.html>
- [45] S.Lloyd, *Phys. Rev. Lett.* **88**, 237901 (2002)
- [46] Ng, Y.Jack Entropy 2008, 10(4), 441-461; DOI: 10.3390/e10040441
- [47]] A. Buonanno, “Gravitational waves”, pp 10-52, from the Les Houches Section LXXXVI, ‘*Particle physics and Cosmology, the fabric of space-time*’ which has F. Bernardeau, C. Grojean, and J. Dalibard as editors
- [48] Joint conclusion of the author, and Dr. Fangyu Li, at talks between the two principals in Chongqing University, November 2009, after ADM 50
- [49] R. Weiss, private discussions with the author, at ADM 50
- [50] B.P. Abbott et al., *Nature* **460**, 990-993 (2009)
- [51] L. P. Grishchuk, *Lect. Notes Phys.* 562, 167 (2001)
- [52] U. Debnath, S. Chakraborty, *Int. Journal of Theor. Phys.* **47**, No. 10, p 2663,(2008).
- [53] U. Sarkar, *Particle and Astroparticle physics*, Taylor and Francis,- Series in High Energy physics, Cosmology and Gravitation – **2008**
- [54] A. Kisselev, and V. Petrov, *Czechoslov. J. Phys.* **56** (2006) A91-A97 ; <http://arxiv.org/abs/hep-ph/0511018>.
- [55] [J. Garriga](#), [T. Tanaka](#) *Phys.Rev.Lett.* **84** (2000) 2778-2781; <http://arxiv.org/abs/hep-th/9911055>.
- [56] V. Rubakov, *Classical Theory of Gauge Fields*, Princeton University press, 2002.
- [57] S. Dubovsky, R. Flauger, A. Starobinsky, I. Tkachev,,report UTTG-06-09, TCC-23-09, <http://arxiv.org/abs/0907.1658>.
- [58] M. Peskins, D. Schroeder, D. *An introduction to Quantum Field Theory*, Westview Press, (1995).
- [59] Ibanez, J., and Verdaguer, E. (1985). *Phys. Rev D*, **31**, 251.
- [60] J. Leach, and W. Lesame, arXIV gr-qc/0502109 v3 26 Apr 2005.
- [61] V. Ruutu,. .V. Eltsov, A. Gill, et. al. *Nature* **382**, 334-336 (25 July 1996).
- [62] U. Debnath and S. Chakraborty, *Int. Journal of Theor. Phys.* **47**, No. 10, page 663 (2008).
- [63] J. Garigga, *Phys Rev. D* **49**, volume 12, pp 6327-6342; <http://128.84.158.114/abs/hep-ph/9308280>.
- [64] B. Kay, *Quantum Field theory in Curved Spacetime*, pp 180-190, in the Encyclopedia of Mathematical physics, Vol 5, *General Relativity; Quantum Gravity; String theory and M. Theory*, With J-P Francoise, G. Naber, and T. Tsun as editors