

New math. for physical applications

S. Kalimuthu

2/349 Kanjampatti B.O, Pollachi Via, Tamil Nadu 642003, India

Email: sennimalaikalimuthu@outlook.com

Mobile: + 91 850 899 1577

Abstract

After establishing the fundamental physics prizes, Yuri Milner said: “*Unlike the Nobel in physics, the Fundamental Physics Prize can be awarded to scientists whose ideas have not yet been verified by experiments, which often occurs decades later. Sometimes a radical new idea “really deserves recognition right away because it expands our understanding of at least what is possible.”*”. Keeping this mind the author formulated two spherical geometrical theorems which may applied for the studies and probes of fundamental particles, quantum gravity, gravitational waves ,dark matter and dark energy.

Keywords: Non Euclidean math., New Geometry, Physical Applications.

PACS:

[04.20.-q](#), [04.50.-h](#), [06.20.Jr](#) (all), [01.70.+w](#), [42.50.-p](#), [04.20.-](#), [04.50.+h](#), [98.80.Jk](#) [04.30.-w](#)


Introduction

The Pauli exclusion principle was postulated in an attempt to explain some of the properties of electrons in an atom. This principle says that in a closed system, no two electrons can occupy the same state..Heisenberg’s uncertainty principle states that the position and momentum of a particle cannot be simultaneously measured with arbitrarily high precision. Special relativity applies only to cases in which objects are moving at a uniform velocity. General relativity, however, is applicable to all forms of accelerated motion. This theory of general relativity arose from Einstein's principle of equivalence. Einstein formulated this principle by examining a given mass in two different states. Einstein’s equivalence principle is any of several related concepts dealing with the equivalence of gravitational and inertial mass, and to Albert Einstein's observation that the gravitational "force" as experienced locally while standing on a massive body (such as the Earth) is actually the same as the *pseudo-force* experienced by an observer in a non- inertial (accelerated) frame of reference. Like these easy and brief principles, the author proposes the following spherical geometrical theorems.


Theorem 1 : There exists a spherical triangle whose interior angle sum adds to 360^0

Theorem 2 : There exists a spherical triangle whose interior angle sum adds to 540^0

Construction



Spherical Figure 1



Spherical Figure 2

First Proof for theorem 1

In spherical figure 1. Consider NB , WE and EN as the three sides of triangle NEW. WE is the equator of spherical figure 1 and both EN and WN are perpendiculars to WE. Since the angle WNE is a straight angle, we get that the sum of the interior angles of spherical triangle WNE is equal to 360 degrees. And hence the proof.

Second Proof for theorem 1

In spherical figure 2, consider WN , NE and EW as the three sides of spherical triangle WNE. Since the angles WNE , NEW and EWN are straight angles, we obtain that the sum of the interior angles of spherical triangle WNE is 540 degrees. And hence the proof.

Let y , z and m are three distinct spherical triangles. Since the interior angle sum of a spherical triangle is more than 180 degree,

$$\text{Let us assume, } y + z = 360^0 + a \quad (1)$$

$$m+z = 360^0 + b \quad (2)$$

$$(1) - (2) \text{ gives, } y - m = a - b \quad (3)$$

$$\text{Squaring (3), } y^2 + m^2 - 2my = a^2 + b^2 - 2ab \quad (3a)$$

Multiplying (1) and (2),

$$y(m+z) + z(m+z) = 360^{02} + 360^0 b + 360^0 a + ab \quad (4)$$

Adding (3a) and (4) we get that,

$$\begin{aligned} y(m+z+y-2m) + m(m+z) + (z+b)(z-b) \\ = a(a-b + 360^0) + 360^0(b + 360^0) \end{aligned}$$

Applying (1) in the first factor, and (2) in the second and third factors of LHS

And putting (2) in the second factor of RHS we have,

$$\begin{aligned} y(360^0 + a - m) + m(m+z) + (z+b)(360^0 - m) \\ = a(a-b + 360^0) + 360^0(m+z) \end{aligned}$$

$$\text{i. e } y(360^0 + a - m) + (m - 360^0)(m+z) = a(a-b + 360^0)$$

$$\text{Putting (3) in RHS, } y(360^0 + a - m) + (m - 360^0)(m+z) = a(y - m + 360^0)y$$

$$(360^0 + a - m - a) + m(m+z+a) = 360^0(a+m+z)$$

$$\text{i.e } y(360^0 - m) + (m + z + a)(m - 360^0) = 0$$

$$\text{i.e } (m - 360^0)(m + z + a - y) = 0$$

$$\text{Assuming (3) in the second factor, } e(m - 360^0)(z + a + b - a) = 0$$

$$\text{i.e } (m - 360^0)(z + b) = 0$$

$$\text{i.e } m = 360^0 \tag{5}$$

So, (5) establishes our first theorem.

Third Proof for theorem 1

Since angles WAB , ABC , and BCE are straight angles they are all each equal to 180 degree.

$$\text{Let } v \text{ be the value of this 180 degree.} \tag{1}$$

$$\text{Let angle } WNB = s, \text{ } ANC = t \text{ and } WBE = u \tag{2}$$

Assuming (1) and (2) and adding we get that,

$$x + y = 2v + s \tag{3}$$

$$y + z = 2v + t \tag{4}$$

$$z + m = 2v + u \tag{5}$$

$$(3) - (4) \text{ gives, } x + t = z + s \tag{6}$$

$$(4) - (5) \text{ yields, } y + u = m + t \tag{7}$$

Squaring (3)

$$x^2 + y^2 + 2xy = 4v^2 + s^2 + 4vs \tag{3a}$$

$$\text{Squaring (7) } m^2 + t^2 + 2mt = y^2 + u^2 + 2yu \tag{7a}$$

Adding (3a) and (7a) we get that,

$$\begin{aligned} (x + u)(x - u) + (m + 2v)(m - 2v) + (t + s)(t - s) \\ + 2y(x - u) + 2mt - 4vs = 0 \end{aligned}$$

$$\begin{aligned} \text{i. e } (x - u) [x + u + 2y] + (m + 2v)(m - 2v) + \\ (t + s)(t - s) + 2mt - 4vs = 0 \end{aligned}$$

Replacing $x + y$ by $2v + s$, $y + u$ by $m + t$, $m - 2v$ by $u - z$ and $t - s$ by $z - x$ we have,

[See eqns. (3) , (5) , (6) and (7)]

$$\begin{aligned} (x - u) [2v + s + m + t] + (m + 2v)(u - z) + \\ (t + s)(t - s) + 2mt - 4vs = 0 \end{aligned}$$

Rearranging, $(m + 2v)[x - u + u - z] + (t + s)[t - s + x - u] + 2mt - 4vs = 0$

$$\text{i. e } (m + 2v)[x - z] + (t + s)[t - s + x - u] + 2mt - 4vs = 0$$

Substituting $s - t$ for $x - z$ and $s + z$ for $x + t$ [See eqn.(6)] we have,

$$(m + 2v)[s - t] + (t + s)[z - u] + 2mt - 4vs = 0$$

$$\text{i.e } t(z - u + 2m - m - 2v) + s(z - u - 4v - m - 2v) = 0$$

Replacing $m + z$ by $2v + u$ [see eqn. (5)] we obtain, $2m + 4v = 0$

$$\begin{aligned} \text{i} \\ \text{i.e } m + 2v = 0, \text{ i.e } m = -2v \end{aligned} \tag{8}$$

It is well known that in geometry minus theta represents the vertically opposite angles. Since vertically opposite angles are equal it implies

$$\text{from (8) that } m = 2v. \tag{9}$$

Comparing (9) and (2) we get that the sum of the interior angles of spherical triangle NCE is equal to four right angles.

(10)

Eqn. (10) proves our first theorem.

Fourth Proof for theorem 1

In the above spherical figure, the angles at W , A , B , C and E are right angles. s , t and u

denotes triangles WNB, ANC and BNE respectively

Considering straight angles WAB , ABC and BCE and adding,

$$x + y = 2v + s \quad (1)$$

$$y + z = 2v + t \quad (2)$$

$$m + z = 2v + u \quad (3)$$

(2) – (3) gives,

$$m + t = y + u \quad (4)$$

Squaring (1),

$$x^2 + y^2 + 2xy = 4v^2 + s^2 + 4vs \quad (5)$$

Squaring (4),

$$m^2 + t^2 + 2mt = y^2 + u^2 + 2yu \quad (6)$$

Adding (5) and (6),

$$x(x+y)+xy+(m+2v)(m-2v)+(t+s)(t-s)+2mt-u(y+u)-yu-4vs=0 \text{ i.e } x(x+y)+ y(x-u)+2mt-u(y+u)+2mt-4vs + (t+s)(t-s)+ =0$$

Applying (3) in the first factor, and (6) in the third factor we have, $x(2v+s)+y(x-u)+2mt-um-ut-4vs + (t+s)(t-s)+=0$

$$2v(x-s)+s(x-2v)+y(x-u)+m(t-u)+t(m-u) + (t+s)(t-s)+=0$$

From (3) we have, $x - s = 2v - y$ and $x - 2v = s - y$.Assuming these values in the above relations,

$$2v (2v - y) + s (s - y))y(x-u)+m(t-u)+t(m-u) + (t+s)(t-s)+=0$$

$$\text{i.e } y (x - u - s - 2v) + 4v^2 + s^2 +m(t-u)+t(m-u) + (t+s)(t-s)+=0$$

Assuming (3) in the first factor, $-y(y + u) + 4v^2 + s^2 + m(t-u) + t(m-u) + (t+s)(t-s) = 0$

Rearranging, $-y(y + u) + 4v^2 + m(t-u) + t(m-u + t + s) + s(s - s - t)$

Putting (4) in the first factor,

$$-y(m + t) + 4v^2 + m(t-u) + t(m-u + t + s) - st = 0$$

$$i. e + 4v^2 + m(t-u - y) + t(m-u + t + s - y - s) = 0$$

Once again assuming (4), $4v^2 - m^2 = 0$

$$i. e m = 2v \tag{7}$$

And (7) is the fourth proof of our first theorem

Proof of theorem 2

In spherical figure 2, consider WN, NE and EW as the three sides of spherical triangle WNE. Since the angles WNE, NEW and EWN are straight angles, we obtain that the sum of the interior angles of spherical triangle WNE is 540 degrees. And hence the proof

Discussion

Let us recall that after the publication of non Euclidean math. work Riemann concluded: "Here after it is up to physicists to apply my findings." Similarly I request the research community to apply my results to theoretical physics and cosmology. There are many burning problems in physics such like quantum gravity, dark matter, dark energy and pre big bang phenomena. When Lobachevsky published his first non Euclidean math in 1824, the whole research community did NOT approve it. They have remarked non Euclidean math. only a mathematical trick. But this concept was widely applied in Einstein's special relativity after 1905 and the formulae of hyperbolic geometry are being applied to study the atomic properties in quantum physics. Riemann's second non Euclidean math. was published in 1854. It was assumed to formulate general relativity in 1915. The hyperbolic math. had to wait for 81 years and the elliptic had to wait for 61 years for recognition and application. Similarly my new findings will be very useful in theoretical physics and cosmology.

There was a trouble with Maxwell's equations. A deep analysis of his equations predicted that light is an electromagnetic wave. And PAM Dirac also encountered such a physical phenomenon. Then Dirac's equations revealed that there exists anti - particles. Similarly the author's finding that the sum of the interior of spherical triangle NCE is equal to four straight

angles will unlock some of the mysteries of cosmological problems such like gravitational waves, dark energy and dark matter. The applications of eqn. (7) to spherical trigonometry and differential equations will predict new cosmological phenomena.

Algebra is the extension of number theory. It occupies almost all the areas of science, technology, and administration. The famous French mathematician used to tell time and again that As long as algebra and geometry have been separated, their progress have been slow and their uses limited; but when these two sciences have been united, they have lent each mutual forces, and have marched together towards perfection. And once Einstein proposed to the scientific community to put all the equations of physics in algebra. These two says are the foundational guide lines for the author for the preparation of this paper. Future probes and studies will surely create a new field of spherical geometry & trigonometry.

TP:

Kepler's law of planetary motions, Galileo's inventions, Einstein's special and general relativity theories, De Broglie's matter waves hypothesis, Pauli's exclusive principle, Heisenberg's uncertainty principle, Fractals geometric idea, Lobachevsky's non Euclidean geometric concept, Riemann's non Euclidean geometric theory, Peter Higg's Bosons papers and many other ground breaking inventions were initially NOT accepted/approved by the research community. Later on these findings were gradually agreeable to the scientists. *In quasicrystals, we find the fascinating mosaics of the Arabic world reproduced at the level of atoms: regular patterns that never repeat themselves. However, the configuration found in quasicrystals was considered impossible, and Dan Shechtman had to fight a fierce battle against established science. The Nobel Prize in Chemistry 2011 has fundamentally altered how chemists conceive of solid matter. The research community, Editors and referees of professional journals, research institutes and chemistry departments completely ignored, insulted and avoided and Dan Shechtman. But he did not bother about this and continued his research and finally won the Nobel prize In this work, by applying algebra to geometry, the author has found challenging results.*

References

- [1] http://mathforum.org/library/topics/elliptic_g/
- [2] <http://www.sjsu.edu/faculty/watkins/sphere.htm>
- [3] <http://www.alibris.com/search/books/isbn/9781140723271>
- [4] <http://www.euclideanspace.com/maths/geometry/space/nonEuclid/spherical/index.htm>
- [5] <http://archive.org/details/textbookonspheri033273mbp>
- [6] http://en.wikipedia.org/wiki/Spherical_trigonometry
- [7] www.wilbourhall.org/pdfs/spherical_trigonometry2.pdf