

Reformulation of relativistic quantum mechanics and field theory equations with space - time sedenions

Victor L. Mironov, Sergey V. Mironov

Institute for physics of microstructures RAS, 603950, Nizhniy Novgorod, GSP-105, Russia

e-mail: mironov@ipmras.ru

(Submitted 13 January 2014, revised 25 February 2014)

We present an alternative type of sixteen-component hypercomplex scalar-vector values named “space-time sedenions”, generating associative noncommutative space-time Clifford algebra. The generalization of relativistic quantum mechanics and field theory on the basis of sedenionic space-time operators and hypercomplex wave functions is discussed.

1. Introduction

The multicomponent hypercomplex numbers and multivectors are widely used for the reformulation of the equations of quantum mechanics and field theory. The early generalization of quantum mechanics and electrodynamics were made on the basis of quaternions [1-5] with scalar-vector structure. The next step was taken on the basis of biquaternions and octonions [6-11] enclosing scalar, pseudoscalar, vector and pseudovector components. These structures take into account the space symmetry with respect to the spatial inversion. However, a consistent relativistic approach requires taking into consideration full time and space symmetries that leads to the sixteen-component space-time algebras.

The well known sixteen-component hypercomplex numbers, sedenions, are obtained from octonions by the Cayley-Dickson extension procedure [12,13]. In this case the sedenion is defined as

$$S = O_1 + O_2 \mathbf{e}, \quad (1)$$

where O_i is an octonion and the parameter of duplication \mathbf{e} is similar to imaginary unit $\mathbf{e}^2 = -1$. The algebra of sedenions has the specific rules of multiplication. The product of two sedenions

$$S_1 = O_{11} + O_{12} \mathbf{e},$$

$$S_2 = O_{21} + O_{22} \mathbf{e},$$

is defined as

$$S_1 S_2 = (O_{11} + O_{12} \mathbf{e})(O_{21} + O_{22} \mathbf{e}) = (O_{11} O_{21} - \bar{O}_{22} O_{12}) + (O_{22} O_{11} + O_{12} \bar{O}_{21}) \mathbf{e}, \quad (2)$$

where \bar{O}_i is conjugated octonion. The sedenionic multiplication (2) allows one to introduce a well defined norm of sedenion. However such procedure of constructing the higher hypercomplex numbers leads to the fact that the sedenions as well as octonions generate normed but nonassociative algebra [14-16]. This greatly complicates the use of the Cayley-Dickson sedenions in the physical applications.

Recently we have developed an alternative approach to constructing the multicomponent values based on our scalar-vector conception realized in associative eight-component octons [17-19] and sixteen-component sedeons [20-23]. In particular, we have demonstrated the method, which allows one to reformulate the equations of relativistic quantum mechanics and field theory on the basis of sedeonic space-time operators and scalar-vector wave functions. In this paper we present an alternative version of the sixteen-component associative space-time algebra and demonstrate some of its application to the generalization of the field theory equations.

2. Sedenionic space-time algebra

It is known, the quaternion is a four-component object

$$\widehat{q} = q_0 \mathbf{a}_0 + q_1 \mathbf{a}_1 + q_2 \mathbf{a}_2 + q_3 \mathbf{a}_3, \quad (3)$$

where components q_ν (Greek indexes $\nu = 0, 1, 2, 3$) are numbers (complex in general), $\mathbf{a}_0 \equiv 1$ is scalar units and values \mathbf{a}_m (Latin indexes $m = 1, 2, 3$) are quaternionic units, which are interpreted as unit vectors. The rules of multiplication and commutation for \mathbf{a}_m are presented in Table 1. We introduce also the space-time basis $\mathbf{e}_t, \mathbf{e}_r, \mathbf{e}_{tr}$, which is responsible for the space-time inversions. The indexes \mathbf{t} and \mathbf{r} indicate the transformations (\mathbf{t} for time inversion and \mathbf{r} for spatial inversion), which change the corresponding values. The value $\mathbf{e}_0 \equiv 1$ is a scalar unit. For convenience we introduce numerical designations $\mathbf{e}_1 \equiv \mathbf{e}_t$ (time scalar unit); $\mathbf{e}_2 \equiv \mathbf{e}_r$ (space scalar unit) and $\mathbf{e}_3 \equiv \mathbf{e}_{tr}$ (space-time scalar unit). The rules of multiplication and commutation for this basis we choose similar to the rules for quaternionic units (see Table 2).

Table 1.

| | | | |
|----------------|-----------------|-----------------|-----------------|
| | \mathbf{a}_1 | \mathbf{a}_2 | \mathbf{a}_3 |
| \mathbf{a}_1 | -1 | \mathbf{a}_3 | $-\mathbf{a}_2$ |
| \mathbf{a}_2 | $-\mathbf{a}_3$ | -1 | \mathbf{a}_1 |
| \mathbf{a}_3 | \mathbf{a}_2 | $-\mathbf{a}_1$ | -1 |

Table 2.

| | | | |
|----------------|-----------------|-----------------|-----------------|
| | \mathbf{e}_1 | \mathbf{e}_2 | \mathbf{e}_3 |
| \mathbf{e}_1 | -1 | \mathbf{e}_3 | $-\mathbf{e}_2$ |
| \mathbf{e}_2 | $-\mathbf{e}_3$ | -1 | \mathbf{e}_1 |
| \mathbf{e}_3 | \mathbf{e}_2 | $-\mathbf{e}_1$ | -1 |

Note that the unit vectors $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$ and the space-time units $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ generate the anticommutative algebras:

$$\begin{aligned} \mathbf{a}_n \mathbf{a}_m &= -\mathbf{a}_m \mathbf{a}_n, \\ \mathbf{e}_n \mathbf{e}_m &= -\mathbf{e}_m \mathbf{e}_n, \end{aligned} \quad (4)$$

for $\mathbf{n} \neq \mathbf{m}$, but $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ commute with $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$:

$$\mathbf{e}_n \mathbf{a}_m = \mathbf{a}_m \mathbf{e}_n, \quad (5)$$

for any \mathbf{n} and \mathbf{m} . Then we can introduce the sixteen-component space-time sedenion \tilde{V} in the following form:

$$\begin{aligned} \tilde{V} &= \mathbf{e}_0 (V_{00} \mathbf{a}_0 + V_{01} \mathbf{a}_1 + V_{02} \mathbf{a}_2 + V_{03} \mathbf{a}_3) + \mathbf{e}_1 (V_{10} \mathbf{a}_0 + V_{11} \mathbf{a}_1 + V_{12} \mathbf{a}_2 + V_{13} \mathbf{a}_3) \\ &+ \mathbf{e}_2 (V_{20} \mathbf{a}_0 + V_{21} \mathbf{a}_1 + V_{22} \mathbf{a}_2 + V_{23} \mathbf{a}_3) + \mathbf{e}_3 (V_{30} \mathbf{a}_0 + V_{31} \mathbf{a}_1 + V_{32} \mathbf{a}_2 + V_{33} \mathbf{a}_3). \end{aligned} \quad (6)$$

The sedenionic components $V_{\nu\mu}$ are numbers (complex in general). Introducing designation of scalar and vector values in accordance with the following relations

$$\begin{aligned} V &= \mathbf{e}_0 V_{00} \mathbf{a}_0, \\ \vec{V} &= \mathbf{e}_0 (V_{01} \mathbf{a}_1 + V_{02} \mathbf{a}_2 + V_{03} \mathbf{a}_3), \\ V_t &\equiv V_1 = \mathbf{e}_1 V_{10} \mathbf{a}_0, \\ \vec{V}_t &\equiv \vec{V}_1 = \mathbf{e}_1 (V_{11} \mathbf{a}_1 + V_{12} \mathbf{a}_2 + V_{13} \mathbf{a}_3), \end{aligned} \quad (7)$$

$$\begin{aligned}
V_r &\equiv V_2 = \mathbf{e}_2 V_{20} \mathbf{a}_0, \\
\vec{V}_r &\equiv \vec{V}_2 = \mathbf{e}_2 (V_{21} \mathbf{a}_1 + V_{22} \mathbf{a}_2 + V_{23} \mathbf{a}_3), \\
V_{tr} &\equiv V_3 = \mathbf{e}_3 V_{30} \mathbf{a}_0, \\
\vec{V}_{tr} &\equiv \vec{V}_3 = \mathbf{e}_3 (V_{31} \mathbf{a}_1 + V_{32} \mathbf{a}_2 + V_{33} \mathbf{a}_3),
\end{aligned}$$

We can represent the sedenion in the following scalar-vector form:

$$\tilde{V} = V + \vec{V} + V_t + \vec{V}_t + V_r + \vec{V}_r + V_{tr} + \vec{V}_{tr}. \quad (8)$$

Thus, the sedenionic algebra encloses four groups of values, which are differed with respect to spatial and time inversion.

- (1) Absolute scalars (V) and absolute vectors (\vec{V}) are not transformed under spatial and time inversion.
- (2) Time scalars (V_t) and time vectors (\vec{V}_t) are changed (in sign) under time inversion and are not transformed under spatial inversion.
- (3) Space scalars (V_r) and space vectors (\vec{V}_r) are changed under spatial inversion and are not transformed under time inversion.
- (4) Space-time scalars (V_{tr}) and space-time vectors (\vec{V}_{tr}) are changed under spatial and time inversion.

Further we will use the symbol 1 instead units \mathbf{a}_0 and \mathbf{e}_0 for simplicity. Introducing the designations of scalar-vector values

$$\begin{aligned}
\vec{V}_0 &= V_{00} + V_{01} \mathbf{a}_1 + V_{02} \mathbf{a}_2 + V_{03} \mathbf{a}_3, \\
\vec{V}_1 &= V_{10} + V_{11} \mathbf{a}_1 + V_{12} \mathbf{a}_2 + V_{13} \mathbf{a}_3, \\
\vec{V}_2 &= V_{20} + V_{21} \mathbf{a}_1 + V_{22} \mathbf{a}_2 + V_{23} \mathbf{a}_3, \\
\vec{V}_3 &= V_{30} + V_{31} \mathbf{a}_1 + V_{32} \mathbf{a}_2 + V_{33} \mathbf{a}_3,
\end{aligned} \quad (9)$$

we can write the sedenion (6) in the following compact form

$$\tilde{V} = \vec{V}_0 + \mathbf{e}_1 \vec{V}_1 + \mathbf{e}_2 \vec{V}_2 + \mathbf{e}_3 \vec{V}_3. \quad (10)$$

On the other hand, introducing designations of space-time sedenion-scalars

$$\begin{aligned}
V_0 &= (V_{00} + \mathbf{e}_1 V_{10} + \mathbf{e}_2 V_{20} + \mathbf{e}_3 V_{30}), \\
V_1 &= (V_{01} + \mathbf{e}_1 V_{11} + \mathbf{e}_2 V_{21} + \mathbf{e}_3 V_{31}), \\
V_2 &= (V_{02} + \mathbf{e}_1 V_{12} + \mathbf{e}_2 V_{22} + \mathbf{e}_3 V_{32}), \\
V_3 &= (V_{03} + \mathbf{e}_1 V_{13} + \mathbf{e}_2 V_{23} + \mathbf{e}_3 V_{33}),
\end{aligned} \quad (11)$$

we can write the sedenion (6) as

$$\tilde{V} = V_0 + V_1 \mathbf{a}_1 + V_2 \mathbf{a}_2 + V_3 \mathbf{a}_3, \quad (12)$$

or introducing the sedenion-vector

$$\vec{V} = \vec{V} + \vec{V}_t + \vec{V}_r + \vec{V}_{tr} = V_1 \mathbf{a}_1 + V_2 \mathbf{a}_2 + V_3 \mathbf{a}_3, \quad (13)$$

we can rewrite the sedenion in following compact form

$$\tilde{V} = V_0 + \vec{V}. \quad (14)$$

Further we will indicate sedenion-scalars and sedenion-vectors with the bold capital letters.

Let us consider the sedenionic multiplication in detail. The sedenionic product of two sedenions $\tilde{\mathbf{A}}$ and $\tilde{\mathbf{B}}$ can be represented in the following form

$$\tilde{\mathbf{A}}\tilde{\mathbf{B}} = (\mathbf{A}_0 + \vec{\mathbf{A}})(\mathbf{B}_0 + \vec{\mathbf{B}}) = \mathbf{A}_0\mathbf{B}_0 + \mathbf{A}_0\vec{\mathbf{B}} + \vec{\mathbf{A}}\mathbf{B}_0 + (\vec{\mathbf{A}} \cdot \vec{\mathbf{B}}) + [\vec{\mathbf{A}} \times \vec{\mathbf{B}}] \quad (15)$$

Here we denoted the sedenionic scalar multiplication of two sedenion-vectors (internal product) by symbol “ \cdot ” and round brackets

$$(\vec{\mathbf{A}} \cdot \vec{\mathbf{B}}) = -\mathbf{A}_1\mathbf{B}_1 - \mathbf{A}_2\mathbf{B}_2 - \mathbf{A}_3\mathbf{B}_3, \quad (16)$$

and sedenionic vector multiplication (external product) by symbol “ \times ” and square brackets,

$$[\vec{\mathbf{A}} \times \vec{\mathbf{B}}] = (\mathbf{A}_2\mathbf{B}_3 - \mathbf{A}_3\mathbf{B}_2)\mathbf{a}_1 + (\mathbf{A}_3\mathbf{B}_1 - \mathbf{A}_1\mathbf{B}_3)\mathbf{a}_2 + (\mathbf{A}_1\mathbf{B}_2 - \mathbf{A}_2\mathbf{B}_1)\mathbf{a}_3. \quad (17)$$

In (16) and (17) the multiplication of sedenionic components is performed in accordance with (11) and Table 2. Thus the sedenionic product

$$\tilde{\mathbf{F}} = \tilde{\mathbf{A}}\tilde{\mathbf{B}} = \mathbf{F}_0 + \vec{\mathbf{F}} \quad (18)$$

has the following components:

$$\begin{aligned} \mathbf{F}_0 &= \mathbf{A}_0\mathbf{B}_0 - \mathbf{A}_1\mathbf{B}_1 - \mathbf{A}_2\mathbf{B}_2 - \mathbf{A}_3\mathbf{B}_3, \\ \mathbf{F}_1 &= \mathbf{A}_1\mathbf{B}_0 + \mathbf{A}_0\mathbf{B}_1 + (\mathbf{A}_2\mathbf{B}_3 - \mathbf{A}_3\mathbf{B}_2), \\ \mathbf{F}_2 &= \mathbf{A}_2\mathbf{B}_0 + \mathbf{A}_0\mathbf{B}_2 + (\mathbf{A}_3\mathbf{B}_1 - \mathbf{A}_1\mathbf{B}_3), \\ \mathbf{F}_3 &= \mathbf{A}_3\mathbf{B}_0 + \mathbf{A}_0\mathbf{B}_3 + (\mathbf{A}_1\mathbf{B}_2 - \mathbf{A}_2\mathbf{B}_1). \end{aligned} \quad (19)$$

Note that in the sedenionic algebra the square of vector is defined as

$$\vec{\mathbf{A}}^2 = (\vec{\mathbf{A}} \cdot \vec{\mathbf{A}}) = -\mathbf{A}_1^2 - \mathbf{A}_2^2 - \mathbf{A}_3^2, \quad (20)$$

and the square of modulus of vector is

$$|\vec{\mathbf{A}}|^2 = -(\vec{\mathbf{A}} \cdot \vec{\mathbf{A}}) = \mathbf{A}_1^2 + \mathbf{A}_2^2 + \mathbf{A}_3^2. \quad (21)$$

3. Spatial rotation and space-time inversion

The rotation of sedenion $\tilde{\mathbf{V}}$ on the angle θ around the absolute unit vector \vec{n} is realized by uncompleted sedenion

$$\tilde{\mathbf{U}} = \cos(\theta/2) + \vec{n} \sin(\theta/2) \quad (22)$$

and by conjugated sedenion $\tilde{\mathbf{U}}^*$:

$$\tilde{\mathbf{U}}^* = \cos(\theta/2) - \vec{n} \sin(\theta/2) \quad (23)$$

with

$$\tilde{\mathbf{U}}\tilde{\mathbf{U}}^* = \tilde{\mathbf{U}}^*\tilde{\mathbf{U}} = 1. \quad (24)$$

The transformed sedenion $\tilde{\mathbf{V}}'$ is defined as sedenionic product

$$\tilde{\mathbf{V}}' = \tilde{\mathbf{U}}^* \tilde{\mathbf{V}} \tilde{\mathbf{U}}, \quad (25)$$

Thus the transformed sedenion $\tilde{\mathbf{V}}'$ can be written as

$$\begin{aligned}\vec{V}' &= [\cos(\theta/2) - \vec{n} \sin(\theta/2)](\vec{V}_0 + \vec{V})[\cos(\theta/2) + \vec{n} \sin(\theta/2)] \\ &= \vec{V}_0 + \vec{V} \cos \theta - \vec{n}(\vec{n} \cdot \vec{V})(1 - \cos \theta) - [\vec{n} \times \vec{V}] \sin \theta.\end{aligned}\quad (26)$$

It is clearly seen that rotation does not transform the sedenion-scalar part, but the sedenionic vector \vec{V} is rotated on the angle θ around \vec{n} .

The operations of time inversion (\hat{R}_t), space inversion (\hat{R}_r) and space-time inversion (\hat{R}_{tr}) are connected with transformations in $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ basis and can be presented as

$$\begin{aligned}\hat{R}_t \vec{V} &= -\mathbf{e}_2 \vec{V} \mathbf{e}_2 = \vec{V}_0 - \mathbf{e}_1 \vec{V}_1 + \mathbf{e}_2 \vec{V}_2 - \mathbf{e}_3 \vec{V}_3, \\ \hat{R}_r \vec{V} &= -\mathbf{e}_1 \vec{V} \mathbf{e}_1 = \vec{V}_0 + \mathbf{e}_1 \vec{V}_1 - \mathbf{e}_2 \vec{V}_2 - \mathbf{e}_3 \vec{V}_3, \\ \hat{R}_{tr} \vec{V} &= -\mathbf{e}_3 \vec{V} \mathbf{e}_3 = \vec{V}_0 - \mathbf{e}_1 \vec{V}_1 - \mathbf{e}_2 \vec{V}_2 + \mathbf{e}_3 \vec{V}_3.\end{aligned}\quad (27)$$

4. Sedenionic Lorentz transformations

The relativistic event four-vector can be represented in the follow sedenionic form:

$$\vec{S} = \mathbf{e}_1 ct + \mathbf{e}_2 \vec{r}. \quad (28)$$

The square of this value is the Lorentz invariant

$$\vec{S} \vec{S} = -c^2 t^2 + x^2 + y^2 + z^2. \quad (29)$$

The Lorentz transformation of event four-vector is realized by uncompleted sedenions

$$\begin{aligned}\tilde{L} &= \text{ch } \vartheta + \mathbf{e}_3 \vec{m} \text{sh } \vartheta, \\ \tilde{L}^* &= \text{ch } \vartheta - \mathbf{e}_3 \vec{m} \text{sh } \vartheta,\end{aligned}\quad (30)$$

where $\text{th } 2\vartheta = v/c$, v is velocity of motion along the absolute unit vector \vec{m} . Note that

$$\tilde{L}^* \tilde{L} = \tilde{L} \tilde{L}^* = 1. \quad (31)$$

The transformed event four-vector \vec{S}' is written as

$$\begin{aligned}\vec{S}' &= \tilde{L}^* \vec{S} \tilde{L} = (\text{ch } \vartheta - \mathbf{e}_3 \text{sh } \vartheta \vec{m})(\mathbf{e}_1 ct + \mathbf{e}_2 \vec{r})(\text{ch } \vartheta + \mathbf{e}_3 \text{sh } \vartheta \vec{m}) = \\ &= \mathbf{e}_1 ct \text{ch } 2\vartheta + \mathbf{e}_1 (\vec{m} \cdot \vec{r}) \text{sh } 2\vartheta \\ &+ \mathbf{e}_2 \vec{r} - \mathbf{e}_2 ct \vec{m} \text{sh } 2\vartheta - 2\mathbf{e}_2 (\vec{m} \cdot \vec{r}) \vec{m} \text{sh}^2 \vartheta + \mathbf{e}_2 (\vec{m} \cdot \vec{r}) \vec{m} (1 - \text{ch } 2\vartheta).\end{aligned}\quad (32)$$

Separating the values with \mathbf{e}_1 and \mathbf{e}_2 we get the well known formulas for time and coordinates transformation [24]:

$$t' = \frac{t - x v / c^2}{\sqrt{1 - v^2 / c^2}}, \quad x' = \frac{x - t v}{\sqrt{1 - v^2 / c^2}}, \quad y' = y, \quad z' = z, \quad (33)$$

where x is the coordinate along the \vec{m} vector.

Let us also consider the Lorentz transformation of the full sedenion \vec{V} .

The transformed sedenion \vec{V}' can be written as sedenionic product

$$\vec{V}' = \tilde{L}^* \vec{V} \tilde{L}. \quad (34)$$

$$\begin{aligned}
\tilde{V}' &= (\text{ch } \vartheta - \mathbf{e}_{\text{tr}} \text{sh } \vartheta \bar{m}) (V_0 + \vec{V}) (\text{ch } \vartheta + \mathbf{e}_{\text{tr}} \text{sh } \vartheta \bar{m}) \\
&= V_0 \text{ch}^2 \vartheta + \mathbf{e}_{\text{tr}} V_0 \mathbf{e}_{\text{tr}} \text{sh}^2 \vartheta - (\mathbf{e}_{\text{tr}} V_0 - V_0 \mathbf{e}_{\text{tr}}) \bar{m} \text{ch } \vartheta \text{sh } \vartheta \\
&\quad + \vec{V} \text{ch}^2 \vartheta - \mathbf{e}_{\text{tr}} \bar{m} \vec{V} \bar{m} \mathbf{e}_{\text{tr}} \text{sh}^2 \vartheta - (\mathbf{e}_{\text{tr}} \bar{m} \vec{V} - \vec{V} \bar{m} \mathbf{e}_{\text{tr}}) \text{ch } \vartheta \text{sh } \vartheta.
\end{aligned} \tag{35}$$

Rewriting the expression (35) with scalar (16) and vector (17) products we get

$$\begin{aligned}
\tilde{V}' &= V_0 \text{ch}^2 \vartheta + \mathbf{e}_{\text{tr}} V_0 \mathbf{e}_{\text{tr}} \text{sh}^2 \vartheta - (\mathbf{e}_{\text{tr}} V_0 - V_0 \mathbf{e}_{\text{tr}}) \bar{m} \text{ch } \vartheta \text{sh } \vartheta \\
&\quad + \vec{V} \text{ch}^2 \vartheta - \mathbf{e}_{\text{tr}} \vec{V} \mathbf{e}_{\text{tr}} \text{sh}^2 \vartheta - 2 \mathbf{e}_{\text{tr}} (\bar{m} \cdot \vec{V}) \mathbf{e}_{\text{tr}} \bar{m} \text{sh}^2 \vartheta \\
&\quad - (\mathbf{e}_{\text{tr}} (\bar{m} \cdot \vec{V}) - (\vec{V} \cdot \bar{m}) \mathbf{e}_{\text{tr}}) \text{ch } \vartheta \text{sh } \vartheta - (\mathbf{e}_{\text{tr}} [\bar{m} \times \vec{V}] - [\vec{V} \times \bar{m}] \mathbf{e}_{\text{tr}}) \text{ch } \vartheta \text{sh } \vartheta.
\end{aligned} \tag{36}$$

Thus, the transformed sedenion has the following components:

$$\begin{aligned}
V' &= V, \\
V'_{\text{tr}} &= V_{\text{tr}}, \\
V'_r &= V_r \text{ch } 2\vartheta - \mathbf{e}_{\text{tr}} (\bar{m} \cdot \vec{V}_t) \text{sh } 2\vartheta, \\
V'_t &= V_t \text{ch } 2\vartheta - \mathbf{e}_{\text{tr}} (\bar{m} \cdot \vec{V}_r) \text{sh } 2\vartheta, \\
\vec{V}' &= \vec{V} \text{ch } 2\vartheta + 2 (\bar{m} \cdot \vec{V}) \bar{m} \text{sh}^2 \vartheta - \mathbf{e}_{\text{tr}} [\bar{m} \times \vec{V}_{\text{tr}}] \text{sh } 2\vartheta, \\
\vec{V}'_{\text{tr}} &= \vec{V}_{\text{tr}} \text{ch } 2\vartheta + 2 (\bar{m} \cdot \vec{V}_{\text{tr}}) \bar{m} \text{sh}^2 \vartheta - \mathbf{e}_{\text{tr}} [\bar{m} \times \vec{V}] \text{sh } 2\vartheta, \\
\vec{V}'_r &= \vec{V}_r - 2 (\bar{m} \cdot \vec{V}_r) \bar{m} \text{sh}^2 \vartheta - \mathbf{e}_{\text{tr}} V_t \bar{m} \text{sh } 2\vartheta, \\
\vec{V}'_t &= \vec{V}_t - 2 (\bar{m} \cdot \vec{V}_t) \bar{m} \text{sh}^2 \vartheta - \mathbf{e}_{\text{tr}} V_r \bar{m} \text{sh } 2\vartheta.
\end{aligned} \tag{37}$$

4. Subalgebras of space-time quaternions and octonions

The sedenionic basis introduced above enables constructing different types of low-dimensional hypercomplex numbers. For example one can introduce space-time complex numbers

$$\begin{aligned}
Z_t &= z_1 + \mathbf{e}_t z_2, \\
Z_r &= z_1 + \mathbf{e}_r z_2, \\
Z_{\text{tr}} &= z_1 + \mathbf{e}_{\text{tr}} z_2,
\end{aligned} \tag{38}$$

which are transformed under space and time conjugation. Moreover we can consider the space-time quaternions, which differ in their properties with respect to the operations of the spatial and time inversion.

$$\hat{q} = q_0 \mathbf{a}_0 + \mathbf{e}_0 (q_1 \mathbf{a}_1 + q_2 \mathbf{a}_2 + q_3 \mathbf{a}_3), \tag{39}$$

$$\hat{q}_t = q_0 \mathbf{a}_0 + \mathbf{e}_t (q_1 \mathbf{a}_1 + q_2 \mathbf{a}_2 + q_3 \mathbf{a}_3), \tag{40}$$

$$\hat{q}_r = q_0 \mathbf{a}_0 + \mathbf{e}_r (q_1 \mathbf{a}_1 + q_2 \mathbf{a}_2 + q_3 \mathbf{a}_3), \tag{41}$$

$$\hat{q}_{\text{tr}} = q_0 \mathbf{a}_0 + \mathbf{e}_{\text{tr}} (q_1 \mathbf{a}_1 + q_2 \mathbf{a}_2 + q_3 \mathbf{a}_3), \tag{42}$$

The absolute quaternion (39) is the sum of the absolute scalar and absolute vector. It remains constant under the transformations of space and time inversion (27). Time quaternion \hat{q}_t , space

quaternion \widehat{q}_r and space-time quaternion \widehat{q}_{tr} are transformed under inversions in accordance with the commutation rules for the basis elements $\mathbf{e}_t, \mathbf{e}_r, \mathbf{e}_{tr}$. For example, performing the operation of time inversion (see (27)) with the quaternion \widehat{q}_t we obtain the conjugated quaternion

$$\overline{\widehat{q}}_t = -\mathbf{e}_r \widehat{q}_t \mathbf{e}_r = q_0 \mathbf{a}_0 - \mathbf{e}_t (q_1 \mathbf{a}_1 + q_2 \mathbf{a}_2 + q_3 \mathbf{a}_3). \quad (43)$$

On the other hand, the sedenionic basis allows one to construct various types of space-time eight-component octonions:

$$\check{G}_t = G_{00} + G_{01} \mathbf{a}_1 + G_{02} \mathbf{a}_2 + G_{03} \mathbf{a}_3 + \mathbf{e}_t G_{10} + \mathbf{e}_t (G_{11} \mathbf{a}_1 + G_{12} \mathbf{a}_2 + G_{13} \mathbf{a}_3), \quad (44)$$

$$\check{G}_r = G_{00} + G_{01} \mathbf{a}_1 + G_{02} \mathbf{a}_2 + G_{03} \mathbf{a}_3 + \mathbf{e}_r G_{20} + \mathbf{e}_r (G_{21} \mathbf{a}_1 + G_{22} \mathbf{a}_2 + G_{23} \mathbf{a}_3), \quad (45)$$

$$\check{G}_{tr} = G_{00} + G_{01} \mathbf{a}_1 + G_{02} \mathbf{a}_2 + G_{03} \mathbf{a}_3 + \mathbf{e}_{tr} G_{30} + \mathbf{e}_{tr} (G_{31} \mathbf{a}_1 + G_{32} \mathbf{a}_2 + G_{33} \mathbf{a}_3). \quad (46)$$

5. Generalized sedenionic equations of relativistic quantum mechanics

The wave function of free quantum particle should satisfy an equation, which is obtained from the Einstein relation for energy and momentum

$$E^2 - c^2 p^2 = m_0^2 c^4 \quad (47)$$

by means of changing classical energy E and momentum \vec{p} on corresponding quantum-mechanical operators:

$$\hat{E} = i\hbar \frac{\partial}{\partial t} \text{ and } \hat{p} = -i\hbar \vec{\nabla}. \quad (48)$$

Here c is the velocity of light, m_0 is the particle rest mass, \hbar is the Planck constant. The Einstein relation (47) can be written using space-time sedenion algebra in the following form:

$$(\mathbf{e}_t E + \mathbf{e}_r c \vec{p} + i \mathbf{e}_{tr} m_0 c^2)(\mathbf{e}_t E + \mathbf{e}_r c \vec{p} + i \mathbf{e}_{tr} m_0 c^2) = 0. \quad (49)$$

Let us consider the wave function in the form of space-time sedenion

$$\tilde{W}(t, \vec{r}) = W_0(t, \vec{r}) + \tilde{W}(t, \vec{r}). \quad (50)$$

Then the generalized sedenionic wave equation for free particle can be written in the symmetric form

$$\left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \vec{\nabla} + \mathbf{e}_{tr} \frac{m_0 c}{\hbar} \right) \left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \vec{\nabla} + \mathbf{e}_{tr} \frac{m_0 c}{\hbar} \right) \tilde{W} = 0. \quad (51)$$

Note that for electrically charged particle in an external electromagnetic field we have the following sedenionic wave equation:

$$\left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} + \mathbf{e}_t \frac{ie}{\hbar c} \varphi - \mathbf{e}_r \vec{\nabla} + \mathbf{e}_r \frac{ie}{\hbar c} \vec{A} + \mathbf{e}_{tr} \frac{m_0 c}{\hbar} \right) \left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} + \mathbf{e}_t \frac{ie}{\hbar c} \varphi - \mathbf{e}_r \vec{\nabla} + \mathbf{e}_r \frac{ie}{\hbar c} \vec{A} + \mathbf{e}_{tr} \frac{m_0 c}{\hbar} \right) \tilde{W} = 0. \quad (52)$$

This equation describes the particles with spin 1/2 in an external electromagnetic field [18, 21].

There is a special class of particles described by the first-order wave equation [25]. For these particles the sedenionic Dirac-like wave equation has the following form:

$$\left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \vec{\nabla} + \mathbf{e}_{tr} \frac{m_0 c}{\hbar} \right) \tilde{W} = 0. \quad (53)$$

In fact, this equation describes the special quantum field with zero field intensities [19]. Analogously the electrically charged particle interacting with external electromagnetic field is described by the following sedenionic first-order wave equation:

$$\left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} + \mathbf{e}_t \frac{ie}{\hbar c} \varphi - \mathbf{e}_r \vec{\nabla} + \mathbf{e}_r \frac{ie}{\hbar c} \vec{A} + \mathbf{e}_{tr} \frac{m_0 c}{\hbar} \right) \tilde{\mathcal{W}} = 0. \quad (54)$$

This equation also describes the particles with spin 1/2 in an external electromagnetic field [19].

5. Generalized sedenionic equations for massive force field

The generalized sedenionic wave equation

$$\left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \vec{\nabla} + \mathbf{e}_{tr} \frac{m_0 c}{\hbar} \right) \left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \vec{\nabla} + \mathbf{e}_{tr} \frac{m_0 c}{\hbar} \right) \tilde{\mathcal{W}} = 0 \quad (55)$$

enables another interpretation. It can be considered as the equation for the force massive field [26]. In this case the parameter m_0 is the mass of quantum of field. Then considering the phenomenological source of field we can propose the following nonhomogeneous wave equation for the field potential:

$$\left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \vec{\nabla} + \mathbf{e}_{tr} \frac{m_0 c}{\hbar} \right) \left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \vec{\nabla} + \mathbf{e}_{tr} \frac{m_0 c}{\hbar} \right) \tilde{\mathcal{W}} = \tilde{\mathcal{J}}. \quad (56)$$

In the special case when the mass is equal to zero the equation (56) coincides with the equation for electromagnetic field in a vacuum. Indeed, choosing the potential as

$$\tilde{\mathcal{W}} = \mathbf{e}_t \varphi + \mathbf{e}_r \vec{A}, \quad (57)$$

and the source of electromagnetic field as

$$\tilde{\mathcal{J}} = -\mathbf{e}_t 4\pi\rho - \mathbf{e}_r \frac{4\pi}{c} \vec{j}, \quad (58)$$

we obtain the following wave equation:

$$\left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \vec{\nabla} \right) \left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \vec{\nabla} \right) (\mathbf{e}_t \varphi + \mathbf{e}_r \vec{A}) = -\mathbf{e}_t 4\pi\rho - \mathbf{e}_r \frac{4\pi}{c} \vec{j}. \quad (59)$$

After the action of the first operator in the left-hand side of equation (59) we obtain

$$\left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \vec{\nabla} \right) (\mathbf{e}_t \varphi + \mathbf{e}_r \vec{A}) = -\frac{1}{c} \frac{\partial \varphi}{\partial t} + \mathbf{e}_{tr} \frac{1}{c} \frac{\partial \vec{A}}{\partial t} + \mathbf{e}_{tr} \vec{\nabla} \varphi + (\vec{\nabla} \cdot \vec{A}) + [\vec{\nabla} \times \vec{A}]. \quad (60)$$

Using the sedenionic definitions of the electric and magnetic fields

$$\begin{aligned} \vec{E} &= -\frac{1}{c} \frac{\partial \vec{A}}{\partial t} - \vec{\nabla} \varphi, \\ \vec{H} &= [\vec{\nabla} \times \vec{A}], \end{aligned} \quad (61)$$

and taking into account the Lorentz gauge condition

$$\frac{1}{c} \frac{\partial \varphi}{\partial t} - (\vec{\nabla} \cdot \vec{A}) = 0, \quad (62)$$

We can rewrite the expression (60) as

$$\left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \bar{\nabla} \right) (\mathbf{e}_t \varphi + \mathbf{e}_r \bar{A}) = -\mathbf{e}_r \bar{E} + \bar{H}. \quad (63)$$

Then the wave equation (59) can be represented in the following form:

$$\left(\mathbf{e}_t \frac{1}{c} \frac{\partial}{\partial t} - \mathbf{e}_r \bar{\nabla} \right) (-\mathbf{e}_r \bar{E} + \bar{H}) = -\mathbf{e}_t 4\pi\rho - \mathbf{e}_r \frac{4\pi}{c} \bar{j}. \quad (64)$$

Performing sedenionic multiplication in the left-hand side of equation (64) we get

$$\begin{aligned} & \mathbf{e}_r \frac{1}{c} \frac{\partial \bar{E}}{\partial t} + \mathbf{e}_t (\bar{\nabla} \cdot \bar{E}) + \mathbf{e}_t [\bar{\nabla} \times \bar{E}] \\ & + \mathbf{e}_t \frac{1}{c} \frac{\partial \bar{H}}{\partial t} - \mathbf{e}_r (\bar{\nabla} \cdot \bar{H}) - \mathbf{e}_r [\bar{\nabla} \times \bar{H}] = -\mathbf{e}_t 4\pi\rho - \mathbf{e}_r \frac{4\pi}{c} \bar{j}. \end{aligned} \quad (65)$$

Separating space-time values we obtain the system of Maxwell equations in the following form:

$$\begin{aligned} \mathbf{e}_t (\bar{\nabla} \cdot \bar{E}) &= -\mathbf{e}_t 4\pi\rho, & (\text{time scalar part}) \\ \mathbf{e}_r [\bar{\nabla} \times \bar{H}] &= \mathbf{e}_r \frac{1}{c} \frac{\partial \bar{E}}{\partial t} + \mathbf{e}_r \frac{4\pi}{c} \bar{j}, & (\text{space vector part}) \\ \mathbf{e}_t [\bar{\nabla} \times \bar{E}] &= -\mathbf{e}_t \frac{1}{c} \frac{\partial \bar{H}}{\partial t}, & (\text{time vector part}) \\ \mathbf{e}_r (\bar{\nabla} \cdot \bar{H}) &= 0. & (\text{space scalar part}) \end{aligned} \quad (66)$$

The system (66) coincides with the Maxwell equations.

6. Discussion

The algebra of sedenions proposed in this article is the anticommutative associative space-time Clifford algebra. The sedenionic basis elements \mathbf{a}_n are responsible for the spatial rotation, while the elements \mathbf{e}_n are responsible for the space-time inversions. Mathematically, these two bases are equivalent, and the different physical properties attributed to them are an important physical essence of our sedenionic hypothesis.

In contrast to the previously discussed sedeonic algebra [20-23], which uses the multiplication rules of basic elements \mathbf{a}'_n and \mathbf{e}'_n proposed by A.Macfarlane [27], the multiplication rules for sedenionic basis elements \mathbf{a}_n and \mathbf{e}_n coincide with the rules for quaternion units introduced by W.R.Hamilton [28]. There is a close connection between these two bases. The transition from the sedeonic basis to sedenionic basis is performed by following replacement:

$$\begin{aligned} \mathbf{a}'_n &= i\mathbf{a}_n, \\ \mathbf{e}'_n &= i\mathbf{e}_n. \end{aligned}$$

There is one disadvantage of sedenions connected with the fact that the square of the vector is a negative value. However, on the other side the sedenionic rules of cross-multiplying do not contain the imaginary unit and this leads to the considerable simplifications in the calculations. But of course, the physical results do not depend on the choice of algebra, so these two algebras are equivalent.

7. Conclusion

Thus in this paper we presented the sixteen-component hypercomplex values sedenions, generating associative noncommutative space-time algebra. We proposed the generalization of the relativistic quantum mechanics equations and equations for the massive field based on sedenionic space-time operators and hypercomplex scalar-vector wave functions.

Acknowledgments

The authors are very thankful to G.V. Mironova for kind assistance and moral support.

References

1. S.L. Adler, "Quaternionic Quantum Mechanics and Quantum Fields", New York: Oxford University Press, 1995.
2. K. Imaeda – A new formulation of classical electrodynamics, *Nuovo cimento*, **32**(1), 138-162, (1976).
3. V. Majernik – Quaternionic formulation of the classical fields, *Advances in Applied Clifford Algebras*, **9**(1), 119-130, (1999).
4. A.J. Davies – Quaternionic Dirac equation, *Physical Review D*, **41**(8), 2628-2630 (1990).
5. C. Schwartz – Relativistic quaternionic wave equation, *Journal of Mathematical Physics*, **47**, 122301 (2006).
6. L.Yu-Fen – Triality, biquaternion and vector representation of the Dirac equation, *Advances in Applied Clifford Algebras*, **12**(2), 109-124 (2002).
7. M. Gogberashvili – Octonionic electrodynamics – *Journal of Physics A: Mathematics in General*, **39**, 7099-7104, (2006).
8. A. Gamba – Maxwell's equations in octonion form, *Nuovo Cimento A*, **111**(3), 293-302, (1998).
9. T. Tolan, K. Özdas, M. Tanisli – Reformulation of electromagnetism with octonions, *Nuovo Cimento B*, **121**(1), 43-55, (2006).
10. M. Gogberashvili - Octonionic version of Dirac equations, *International Journal of Modern Physics A*, **21**(17), 3513-3523 (2006).
11. S. De Leo, K. Abdel-Khalek – Octonionic Dirac equation, *Progress in Theoretical Physics*, **96**, 833-846 (1996).
12. L.E. Dickson - On quaternions and their generalization and the history of the eight square theorem", *Annals of Mathematics (Second Series)*, **20** (3), 155–171 (1919).
13. K. Imaeda, M. Imaeda – Sedenions: algebra and analysis, *Applied Mathematics and Computations*, **115**, 77-88 (2000).
14. K. Carmody – Circular and hyperbolic quaternions, octonions, and sedenions, *Applied Mathematics and Computations*, **28**, 47-72 (1988).
15. K. Carmody – Circular and hyperbolic quaternions, octonions, and sedenions – further results, *Applied Mathematics and Computations*, **84**, 27-47 (1997).
16. J. Köpflinger – Dirac equation on hyperbolic octonions, *Applied Mathematics and Computations*, **182**, 443-446 (2006).
17. V.L. Mironov, S.V. Mironov – Octonic representation of electromagnetic field equations, *Journal of Mathematical Physics*, **50**, 012901 (2009).
18. V.L. Mironov, S.V. Mironov – Octonic second-order equations of relativistic quantum mechanics, *Journal of Mathematical Physics*, **50**, 012302 (2009).
19. V.L. Mironov, S.V. Mironov – Octonic first-order equations of relativistic quantum mechanics, *International Journal of Modern Physics A*, **24**(22), 4157-4167 (2009).
20. V.L. Mironov, S.V. Mironov – Noncommutative sedeons and their application in field theory, *ArXiv* <http://arxiv.org/abs/1111.4035>, (2011).
21. V.L. Mironov, S.V. Mironov – Reformulation of relativistic quantum mechanics equations with non-commutative sedeons, *Applied Mathematics*, **4**(10C), 53-60 (2013).

22. V.L. Mironov, S.V. Mironov, S. A. Korolev – Sedeonic theory of massless fields, ArXiv <http://arxiv.org/abs/1206.5969>, (2012).
23. S.V. Mironov, V.L. Mironov – Sedeonic theory of massive fields, ViXra <http://vixra.org/abs/1311.0005>, (2013).
24. L.D. Landau and E.M. Lifshits, “Classical Theory of Fields” (4th ed.). New York: Pergamon Press, Vol. 2 of the Course of Theoretical Physics. (1975).
25. P.A.M. Dirac, “The Principles of Quantum Mechanics”, Clarendon Press, Oxford, (1958).
26. C. Cafaro, S.A. Ali – The spacetime algebra approach to massive classical electrodynamics with magnetic monopoles, *Advances in Applied Clifford Algebras*, **17**, 23-36 (2006).
27. A. Macfarlane - Hyperbolic quaternions, *Proceedings of the Royal Society at Edinburgh*, 1899-1900 session, pp. 169–181 (1900).
28. W.R. Hamilton, "Lectures on Quaternions", Royal Irish Academy, (1853).