

SOLUTION OF TWO QUESTIONS CONCERNING THE DIVISOR FUNCTION AND THE PSEUDO – SMARANDACHE FUNCTION

Zhong Li

Abstract In this paper we completely solve two questions concerning the divisor function and the pseudo – Smarandache function.

Key words divisor function, pseudo – Smarandache function, functional equation

1 Introduction

Let \mathbb{N} be the set of all positive integers . For any $n \in \mathbb{N}$, let

$$(1) \quad d(n) = \sum_{d|n} 1,$$

$$(2) \quad Z(n) = \min \{ a \mid a \in \mathbb{N}, n \mid \sum_{j=1}^a j \}$$

Then $d(n)$ and $Z(n)$ are called the divisor function and the pseudo – Smarandache function of n , respectively, In^[1], Ashbacher posed the following unsolved questions.

Question 1 How many solutions n are there to the functional equation.

$$(3) \quad Z(n) = d(n), n \in \mathbb{N}?$$

Question 2 How many solutions n are there to the functional equation.

$$(4) \quad Z(n) + d(n) = n, n \in \mathbb{N}?$$

In this paper we completely solve the above questions as follows.

Theorem 1 The equation (3) has only the solutions $n = 1, 3$ and 10 .

Theorem 2 The equation (4) has only the solution $n = 56$.

2 Proof of Theorem 1

A computer search showed that (3) has only the solutions $n = 1, 3$ and 10 with $n \leq 10000$ (see [1]).

We now let n be a solution of (3) with $n \neq 1, 3$ or 10 . Then we have $n > 10000$. Let

$$(5) \quad n = p_1^{r_1} p_2^{r_2} \cdots p_k^{r_k}$$

be the factorization of n . By [2, Theorem 273], we get from (1) and (5) that

$$(6) \quad d(n) = (r_1 + 1)(r_2 + 1) \cdots (r_k + 1).$$

On the other hand, since $\sum_{j=1}^a j = a(a+1)/2$ for any $a \in \mathbb{N}$, we see from (2) that $n \mid Z(n)(Z(n)+1)/2$. It implies that $Z(n)(Z(n)+1)/2 \geq n$. So we have

$$(7) \quad Z(n) \geq \sqrt{2n + \frac{1}{4}} - \frac{1}{2}$$

Hence, by (3), (5), (6) and (7), we get

$$(8) \quad 1 > \sqrt{2} \prod_{i=1}^k \frac{p_i^{r_i/2}}{r_i + 1} - \frac{1}{2} \prod_{i=1}^k \frac{1}{r_i + 1}$$

If $p_1 > 3$, then from (8) we get $p_1 \geq 5$ and

$$1 \geq \sqrt{2} \left(\frac{\sqrt{5}}{2}\right)^k - \frac{1}{2^{k+1}} > 1,$$

a contradiction. Therefore, if (8) holds, then either $p_1 = 2$ or $p_1 = 3$. By the same method, then n must satisfy one of the following conditions.

$$(i) p_1 = 2 \text{ and } r_1 \leq 4 .$$

$$(ii) p_1 = 3 \text{ and } r_1 = 1 .$$

However, by (8), we can calculate that $n < 10000$, a contradiction. Thus, the theorem is proved.

3 Proof of Theorem 2

A computer search showed that (4) has only the solution $n = 56$ with $n \leq 10000$ (see [1]). We now let n be a solution of (4) with $n \neq 56$. Then we have $n > 10000$. We see from (4) that

$$(9) \quad Z(n) \equiv -d(n) \pmod{n}$$

It implies that.

$$(10) \quad Z(n) + 1 \equiv 1 - d(n) \pmod{n}$$

By the proof of Theorem 1, we have $n \mid Z(n)(Z(n) + 1)/2$, by (2). It can be written as

$$(11) \quad Z(n)(Z(n) + 1) \equiv 0 \pmod{n}.$$

Substituting (9) and (10) into (11), we get

$$(12) \quad d(n)(d(n) - 1) \equiv 0 \pmod{n}.$$

Notice that $d(n) > 1$ if $n > 1$. We see from (12) that

$$(13) \quad (d(n))^2 > n$$

Let (5) be the factorization of n . By (5), (6) and (13), we obtain

$$(14) \quad 1 > \prod_{i=1}^k \frac{p_i^{r_i}}{(r_i + 1)^2}$$

On the other hand, it is a well known fact that $Z(p^r) \geq p^r - 1 > (r + 1)^2$ for any prime power p^r with $p^r > 32$. We find from (14) that $k \geq 2$.

If $p_1 > 3$, then $p_i^{r_i} / (r_i + 1)^2 \geq 5/4 > 1$ for $i = 1, 2, \dots, k$, It implies that if (14) holds, then either $p_1 = 2$ or $p_1 = 3$. By the same method, then n must satisfy one of the following conditions:

(i) $p_1 = 2, p_2 = 3$ and $(r_1, r_2) = (1, 1), (2, 1), (3, 1), (4, 1), (5, 1), (6, 1), (1, 2), (2, 2), (3, 2), (4, 2)$ or $(5, 2)$.

(ii) $p_1 = 2, p_2 > 3$ and $r_1 \leq 5$.

(iii) $p_1 = 3$ and $r_1 = 1$.

However, by (14), we can calculate that $n < 10000$, a contradiction. Thus, the theorem is proved.

References

- [1] C. Ashbacher, The pseudo – Smarandache function and the classical functions of number theory, *Smarandache Notions J.*, 9(1998), 78 – 81.
 [2] G. H. Hardy and E. M. Wright, *An Introduction to the Theory of Numbers*, Oxford, Oxford Univ. Press, 1937.

Department of Mathematics
 Maoming Educational College
 Maoming, Guangdong
 P. R. China