

A Derivation of GravitoElectroMagnetic (GEM) Proca-type Equations in Fractional Space

Victor Christianto¹, email: victorchristianto@gmail.com

Abstract

In a recent paper, M. Zubair et al. described a novel approach for fractional space generalization of the differential electromagnetic equations. A new form of vector differential operator Del, and its related differential operators, is formulated in fractional space. Using these modified vector differential operators, the classical Maxwell equations have been worked out for fractal media. In the meantime, there are other papers discussing fractional Maxwell equations. However, so far there is no derivation of Proca equations and Proca-type equations for GEM in fractional space. Therefore in this paper I present for the first time a derivation of GravitoElectroMagnetic (GEM) Proca-type equations in fractional space. Considering that Proca equations may be used to explain some electromagnetic effect in superconductor, then fractional GEM Proca-type equations may be expected to explain some gravitomagnetic effects of superconductor for fractal media. It is our hope, that this paper may stimulate further investigation and experiments in particular with respect to gravitomagnetic effects.

Introduction

There has been much interests to study different physical phenomenon in fractional dimensional space during the last few decades. It is also important to mention that the experimental measurement of the dimension of real world is 3 ± 10^{-6} , not exactly 3.[1]

In a recent paper, M. Zubair et al. described a novel approach for fractional space generalization of the differential electromagnetic equations.[1] A new form of vector differential operator Del, and its related differential operators, is formulated in fractional space. Using these modified vector differential operators, the classical Maxwell equations have been worked out for fractal media. In the meantime, there are other papers discussing fractional Maxwell equations.[2][3] However, so far there is no derivation of Proca equations and Proca-type equations for GEM in fractional space. Therefore in this paper I present for the first time a derivation of GravitoElectroMagnetic (GEM) Proca-type

¹ URL: <http://www.sciprint.org>, <http://independent.academia.edu/VChristianto>

equations in fractional space. Considering that Proca equations may be used to explain some electromagnetic effect in superconductor[4], then fractional GEM Proca-type equations may be expected to explain some gravitomagnetic effects of superconductor for fractal media.[5] It is our hope, that this paper may stimulate further investigation and experiments in particular with respect to gravitomagnetic effects.

Review of previous result: Maxwell equations in fractional space

I will not re-derive Maxwell equations here. For a good reference on *Classical Electrodynamics*, see for example Julian Schwinger et al.'s book [9]. Penrose also discusses Maxwell equations shortly in his book: *The Road to Reality*. [10]

Zubair et al. were able to write a differential form of Maxwell equations in far-field region in the fractional space as follows [1]:

$$\operatorname{div}_D D = \rho_v, \quad (1)$$

$$\operatorname{div}_D B = 0, \quad (2)$$

$$\operatorname{curl}_D E = -\frac{\partial B}{\partial t}, \quad (3)$$

$$\operatorname{curl}_D H = J + \frac{\partial D}{\partial t}, \quad (4)$$

and the continuity equation in fractional space as:

$$\operatorname{div}_D J = -\frac{\partial \rho_v}{\partial t}, \quad (5)$$

where div_D and curl_D are defined as follows [1]:

$$\operatorname{div}_D F = \nabla_D \cdot F = \frac{\partial F_x}{\partial x} + \frac{1}{2} \frac{(\alpha_1 - 1)F_x}{x} + \frac{\partial F_y}{\partial y} + \frac{1}{2} \frac{(\alpha_2 - 1)F_y}{y} + \frac{\partial F_z}{\partial z} + \frac{1}{2} \frac{(\alpha_3 - 1)F_z}{z}, \quad (6)$$

$$\operatorname{curl}_D F = \nabla_D \times F = \begin{vmatrix} \hat{x} & \hat{y} & \hat{z} \\ \frac{\partial}{\partial x} + \frac{1}{2} \frac{\alpha_1 - 1}{x} & \frac{\partial}{\partial y} + \frac{1}{2} \frac{\alpha_2 - 1}{y} & \frac{\partial}{\partial z} + \frac{1}{2} \frac{\alpha_3 - 1}{z} \\ F_x & F_y & F_z \end{vmatrix}, \quad (7)$$

where parameters ($0 < \alpha_1 \leq 1$, $0 < \alpha_2 \leq 1$ and $0 < \alpha_3 \leq 1$) are used to describe the measure distribution of space where each one is acting independently on a single coordinate and the total dimension of the system is $D = \alpha_1 + \alpha_2 + \alpha_3$. [1]

Proca Equations in Fractional Space

Proca equations can be considered as an extension of Maxwell equations, and they have been derived in various ways, see for instance [4][6][7]. It can be shown that Proca equations can be derived from first principles [6], and also that Proca equations may have link with Klein-Gordon equation [7]. In this paper I will not attempt to re-derive Proca equations, instead I will use Proca equations as described in [6]. Then I will derive the Proca equations in fractional space, in accordance with Zubair et al.'s approach as outlined above. [1]

According to Blackledge [7], Proca equations can be written as follows:

$$\nabla \cdot \vec{E} = \frac{\rho}{\epsilon_0} - \kappa^2 \phi, \quad (8)$$

$$\nabla \cdot \vec{B} = 0, \quad (9)$$

$$\nabla \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}, \quad (10)$$

$$\nabla \times \vec{B} = \mu_0 j + \epsilon_0 \mu_0 \frac{\partial \vec{E}}{\partial t} + \kappa^2 \vec{A}, \quad (11)$$

where:

$$\nabla \phi = -\frac{\partial \vec{A}}{\partial t} - \vec{E}, \quad (12)$$

$$\vec{B} = \nabla \times \vec{A}, \quad (13)$$

$$\kappa = \frac{mc_0}{\hbar}. \quad (14)$$

Therefore, by using the definition in equation (6) and (7), we can arrive at Proca equations in fractional space from (8) through (13), as follows:

$$\operatorname{div}_D \vec{E} = \frac{\rho}{\varepsilon_0} - \kappa^2 \phi, \quad (15)$$

$$\operatorname{div}_D \vec{B} = 0, \quad (16)$$

$$\operatorname{curl}_D \vec{E} = -\frac{\partial \vec{B}}{\partial t}, \quad (17)$$

$$\operatorname{curl}_D \vec{B} = \mu_0 \mathbf{j} + \varepsilon_0 \mu_0 \frac{\partial \vec{E}}{\partial t} + \kappa^2 \vec{A}, \quad (18)$$

where:

$$\nabla_D \phi = -\frac{\partial \vec{A}}{\partial t} - \vec{E}, \quad (19)$$

$$\vec{B} = \operatorname{curl}_D \vec{A}, \quad (20)$$

and Del operator ∇_D can be defined as follows [1]:

$$\nabla = \left(\frac{\partial}{\partial x} + \frac{1}{2} \frac{\alpha_1 - 1}{x} \right) \hat{x} + \left(\frac{\partial}{\partial y} + \frac{1}{2} \frac{\alpha_2 - 1}{y} \right) \hat{y} + \left(\frac{\partial}{\partial z} + \frac{1}{2} \frac{\alpha_3 - 1}{z} \right) \hat{z}. \quad (21)$$

To my best knowledge so far, the above expression of Proca equations in fractional space has not been proposed elsewhere before.

GravitoElectroMagnetic (GEM) Proca-type Equations in Fractional Space

The term GravitoElectroMagnetism (GEM) refers to the formal analogies between Newton's law of gravitation and Coulomb's law of electricity. The theoretical analogy between the electromagnetic and the gravitational field equations has been first suggested by Heaviside in 1893, see [8]. The fields of GEM can be defined in close analogy with the classical electrodynamics. Therefore, if we can consider Proca equations as extension of Maxwell equations, then we can also find Proca-type equations for GEM.

In accordance with Demir [8], the Proca-type equations for GEM can be expressed straightforward from their electromagnetic counterpart as follows (Here I use Demir's notations instead of Blackledge's notations):

$$\nabla \cdot \vec{E}_g = -\rho_e - \kappa_g^2 \phi, \quad (22)$$

$$\nabla \cdot \vec{H}_g = 0, \quad (23)$$

$$\nabla \times \vec{E}_g = -\frac{\partial \vec{H}_g}{\partial t}, \quad (24)$$

$$\nabla \times \vec{H}_g = -J_g^e + \frac{\partial \vec{E}_g}{\partial t} + \kappa_g^2 \vec{A}_g^e, \quad (25)$$

where the fields E_g and H_g can be defined in terms of the potentials just as given in equation (12) and (13), and the term κ_g represents the inverse Compton wavelength of the graviton, [8]

$$\kappa_g = \frac{m_g c}{\hbar}. \quad (26)$$

Now I will present the Proca-type equations for GEM in fractional space using the same method as described in the previous section and equation (6) and (7), which can be written as follows:

$$div_D \vec{E}_g = -\rho_e - \kappa_g^2 \phi, \quad (27)$$

$$div_D \vec{H}_g = 0, \quad (28)$$

$$curl_D \vec{E}_g = -\frac{\partial \vec{H}_g}{\partial t}, \quad (29)$$

$$curl_D \vec{H}_g = -J_g^e + \frac{\partial \vec{E}_g}{\partial t} + \kappa_g^2 \vec{A}_g^e, \quad (30)$$

To the best of my knowledge, the above expression of Proca-type equations for GEM in fractional space has not been proposed elsewhere before.

Fractional Helmholtz equation and solution of classical wave equation in fractional space

It is worth noting here that Zubair also wrote Helmholtz equations in fractional space for E and H field as a consequence of his Maxwell equations in fractional space, as follows [1]:

$$\nabla_D^2 E - \mu \varepsilon \frac{\partial^2 E}{\partial t^2} = 0, \quad (31)$$

$$\nabla_D^2 H - \mu\epsilon \frac{\partial^2 H}{\partial t^2} = 0. \quad (32)$$

In another paper, Zubair, Mughal & Naqvi give a solution of this kind of Helmholtz equation in fractional space [11]. The Laplacian operator in D-dimensional fractional space is defined as follows:

$$\nabla_D^2 = \frac{\partial^2}{\partial x^2} + \frac{\alpha_1 - 1}{x} \frac{\partial}{\partial x} + \frac{\partial^2}{\partial y^2} + \frac{\alpha_2 - 1}{y} \frac{\partial}{\partial y} + \frac{\partial^2}{\partial z^2} + \frac{\alpha_3 - 1}{z} \frac{\partial}{\partial z}. \quad (33)$$

Then they derive a solution of equation (31) with the help of Bessel equation [11].

It is also interesting to note here, that Shpenkov has suggested that a classical wave equation which is essentially the same with Helmholtz equation can be used to derive a periodic table of elements which is near to Mendeleev's periodic table [12][13]. This result is in contradiction to spherical solution of Schrodinger equation which does not explain any atom, except perhaps hydrogen[14][15]. Therefore it seems worth to discuss what the effect of an extension of classical wave equation in fractional space to the structure of atoms and molecules is. A review of Shpenkov's interpretation and use of classical wave equation has been submitted by this writer [16].

Since the classical wave equation as described by Shpenkov is the same with the equation of vibrating string in 1-dimension, then it seems to compare the solution of equation (31) with solution of fractal vibrating string. A few papers have been written discussing this fractal vibrating string in detail, which can be found elsewhere. [17][18] It appears to be worth to study spherical solution of this fractal vibrating string equation in order to verify Shpenkov's results, including his periodic table of elements. One possible way to find such a solution of fractal vibrating string is to obtain numerical solution of such an equation, by a method of converting fractional differential equation to partial differential equation as proposed by He & Li [19]. After a partial differential equation (PDE) is obtained then it seems not so difficult to find its numerical solution with computer algebra packages like Mathematica, Maple or MatLab.

Here I also propose to investigate further the spherical solution of Helmholtz equation corresponding to Proca-type equations for GEM in fractional space. This kind of investigation may be useful for the study of gravitomagnetic effect and gravitational wave.

Concluding remarks

In a recent paper, M. Zubair et al. described a novel approach for fractional space generalization of the differential electromagnetic equations. A new form of vector

differential operator Del, and its related differential operators, is formulated in fractional space. Using these modified vector differential operators, the classical Maxwell equations have been worked out for fractal media. In the meantime, there are other papers discussing fractional Maxwell equations. However, so far there is no derivation of Proca equations and Proca-type equations for GEM in fractional space. Therefore in this paper I present for the first time a derivation of GravitoElectroMagnetic (GEM) Proca-type equations in fractional space. Considering that Proca equations may be used to explain some electromagnetic effect in superconductor, then fractional GEM Proca-type equations may be expected to explain some gravitomagnetic effects of superconductor for fractal media. It is our hope, that this paper may stimulate further investigation and experiments in particular with respect to gravitomagnetic effects.

Here I also propose to investigate further the spherical solution of Helmholtz equation corresponding to Proca-type equations for GEM in fractional space. This kind of investigation may be useful for the study of gravitomagnetic effect and gravitational wave.

Acknowledgments

Special thanks to Dr. George Shpenkov for sending his papers to this writer. And many thanks to Prof. Akito Takahashi and Prof. L. Wilardjo for their comments on Shpenkov's work.

Version 1.0: April 13th, 2014

VC, email: victorchristianto@gmail.com

References

- [1] Zubair, M., Mughal, M.J., Naqvi, Q.A., & Rizvi, A.A. 2011. Differential Electromagnetic Equations in Fractional Space. *Progress in Electromagnetic Research*, Vol. 114, 255-269. URL: <http://www.jpier.org/PIER/pier114/15.11011403.pdf>
- [2] Tarasov, Vasily E. 2008. Fractional vector calculus and fractional Maxwell equations. *Annals of Physics* 323, 2756-2778, URL: <http://theory.sinp.msu.ru/~tarasov/PDF/AP2008.pdf>; [2a] also in arXiv: 0907.2363 [math-ph], URL: <http://arxiv.org/abs/0907.2363>

- [3] Ostoja-Starzewski, Martin. 2012. Electromagnetism in anisotropic fractal media. *Z. Angew. Math. Phys.*(ZAMP) DOI 10.1007/s00033-012-0230-z . URL: http://mechanical.illinois.edu/media/uploads/web_sites/82/files/zamp_2012.20121002.506b39d642e987.85850489.pdf; [3a] Also in arXiv: 1106.1491 [math-ph], URL: <http://arxiv.org/pdf/1106.1491.pdf>.
- [4] Tajmar, M. 2008. Electrodynamics in Superconductor explained by Proca equations. arXiv:0803.3080
- [5] de Matos, C.J., & Tajmar, M. 2006. Gravitomagnetic London Moment and the Graviton Mass inside a Superconductor. arXiv:gr-qc/0602591
- [6] Gondran, Michel. 2009. Proca equations derived from first principles. arXiv:0901.3300 [quant-ph], URL: <http://arxiv.org/pdf/0901.3300.pdf>
- [7] Blackledge, Jonathan M. 2007. An Approach to Unification using a Linear Systems Model for the Propagation of Broad-band Signals. *ISAST Transaction on Electronics and Signal Processing*, Vol. 1, No. 1, 2007. URL: <http://eleceng.dit.ie/papers/100.pdf>
- [8] Demir, Suleyman. 2013. Space-time algebra for the generalization of gravitational field equations. *Pramana* Vol. 80 No. 5 (Indian Academy of Sciences), May 2013, 811-823. URL: <http://www.ias.ac.in/pramana/v80/p811/fulltext.pdf>
- [9] Schwinger, Julian, DeRaad, Jr., Lester L., Milton, Kimball A., & Tsai, Wu-yang. 1998. *Classical Electrodynamics*. Reading, Massachusetts: Perseus Books. 591 p.
- [10] Penrose, Roger. 2004. *The Road to Reality: A Complete Guide to the Laws of the Universe*. London: Jonathan Cape. 1123 p. URL: <http://staff.washington.edu/freitz/penrose.pdf>
- [11] Zubair, M., Mughal, M.J., & Naqvi, Q.A. 2011. On electromagnetic wave propagation in fractional space. *Nonlinear Analysis: Real World Applications*. doi:10.1016/j.nonrwa.2011.04.010.
- [12] Shpenkov, George P. 2013. *Dialectical View of the World: The Wave Model (Selected Lectures)*. Volume I: Philosophical and Mathematical Background. URL: <http://shpenkov.janmax.com/Vol.1.Dialectics.pdf>
- [13] _____. 2006. An Elucidation of the Nature of the Periodic Law, Chapter 7 in "*The Mathematics of the Periodic Table*", Rouvray, D. H. and King, R. B., ed., Nova Science Publishers, NY, pp. 119-160

- [14] Shpenkov, George P. & Kreidik, Leonid G. 2005. Schrödinger's error in principle. *Galilean Electrodynamics* Vol. 16, No. 3, 51-56, (2005); URL: <http://shpenkov.janmax.com/Blunders.pdf>
- [15] Kreidik, Leonid G., & Shpenkov, George P. 2002. Important Results of Analyzing Foundations of Quantum Mechanics, *Galilean Electrodynamics & QED-EAST*, Vol. 13, Special Issues No. 2, 23-30; URL: <http://shpenkov.janmax.com/QM-Analysis.pdf>
- [16] Christianto, Victor. 2014. The Spherical Solution of Schrodinger equation does not agree with any experiment. Submitted to *Prespacetime Journal*, April 2014. Available at: http://vixra.org/author/Victor_Christianto
- [17] Tarasov, Vasily E. 2005. Wave Equation for Fractal Solid String. *Mod. Phys. Lett. B*, Vol. 19, No. 15, 721-728. URL: <http://theory.sinp.msu.ru/~tarasov/PDF/MPLB2005-1.pdf>
- [18] Hu, Ming-Sheng, Agarwal, Ravi P., & Yang, Xiao-Jun. 2012. Local Fractional Fourier Series with Application to Wave Equation in Fractal Vibrating String. *Abstract and Applied Analysis* Vol. 2012, Article ID 567401, 15p. doi:10.1155/2012/567401. URL: <http://downloads.hindawi.com/journals/aaa/2012/567401.pdf>
- [19] He, Ji-Huan, & Li, Zheng-Biao. 2012. Converting Fractional Differential equations into Partial Differential Equations. *Thermal Science*, Year 2012, Vol. 16, No. 2, 331-334, URL: <http://thermalscience.vinca.rs/pdfs/papers-2011/TSCI110503068H.pdf>