

On the Birch and Swinnerton-Dyer conjecture & ERG Theory

Nicholas R. Wright

Natchitoches, Louisiana, USA

wnicholas2222@yahoo.com

Abstract

We prove the integrality and modularity of the Birch and Swinnerton-Dyer conjecture with ERG Theory. Inspection of the conjecture shows that it is a phenomenological model. Thus, a solution could be found through regression analysis. Numerical verification is possible through nominative determinism/visibility theory. By adding adaptive learning (AL) to the model, the model admits an important time variation in beliefs, which would be ruled out under rational expectations. Entropy can be given from a detailed molecular analysis of the system. In summary, perception consists of the selection, organization, and interpretation of stimuli. These factors affect the conduct of work. We include two inequalities on the log-volume change associated to appropriately chosen deformations.

Keywords: Birch and Swinnerton-Dyer conjecture, ERG Theory, nominative determinism, adaptive learning, belief propagation, number theory

Introduction

The Birch and Swinnerton-Dyer conjecture (BSD), intuitively, specifies the rank of $E(Q)$ will correspond to the value of $L(E,s)$ at 1: the larger r is, the “smaller” $L(E,1)$ is. However, the value of $L(E,s)$ at $s = 1$ does not make sense since the product of $L_p(E,s)$ only converges when $\text{Re}(s) > 3/2$. Nevertheless, if $L(E,s)$ can be continued to an analytic function on the whole of \mathbb{C} , it may be reasonable to believe that the behavior of $L(E,s)$ at $s = 1$ contains the arithmetic information of the rank of $E(Q)$. So let E be an elliptic curve over a global field k , then the order of vanishing of $L(E,s)$ at $s=1$ is equal to the rank of $E(k)$.

Conjecturally there is just one essential type of global L-function, with two descriptions (coming from an algebraic variety and coming from an automorphic representation). We use Nominative Determinism (ND) to show that these two ‘global’ L-functions represent electrical and dynamical systems, respectively. ND is a theory that authors gravitate to the area of research which fits their surname, especially specialties in a field of research. ND can be construed as Mazur’s Visibility Theory. The Hasse principle will support ND as numerical verification of BSD arithmetic rank $r(E)$. We use Mordell-Weil theorem to prove BSD analytic rank $r_{\text{an}}(E)$.

Clayton Alderfer’s ERG theory of motivation will show as a secondary background to BSD. The theory utilizes three frustration-regression (as opposed to satisfaction-progression) levels: Existence, Relatedness (or Relationship), and Growth. ND shows the potential of BSD as the life of a bacteria or sperm: Birth, Swimming, and Death. This pseudo-isolation is congruent with ‘sterbenwold’ folklore scenario. This paper acknowledges that predictive-modeling at the expense of sound reason is a recipe for disaster. Therefore, this paper will focus less on the curse of universality (although explained), and more so on the curse of dimensionality.

These traditional problems often lay unsolved for centuries, and mathematicians gradually came to understand their depth (in some cases), rather than treat them as puzzles. So we will answer the typical questions asked in Diophantine analysis:

1. Are there any solutions?
2. Are there any solutions beyond some that are easily found by inspection?
3. Are there finitely or infinitely many solutions?
4. Can all solutions be found in theory?
5. Can one in practice compute a full list of solutions?

Nominative Determinism (ND)

The nominative is considered a compulsion of the name. It involves the use of heuristics as a decision rule that quickly eliminates alternatives in a bounded rationality model. It is also possible of satisficing, where an alternative is identified as an “acceptable” solution. Using a content theory of motivation such as ERG Theory or Abraham Maslow’s Hierarchy Theory will enhance the solution. As content theories (as opposed to process theories) both the ERG Theory and Maslow’s Hierarchy Theory look at behaviors and other needs that are found within people as opposed to external environmental factors. Moreover, when there are rational solutions there may or may not be infinitely many. In this case, we give them as local bifurcations. The nominative can also appear like spherical mirrors: virtual, erect, and enlarged while concave and diminished while convex. BSD gives us Bryan Birch and Peter Swinnerton-Dyer. We can see that their nominative compulsion is bounded within ERG Theory, but we wish to look more into specialties within their field. Although Birch’s name appears first, we will focus on the elder Swinnerton-Dyer in this exercise and return to Birch later. When considering numbers we come to measurement. Legacy units of measurement have certain advantages. There is the erg, dyne, foe, etc. Considering that we have determined that the nominative is bounded within ERG Theory by ND, we can also assume the erg and dyne. We can also assume the foe, which is shorthand for fifty-one ergs. The nominative compulsion arrives, deepens, and culminates metaphorical or literal. So a foe can be a zero and/or tiny rational number. If we merge this understanding with a criterion like the Néron–Ogg–Shafarevich criterion, we can find isogenous elliptic curves. We can then examine BSD as two “global” L-functions to diffuse, de-fuse, or resolve the conflicts, as opposed to an alternative form of calculus. These “globals” are also prone to stress-related problems. Given this stress, we should expect BSD to be a formula on a molecular (or atomic) magnitude. BSD may serve as an ideal gas formula. Using a regulator (Birch) and period (Swinnerton-Dyer) thought experiment, we proceed to prove that $\text{III}(E)$ is an electrical system while the analytic rank is a dynamical system that can be host to measurements to include Higgs for higher derivatives of $L(E, s)$.

Nekovar Quaternion Algebra B

Turning now to Birch and ND, we first recognize a ‘tree’. Given more quality we may say a log. \ln and \log are logarithms to the base e and base 10. The possibility to use u-substitution integration with $\ln(x)$ becomes overwhelmingly obvious. January Nekovar (2009) has, intuitively, written this out in quaternion algebra and should be referred to from there. So, $r_{\text{an}}(E) \leq 1$. Cursed with universality, unless approached through ND, we have $\log U = \ln i$ or $L(E, 1) = \Omega(E)$. $\log(u)$ or $\exp(u)$ become more probable answers than the polar forms $\ln(i)$ or $(\pi/2)i$. Any other solution will give u-substitution integration with $\ln(x)$ and thus negative numbers. This gives ND apleness. If and only if U , it is assumed

that a voltage V is impressed across. There is no such thing as zero gravity, thus $L(E, 1) \neq 0$. With U as an internal energy we have enough information to plug in several laws of science: Boyle's Law, Charles' Law, Gay-Lussac's Law, Faxen's laws, and Stefan-Boltzmann Law because the emissivity of a blackbody is unity. Even thermal resistance, R , can be a universal gas constant. To explain polar forms, we show that T becomes $\ln(e) = 1$, in electromagnetics the absolute potential at infinity (at $r = \infty$) is zero, and that the unique temperature at which P and V would reach zero is called absolute zero. To understand this in philosophical terms, we ask how far can a dog run into a forest? It would be half minus one. There's an instantaneous point where he's neither running into or out of the forest. The classic sterbenwold scenario is whether a tree in the middle of the forest makes a sound when it falls. Following the ideal gas law, K is called the universal gas constant. If the volume contains m kilograms of gas that has a molecular (or atomic) mass M , then $n = m/M$. Under standard conditions, 1 kmol of ideal gas occupies a volume of 22.4 m^3 . When viewed as an electrical analytical rank, we can also use the right-hand rule of magnetic fields to verify ND without being negligible. With that understanding, we also find how Jerrold B. Tunnell's Theorem explains electrical systems by means of a Pythagorean Theorem for a right triangle and its phasors. ND also explains how: tunnel, tonal, and ton. Even the obsolete measurement called a funal is relevant, its pounds force being similar to the kmol. The important nominative compulsion is: the null.

The Hasse principle

Also known as the local-global principle, the Hasse principle becomes a superb method of defining global L-functions. A more formal version of the Hasse principle states that certain types of equations have a rational solution if and only if they have a solution in the real numbers and in the p-adic numbers for each prime p . Using ND, we know that Hasse means 'hate' in German. We can also find 'heißen'. Heißen means 'to name' or 'to call'. This mimics the two methods of naming a real number and calling a p-adic number. The concept of 'closeness' serves the mathematics well as a form of 'meaning' (Heißen also means 'to mean'). Considering continuously changing magnetic flux, a t-adic number will be necessary and has already been under development. T-adic would represent temperature as utility, and time as power. Victor Kolyvagin (1989) showed that a modular elliptic curve E for which $L(E, 1)$ is not zero has rank 0, and a modular elliptic curve E for which $L(E, 1)$ has a first-order zero at $s = 1$ has rank 1. Remember, G being a one-sided limit is not equal to 0. Kolyvagin is apparently examining temperature in respect to time-space, mass, energy, and gravity.

Adaptive Learning (AL)

Since BSD is essentially meaningless, ND is as much of an ample tool and has probably been used unknowingly under several names. A second ample tool is Adaptive Learning (AL), and is sometimes known as Belief Propagation (BP). It is a statistical formula to construct artificial intelligence. AL is particularly of value in politics and economics. Several popular formulas include the Metropolis-Hastings algorithm, the Metropolis algorithm, and Gibbs sampling. What all of these formulas share is Bayes Law: $P(x|z) \sim L(x;z)P(x)$. John Tate once remarked: Let $A = A_f$ be a modular abelian variety attached to a new-form, and let $r = \text{ord}_{s=1} L(A, s)$ be the analytic rank of A , then for arbitrary abelian varieties the value of a function relates where it is not known to be defined to the order of a group that is not known to be finite. The logic is congruent with AL. If $a > 1$ accept $x^{(t+1)} = x'$ else accept with probability a if rejected: $x^{(t+1)} = x^{(t)}$. In other words, we argue that if agents are assumed to possess a slightly larger amount of knowledge, by correctly noticing that $C_t = Y_t$ holds at all times, the aggregate laws of motion would reduce to those obtained under rational expectations. Because of Faltings's theorem, this is false unless $C = A$. Therefore, we have $C = A = C_t = Y_t$. For sociologist, "culture" counts as an independent force alongside of "structure". That is when we have terms as culture of poverty, culture of corruption, culture of dependency, etc. Here the regimes are assumed to be observed; therefore, the estimation of the model basically reduces to an estimation with an added dummy variable (S_t). As a rough rule of thumb, the distribution of k tilts toward 0 when $S_t = 1$. Showing, repetition is the father of learning.

Dark Globe Theory

Consistent with the Taniyama-Shimura conjecture and Mordell-Weil theorem, BSD conjecture should be raised to the status of an axiom. The BSD Axiom implies a proof of several equivalent fundamental conjectures in Diophantine geometry, including the *abc* conjecture over any number field. The axiom extends substantially the scope of arithmetic geometry. It may in fact be called Dark Globe Theory is some parts of the world (Germanic-speaking Europe). With this in mind we presume that the axiom will take on additional features; such as the "Swampland" concept, DNA supercoils, and Dark Flow. BSD teaches us to think of the world with two branes, and only two branes, in the same vein that Edward Witten (M-theory) taught us to think in multiple dimensions. House's Path-Goal Theory is a dyadic theory of supervision that applies. It concerns the relationship between supervisor, subordinate, and satisfaction. Here, we have the vertical dyad linkages again. It is a p-adic/real number relation. In some respects, the basis of BSD is not to find why $3 + 2 = 5$, but why is 2 an invented number. This new axiom would imply superparticles, chemical reaction, nuclear reaction, and radioactive decay. Even conduction, convection, radiation, and redox (reduction-oxidation reaction). Moreover, it can be shown that P must be of

infinite order. Conversely, such a point of infinite order gives us back a rational triple (a,b,c) . Thus showing that n is a congruent number is equivalent to showing that the elliptic curve E_n has positive rank. Since the null holds true, the converse is also true if we further assume the BSD axiom. Therefore the proof of the BSD conjecture will lead to a solution to the congruent number problem using a finite amount of computation. D. R. Heath-Brown (2004) explains the conditions of dynamical systems: heat and Brownian motion. Mazur-Rubin (2010) proved that if the BSD conjecture holds for all elliptic curves over all number fields, then Hilbert's 10th problem has a negative answer over O_k for any number field K . Therefore, BSD only holds for the lower dynamical systems and not the upper electrical systems (e.g. foe, erg, dyne, etc.). John William Scott Cassels (1965) proves schematics to statistics. Noam Elkies (2006) proves ethics to physics. The Parity conjecture proves the dyne and Paul Monsky (1996) proved determinism. Interestingly, Dokchitser-Dokchitser (2009) is p-parity proving the T-symmetry. Further, like a micro standard deviation, the Gross-Zagier (1986) formula relating the central derivative $L'(E,1)$ and the heights of Heegner points on E defined over an imaginary quadratic field could be another way of saying Higgs boson. Finally, Jan Nekovar (2006) has proven real numbers proving the rationality of Hasse. In Czech, Nekovar means "a smith who is better to keep out of the way". The analytic rank of A_f is equal to the Mordell-Weil rank of A_f .

Conclusion

We find that elliptic curve E over global field k is a finite measurement of erg (energy and work equal to 10^{-7} joules) and the order of $L(E,s)$ at $s = 1$ is equal to the rank of $E(k)$. Due to global field k having qualities of infinity, BSD conjecture would imply that *neither* elliptic curve E nor global field k as having a solution. Inspection of the conjecture shows that it is a phenomenological model. Thus, a solution could be found using regression analysis. This paper has found a solution using regression analysis. Raising BSD conjecture to the status of axiom will eliminate the human side of the equation, therefore allowing one to compute a full list of solutions.

Conflict of Interest

The author claims no conflict of interest.

References

- Abel, Ernest L. (2010). "Influence of Names on Career Choices in Medicine". *Names: A Journal of Onomastics* 58 (2): 65–74. doi:10.1179/002777310X12682237914945.
- Arnolds, C.A. & Boshoff, C. (2002). Compensation, esteem valence and job performance: An empirical assessment of Alderfer's ERG theory. *International Journal of Human Resource Management*, 13(4), pp. 697-719. Database: Business Source Elite
- Cremona, John (2011). "Numerical evidence for the Birch and Swinnerton-Dyer Conjecture". Talk at the BSD 50th anniversary conference, May 2011.
- Dellaert, Frank (2005) "Markov chain Monte Carlo Basics". ICCV05 Tutorial: MCMC for Vision. Retrieved from http://vcla.stat.ucla.edu/old/MCMC/MCMC_tutorial/Lect2_Basic_MCMC.pdf
- Heath-Brown, D. R. (2004). "The Average Analytic Rank of Elliptic Curves". *Duke Mathematical Journal* 122 (3): 591–623. doi:10.1215/S0012-7094-04-12235-3. MR 2057019.
- Kolyvagin, Victor (1989). "Finiteness of $E(Q)$ and $X(E, Q)$ for a class of Weil curves". *Math. USSR Izv.* 32: 523–541. doi:10.1070/im1989v032n03abeh000779.
- Limb, C.; Limb, R.; Limb, C.; Limb, D. (2015). "Nominative determinism in hospital medicine". *The Bulletin* 97 (1): 24–26. doi:10.1308/147363515X14134529299420.
- Nekovář, Jan (2009). "On the parity of ranks of Selmer groups IV". *Compositio Mathematica* 145 (6): 1351–1359. doi:10.1112/S0010437X09003959.
- Rubin, Karl (1991). "The 'main conjectures' of Iwasawa theory for imaginary quadratic fields". *Inventiones Mathematicae* 103 (1): 25–68. doi:10.1007/BF01239508. Zbl 0737.11030.
- Stein, William (2006). "Studying the Birch and Swinnerton-Dyer conjecture for modular abelian varieties using Magma". *Discovering Mathematics with Magma*, Springer. pp. 93-116. Print ISBN 978-3-540-37632-3. Online ISBN 978-3-540-37634-7.
- Tunnell, Jerrold B. (1983). "A classical Diophantine problem and modular forms of weight $3/2$ ". *Inventiones Mathematicae* 72 (2): 323–334. doi:10.1007/BF01389327. Zbl 0515.10013.
- Wiles, Andrew (2006). "The Birch and Swinnerton-Dyer conjecture". In Carlson, James; Jaffe, Arthur; Wiles, Andrew. *The Millennium prize problems*. American Mathematical Society. pp. 31–44. ISBN 978-0-8218-3679-8. MR 2238272.