

# Fermi's golden rule: its derivation and breakdown by an ideal model

J. M. Zhang<sup>1,2,\*</sup>

<sup>1</sup>*Fujian Provincial Key Laboratory of Quantum Manipulation and New Energy Materials,  
College of Physics and Energy, Fujian Normal University, Fuzhou 350007, China*

<sup>2</sup>*Fujian Provincial Collaborative Innovation Center for Optoelectronic  
Semiconductors and Efficient Devices, Xiamen, 361005, China*

Fermi's golden rule is of great importance in quantum dynamics. However, in many textbooks on quantum mechanics, its content and limitation are obscured by the approximations and arguments in the derivation, which are inevitable because of the generic setting considered. Here we propose to introduce it by an ideal model, in which the quasi-continuum band consists of equal-distant levels extending from  $-\infty$  to  $+\infty$ , and each of them couples to the discrete level with the same strength. For this model, the transition probability in the first order perturbation approximation can be calculated analytically by invoking the Poisson summation formula. It turns out to be a *piecewise linear* function of time, demonstrating on one hand the key features of Fermi's golden rule, and on the other hand that the rule breaks down beyond the Heisenberg time, even when the first order perturbation approximation itself is still valid.

## I. INTRODUCTION

As an example of Stigler's law of eponymy, the so-called Fermi's golden rule was actually first derived by Dirac<sup>5</sup> instead of Fermi, although the title "golden" was given by the latter.<sup>2</sup> It is no wonder that Fermi considered the rule a golden one in view of its instrumental role in his theory of Beta decay.<sup>3,4</sup>

As a nontrivial result of the lowest order time-dependent perturbation theory, Fermi's golden rule is introduced in almost every textbook on quantum mechanics. It is about the transition dynamics for such a scenario. Initially, the system is in some eigenstate  $|b\rangle$  of an unperturbed Hamiltonian  $H_0$  with eigenenergy  $E_b$ . Besides the level  $|b\rangle$ ,  $H_0$  has a quasi-continuum  $\{|n\rangle\}$  with eigenenergies  $\{E_n\}$ . It is assumed that  $E_n$  is non-degenerate and increases with  $n$ , and that the quasi-continuum covers an interval  $[a, b]$ , to which  $E_b$  belongs, i.e.,  $a < E_b < b$ . Then at  $t = 0$  a perturbation  $V$  is turned on, which couples  $|b\rangle$  and  $\{|n\rangle\}$  (and possibly, states within the continuum band too, say  $|n_1\rangle$  and  $|n_2\rangle$ ) with strength  $g_n = \langle n|V|b\rangle$ . It is assumed that  $g_n$  is a slowly varying function of  $n$ , and therefore, it is legitimate to introduce a continuous function  $g(E)$  such that  $g_n = g(E_n)$ . Because of the newly introduced coupling, the system transits towards the quasi-continuum. Fermi's golden rule then states that, in the first order perturbation theory, the probability  $P$  of finding the system in the continuum grows linearly in time, and the rate of increase, the so-called transition rate, is of the expression

$$w = \frac{dP}{dt} = \frac{2\pi}{\hbar} |g(E_b)|^2 \rho(E_b). \quad (1)$$

Here  $\rho(\cdot)$  is the density of states of the quasi-continuum. It is defined as  $\rho(E)dE$  is the number of levels in the interval  $[E, E + dE]$ .

In many textbooks on quantum mechanics, formula (1) is derived for a generic case. In other words, no concrete constraint is placed on the quasi-continuum spectrum

$\{E_n\}$  or the couplings  $\{g_n\}$ , except for the conditions above, which are either implicitly or explicitly assumed. The generality of this approach is definitely appreciable, but exactly because of its generality, many approximations and arguments (within the first order perturbation framework) are inevitable, which obscure the content and limitation of the rule, especially for a novice. For example, in deriving (1), in most if not all textbooks,<sup>5-14</sup> the limit of  $t \rightarrow \infty$  is taken. However, intuitively, one would not expect the first order perturbation theory to hold any more in the long time limit of  $t \rightarrow \infty$ . Actually, to be consistent with the first order perturbation theory, the condition  $P \ll 1$  should be satisfied, and the linear behavior of (1) cannot last forever. Although this dilemma is just superficial (the  $t \rightarrow \infty$  limit is essentially achieved already for a short and finite  $t$ ), it does stand as an obstacle for the beginners.

It is therefore desirable to have a model for which the calculation can be done as rigorously and as straightforwardly as possible. Such a model indeed exists, and as a matter of fact, it has been known for decades.<sup>16</sup> Yet, unfortunately, to the best knowledge of the author, it has not been introduced into any quantum mechanics textbook, neither in the main text nor in the exercise part. The model is very simple—in hindsight, it can actually be motivated by the expression (1). The transition rate  $w$  is proportional to the local values of the density of states and the coupling strength squared. Hence, in the model, the level spacing  $E_{n+1} - E_n$  and the coupling  $g_n$  are simply taken as constant. For this model, the transition probability in the first order approximation can be calculated rigorously for arbitrary time  $t$ . It is actually a piecewise linear function of time, showing kinks with a period of  $t_H$ , the Heisenberg time associated with the spectrum. Therefore, on one hand, the approximations and arguments in the general case, in particular, the  $t \rightarrow \infty$  limit, are avoided; on the other hand, it presents an example demonstrating vividly that Fermi's golden rule breaks down beyond the Heisenberg time, even when the first order approximation itself is still valid. As far

as the author knows, the Heisenberg time as an upper bound for the validity domain of Fermi's golden rule is pointed out only by Baym.<sup>14</sup> In many books, it is simply given by the non-depletion condition  $t \ll 1/w$ .<sup>5,6,8,12</sup>

In the following, we shall first present the general formalism of the first order time-dependent perturbation theory and derive Fermi's golden rule in the generic case in Sec. II. Then, we proceed to consider the special case of the ideal model in Sec. III. It is interesting that the model is not only solvable up to the first order approximation, but can be solved exactly by summing up all orders of terms. For completeness, this is done in Sec. IV.

## II. THE GENERIC CASE

Let us first review the derivation of (1) in the generic case. The time dependent Schrödinger equation is

$$i\hbar \frac{\partial |\psi\rangle}{\partial t} = (H_0 + \lambda V)|\psi\rangle. \quad (2)$$

The initial condition is  $|\psi(t=0)\rangle = |b\rangle$ . Here we have introduced a control parameter  $\lambda$ , whose value will be set to 1 in the end. Expand the wave function  $|\psi(t; \lambda)\rangle$  as a power series of  $\lambda$ ,

$$|\psi(t; \lambda)\rangle = \sum_{s=0}^{\infty} \lambda^s |\psi_s(t)\rangle. \quad (3)$$

By the initial condition of  $|\psi\rangle$ , we have  $|\psi_0(0)\rangle = |b\rangle$  and  $|\psi_s(0)\rangle = 0$  for  $s \geq 1$ . Plugging (3) into (2), and equating the coefficients of the powers of  $\lambda$ , we get up to  $s = 1$ ,

$$i\hbar \frac{\partial |\psi_0\rangle}{\partial t} = H_0 |\psi_0\rangle, \quad (4a)$$

$$i\hbar \frac{\partial |\psi_1\rangle}{\partial t} = H_0 |\psi_1\rangle + V |\psi_0\rangle. \quad (4b)$$

From (4a) we solve  $|\psi_0(t)\rangle = e^{-iE_b t/\hbar} |b\rangle$ . Substituting this result into (4b), and projecting both sides of (4b) onto  $|n\rangle$ , we get

$$i\hbar \frac{\partial}{\partial t} \langle n | \psi_1 \rangle = E_n \langle n | \psi_1 \rangle + g_n e^{-iE_b t/\hbar}. \quad (5)$$

By Duhamel's principle,<sup>15</sup> we then easily solve

$$\begin{aligned} \langle n | \psi_1(t) \rangle &= \frac{1}{i\hbar} \int_0^t d\tau g_n e^{-iE_b \tau/\hbar} e^{-iE_n(t-\tau)/\hbar} \\ &= g_n \frac{1 - e^{i(E_n - E_b)t/\hbar}}{E_n - E_b} e^{-iE_n t/\hbar}. \end{aligned} \quad (6)$$

Therefore, to the first order approximation, the probability of finding the system in the quasi-continuum band

is

$$\begin{aligned} P &= \sum_n |\langle n | \psi_1(t) \rangle|^2 \\ &= \sum_n |g_n|^2 \frac{4 \sin^2((E_n - E_b)t/2\hbar)}{(E_n - E_b)^2} \\ &= \frac{2t}{\hbar} \sum_n |g_n|^2 \frac{\sin^2((E_n - E_b)t/2\hbar)}{(E_n - E_b)^2 (t/2\hbar)}. \end{aligned} \quad (7)$$

Under certain condition, the summation can be approximated by an integral

$$P = \frac{2t}{\hbar} \int_a^b d\epsilon \rho(\epsilon) |g(\epsilon)|^2 \frac{\sin^2((\epsilon - E_b)t/2\hbar)}{(\epsilon - E_b)^2 (t/2\hbar)}. \quad (8)$$

The weight function (as a function of  $\epsilon$ ) in the integral

$$\frac{\sin^2((\epsilon - E_b)t/2\hbar)}{(\epsilon - E_b)^2 (t/2\hbar)} \quad (9)$$

consists of lumps whose width scales as  $1/t$ , and whose height scales as  $t$ . In the limit of  $t \rightarrow \infty$ , it converges to the delta function  $\pi\delta(\epsilon - E_b)$ . In this limit,

$$P = \frac{2\pi t}{\hbar} \rho(E_b) |g(E_b)|^2. \quad (10)$$

We then get the golden rule (1). Here we have invoked two approximations ensuing (7), which is an approximation by itself. From (7) to (8), in replacing the summation by an integral, the sampling step-length (i.e., level spacing  $E_{n+1} - E_n$ ) should be much smaller than the characteristic length of variation of the function, which is  $2\pi\hbar/t$ . That is,

$$t \ll 2\pi\hbar\rho(E_b). \quad (11)$$

From (8) to (10), in replacing the weight function (9) by the delta function  $\pi\delta(\epsilon - E_b)$ , the width of the lumps of the former should be much smaller than the width of the interval  $[a, b]$ , or more accurately,

$$\frac{2\pi\hbar}{\min\{|a - E_b|, |b - E_b|\}} \ll t. \quad (12)$$

Both conditions of (11) and (12) are within the first order perturbation approximation itself, and can be satisfied simultaneously. We note that while condition (12) is mentioned in many books,<sup>5,6,8,14</sup> condition (11) is pointed out only by Baym.<sup>14</sup> As we shall see in the ideal model below, the right hand side of (11) defines sharply the boundary beyond which Fermi's golden rule breaks down.

## III. THE IDEAL MODEL

In retrospect, the ideal model can be motivated by the expression of the golden rule (1) or (10). The transi-




FIG. 1: (Color online) Periodic sampling and summation of the function  $\text{sinc}^2 x \equiv \sin^2 x/x^2$ , which is the essential part in the definition of  $W_\alpha(T)$  [see Eq. (16)]. The sampling period is  $T$ , and the shift is determined by  $\alpha$ . In this figure,  $\alpha = 3/7$ .

tion rate is proportional to the local values of the density of state  $\rho(E)$  and the square of the coupling  $|g(E)|^2$  at  $E = E_b$ . In the ideal model, both functions are simply taken as constant. That is,  $E_n = n\Delta$  with  $\Delta$  being the spacing between two adjacent levels, and  $g_n = g$ . More specifically, the unperturbed Hamiltonian and the coupling are, respectively,<sup>16</sup>

$$H_0 = E_b|b\rangle\langle b| + \sum_{n=-\infty}^{\infty} n\Delta|n\rangle\langle n|, \quad (13a)$$

$$V = g \sum_{n=-\infty}^{\infty} (|b\rangle\langle n| + |n\rangle\langle b|). \quad (13b)$$

Note that the quasi-continuum extends from  $-\infty$  to  $+\infty$ , and we have assumed that the coupling strength  $g$  is real, which is always achievable by adjusting the phases of the levels  $|n\rangle$ . This model can be approximately realized with a two-level atom in a multi-mode cavity.<sup>17–20</sup> In this system, the discrete level  $|b\rangle$  corresponds to the atom in the excited state and all cavity modes empty, while each  $|n\rangle$  corresponds to the atom in the ground state and the  $n$ th cavity mode being excited. The mode spacing and the atom-mode couplings (if the atom is appropriately located) are, to a good extent, constant. Only those modes near resonant with the atomic resonance frequency participate significantly in the dynamics. Therefore, it is legitimate to assume that there are infinite number of cavity modes extending from  $-\infty$  to  $+\infty$ .

For the concrete model (13), the general formula (7) reduces to periodic sampling and summation of the function  $\text{sinc}^2 x \equiv \sin^2 x/x^2$ . Specifically, introducing the dimensionless time  $T \equiv \Delta t/2\hbar$  and the offset parameter (here  $[\cdot]$  is the floor function)

$$\alpha = E_b/\Delta - [E_b/\Delta], \quad (14)$$

which characterizes the location of  $E_b$  relative to the quasi-continuum spectrum, the transition probability can be written as

$$P = \left(\frac{4g^2}{\Delta^2}\right) W_\alpha(T). \quad (15)$$

Here the function  $W_\alpha$  is defined as

$$W_\alpha(T) \equiv T^2 \sum_{m=-\infty}^{\infty} \text{sinc}^2[(m - \alpha)T], \quad (16)$$

where the infinite summation is a very regular one—it samples the  $\text{sinc}^2 x$  function uniformly with the period given by  $T$  and the offset determined by  $\alpha$  (see Fig. 1). It is apparent that  $W_\alpha = W_{-\alpha} = W_{1-\alpha}$ , i.e.,  $W_\alpha$  is an even and periodic function of  $\alpha$ . Note that in (15), the time dependence is only through the function  $W_\alpha$  and the coupling strength  $g$  appears only in the prefactor.

The primary concern is to calculate  $W_\alpha$ . For this purpose, we have the standard tool of Poisson summation formula.<sup>21</sup> By this formula, a periodic sampling and summation of a function  $f(x)$ ,

$$I = \sum_{n=-\infty}^{+\infty} f(a + nT) \quad (17)$$

can be converted to a weighted periodic sampling of its Fourier transform

$$F(q) = \int_{-\infty}^{+\infty} dx f(x) e^{-iqx}. \quad (18)$$

That is,

$$I = \frac{1}{T} \sum_{n=-\infty}^{+\infty} F\left(\frac{2\pi n}{T}\right) \exp\left(\frac{i2\pi na}{T}\right). \quad (19)$$

In our case, we need to calculate the Fourier transform of the function  $\text{sinc}^2 x$ . For those who are familiar with the Fraunhofer diffraction of a single slit,<sup>22</sup> it is a basic mathematical fact that the Fourier transform  $F_1$  of the function  $\text{sinc} x$  is the window function.<sup>23,24</sup> Namely,

$$F_1(q) = \begin{cases} \pi, & |q| \leq 1, \\ 0, & |q| > 1. \end{cases} \quad (20)$$

By the convolution theorem, the Fourier transform  $F_2$  of  $\text{sinc}^2 x$  is then simply the self-convolution of  $F_1$ ,

$$\begin{aligned} F_2(q) &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dp F_1(p) F_1(q - p) \\ &= \begin{cases} \frac{\pi}{2}(2 - |q|), & |q| \leq 2, \\ 0, & |q| > 2, \end{cases} \end{aligned} \quad (21)$$

which is a triangle function nonvanishing only on the

support  $[-2, 2]$ . The fact that the Fourier transform of  $\text{sinc}^2 x$  has only a finite support is actually a consequence of the Paley-Wiener theorem.<sup>25</sup> The function  $\text{sinc}^2 x$  is an entire function and is of exponential type 2.<sup>26</sup> Hence, by the Paley-Wiener theorem, its Fourier transform is supported on  $[-2, 2]$ .

The fact that  $F_2$  is nonvanishing only on a finite interval means that invoking the Poisson summation formula really simplifies the original summation problem in (16), because in (19) there will be only a finite number of nonzero terms. For example, if  $0 < T \leq \pi$ , only the  $n = 0$  term is nonzero, and consequently, in this interval,

$$W_\alpha(T) = \pi T, \quad (22)$$

regardless of the value of  $\alpha$ . In terms of the real time  $t$ , the result is that, for  $0 < t < t_H \equiv 2\pi\hbar/\Delta$ , the Heisenberg time,

$$P = \frac{2\pi}{\hbar} \left( \frac{g^2}{\Delta} \right) t. \quad (23)$$

This is nothing but Fermi's golden rule. We have obtained it without using any approximation or argument. The fact that it is independent of the parameter  $\alpha$  is in accord with the coarse-graining spirit implicitly assumed in the usual derivation of the rule.

However, in the more general case of  $m\pi < T \leq (m+1)\pi$ , there are  $2m+1$  nonzero terms in the summation. It is straightforward to get ( $\theta \equiv 2\pi\alpha$ )

$$\begin{aligned} W_\alpha(T) &= \pi \left( \sum_{n=-m}^m \exp(in\theta) \right) T - 2\pi^2 \sum_{n=1}^m n \cos(n\theta) \\ &= \pi \frac{\sin \frac{(2m+1)\theta}{2}}{\sin \frac{\theta}{2}} T - 2\pi^2 \frac{\partial}{\partial \theta} \left( \sum_{n=1}^m \sin n\theta \right) \\ &= \pi \frac{\sin \frac{(2m+1)\theta}{2}}{\sin \frac{\theta}{2}} (T - m\pi) + \pi^2 \frac{\sin^2 \frac{m\theta}{2}}{\sin^2 \frac{\theta}{2}}. \end{aligned} \quad (24)$$

This formula was first obtained by Kyrölä using a different method.<sup>27</sup> We see that  $W_\alpha$  is still a linear function of  $T$  on the interval  $[m\pi, (m+1)\pi]$ , but its slope now depends on both the interval and the offset parameter  $\alpha$ . Therefore,  $W_\alpha$  is a piecewise linear function of  $T$  and has kinks at  $T = m\pi$  (correspondingly,  $t = mt_H$ ) periodically. The point is that, it is continuous but nonsmooth. Apparently, by making  $g$  sufficiently weak, one can make  $P$  arbitrarily small at an arbitrary time  $t$ . Therefore, for a sufficiently small  $g$ , the first order perturbation approximation can be valid far beyond the critical time  $t_H$ . Yet, Fermi's golden rule is no longer valid there.

In Fig. 2, the function  $W_\alpha(T)$  is plotted for four different values of  $\alpha$ . Before  $t_H$ , all curves coincide as (22) says; but beyond  $t_H$ , the curves fan out and have completely different trajectories. Except for the case of  $\alpha = 0$ , namely when  $E_b$  is degenerate with a certain level  $|n\rangle$ ,  $W_\alpha$  shows collapses and revivals. Actually, by (24), at




FIG. 2: (Color online) The function  $W_\alpha(T)$  [see Eq. (16)] for four different values of  $\alpha$ . Each curve is piecewise linear and all the curves coincide in the first interval of  $0 \leq T \leq \pi$ . Note the two particular cases of  $\alpha = 0$  and  $\alpha = 0.5$ . For the former,  $W_\alpha$  diverges quadratically with  $T$ ; for the latter,  $W_\alpha$  returns to zero (as a phenomenon of collapse and revival) repeatedly.

$$T = m\pi,$$

$$W_\alpha(m\pi) = \pi^2 \frac{1 - \cos m\theta}{1 - \cos \theta}. \quad (25)$$

Thus for a generic value of  $\alpha$ ,  $W_\alpha(m\pi)$  is an almost periodic function of  $m$ .

We note that the piecewise linear behavior of the transition probability as illustrated in Fig. 2 has been numerically observed in the contexts of photoexcitation of a molecule,<sup>27</sup> spontaneous decay of a two-level atom in a multi-mode optical cavity,<sup>17,18</sup> and transition dynamics of a Bloch state in a periodically driven tight binding model.<sup>28</sup> In the third of these systems, the model (13) is actually not realized exactly. Except for the superficial time dependence, the perturbation  $V$  contains intra-continuum terms, and the level spacing is not strictly constant. But to the first order of  $g$ , the intra-continuum couplings do not enter the dynamics, and only the Bloch states near resonance play an important role in the dynamics, for which the equidistant level spacing condition is a good approximation. Therefore, the perturbation theory above still holds.

#### IV. EXACT SOLUTION OF THE TRANSITION PROBLEM

The transition problem for the ideal model can actually be solved exactly. This was originally done by Stey and Gibberd using the Laplace transform technique<sup>16</sup> and later by Lefebvre and Savolainen using the memory function method.<sup>29</sup> In the following, a solution in line with the perturbation expansion (3) will be presented.

To determine  $|\psi_s(t)\rangle$  in (3), let us transform into the in-

interaction picture.<sup>10</sup> Define  $|\psi(t)\rangle = e^{-iH_0t/\hbar}|\tilde{\psi}(t)\rangle$ . The time evolution equation for  $|\tilde{\psi}(t)\rangle$  is

$$i\hbar\frac{\partial}{\partial t}|\tilde{\psi}(t)\rangle = \tilde{V}(t)|\tilde{\psi}(t)\rangle, \quad (26)$$

with

$$\begin{aligned} \tilde{V}(t) &= e^{iH_0t/\hbar}V e^{-iH_0t/\hbar} \\ &= g \sum_{n=-\infty}^{\infty} \left( |b\rangle\langle n|e^{i(E_b-E_n)t/\hbar} + \text{h.c.} \right). \end{aligned} \quad (27)$$

The time evolution operator in the interaction picture can be constructed formally as

$$\begin{aligned} S(t) &= I + \int_0^t dt_1 \left( \frac{1}{i\hbar} \tilde{V}(t_1) \right) \\ &+ \int_0^t dt_1 \int_{t_1}^t dt_2 \left( \frac{1}{i\hbar} \tilde{V}(t_2) \right) \left( \frac{1}{i\hbar} \tilde{V}(t_1) \right) + \dots \end{aligned} \quad (28)$$

This is the so-called Dyson series, in which the  $s$ th term corresponds to the  $s$ th term in (3). The matrix element  $S_{bb}(t) \equiv \langle b|S(t)|b\rangle$  is of the form

$$S_{bb}(t) = \sum_{n=0}^{\infty} \left( \frac{g}{i\hbar} \right)^{2n} C_n, \quad (29)$$

with

$$\begin{aligned} C_n &= \sum_{m_1=-\infty}^{\infty} \dots \sum_{m_n=-\infty}^{\infty} \int_0^t dt_1 \int_{t_1}^t dt_2 \dots \int_{t_{2n-1}}^t dt_{2n} e^{i\sum_{j=1}^n (E_b-E_{m_j})(t_{2j}-t_{2j-1})/\hbar} \\ &= \int_0^t dt_1 \int_{t_1}^t dt_2 \dots \int_{t_{2n-1}}^t dt_{2n} \sum_{m_1=-\infty}^{\infty} \dots \sum_{m_n=-\infty}^{\infty} e^{i\sum_{j=1}^n (E_b-E_{m_j})(t_{2j}-t_{2j-1})/\hbar}. \end{aligned} \quad (30)$$

Note that in (29) the odd terms in  $g$  drop out. The reason is simply that the system has to jump even times between the discrete level and the quasi-continuum band to return to the discrete level. Once we have calculated  $S_{bb}(t)$ , we obtain the survival probability  $P_i$  as

$$\begin{aligned} P_i &= |\langle b|e^{-iHt/\hbar}|b\rangle|^2 = |\langle b|e^{-iH_0t/\hbar}S(t)|b\rangle|^2 \\ &= |\langle b|e^{-iE_b t/\hbar}S(t)|b\rangle|^2 = |S_{bb}(t)|^2. \end{aligned} \quad (31)$$

It is ready to see that the dependence of  $P_i$  on  $E_b$  should be through the parameter  $\alpha$ .

In the second line of (30), we need to do the summation

$$\sum_{m=-\infty}^{\infty} e^{im\Delta t/\hbar}. \quad (32)$$

Again, the Poisson summation formula is useful. The

function relevant is  $f(x) = e^{ix}$ , which is sampled with a period of  $\Delta t/\hbar$ . The Fourier transform of  $f$  is

$$F(q) = \int_{-\infty}^{\infty} dx e^{ix} e^{-iqx} = 2\pi\delta(q-1). \quad (33)$$

Therefore by the formula (19) ( $t_H = 2\pi\hbar/\Delta$ ),

$$\begin{aligned} \sum_{m=-\infty}^{\infty} e^{im\Delta t/\hbar} &= \frac{\hbar}{\Delta t} \sum_{m=-\infty}^{\infty} 2\pi\delta\left(\frac{2\pi m\hbar}{\Delta t} - 1\right) \\ &= t_H \sum_{m=-\infty}^{\infty} \delta(t - mt_H). \end{aligned} \quad (34)$$

Substituting (34) into (30), we get

$$\begin{aligned}
C_n &= t_{\text{H}}^n \int_0^t dt_1 \int_{t_1}^t dt_2 \cdots \int_{t_{2n-1}}^t dt_{2n} \prod_{j=1}^n \left( e^{iE_b(t_{2j}-t_{2j-1})/\hbar} \sum_{m_j=-\infty}^{\infty} \delta[(t_{2j}-t_{2j-1})-m_j t_{\text{H}}] \right) \\
&= t_{\text{H}}^n \int_0^t dt_1 \int_{t_1}^t dt_2 \cdots \int_{t_{2n-1}}^t dt_{2n} \prod_{j=1}^n \left( e^{iE_b(t_{2j}-t_{2j-1})/\hbar} \sum_{m_j=0}^{\infty} \delta[(t_{2j}-t_{2j-1})-m_j t_{\text{H}}] \right) \\
&= t_{\text{H}}^n \sum_{m_1=0}^{\infty} \sum_{m_2=0}^{\infty} \cdots \sum_{m_n=0}^{\infty} \int_0^t dt_1 \int_{t_1}^t dt_2 \cdots \int_{t_{2n-1}}^t dt_{2n} \prod_{j=1}^n \left( e^{iE_b(t_{2j}-t_{2j-1})/\hbar} \delta[(t_{2j}-t_{2j-1})-m_j t_{\text{H}}] \right). \quad (35)
\end{aligned}$$

Now suppose  $0 < t < t_{\text{H}}$ . In this case,  $0 \leq t_{2j} - t_{2j-1} < t_{\text{H}}$ , and hence the only contributing term would be  $m_j = 0$ ,  $j = 1, \dots, n$ . Integrating out  $t_{2j}$  and noting that  $\int_0^{\infty} dt \delta(t) = 1/2$ , we get

$$\begin{aligned}
C_n &= t_{\text{H}}^n \int_0^t dt_1 \int_{t_1}^t dt_3 \cdots \int_{t_{2n-3}}^t dt_{2n-1} \left( \frac{1}{2} \right)^n \\
&= \frac{1}{n!} \left( \frac{t_{\text{H}} t}{2} \right)^n. \quad (36)
\end{aligned}$$

By (29), this leads to

$$S_{bb}(t) = \exp\left(-\frac{\pi g^2}{\Delta \hbar} t\right) = \exp\left(-\frac{\gamma}{2} t\right), \quad (37)$$

where  $\gamma \equiv 2\pi g^2/\Delta \hbar$ . It is purely exponential. Note that as in (22), there is no dependence on  $\alpha$ .

Next suppose  $t_{\text{H}} < t < 2t_{\text{H}}$ . In this case, besides the possibility of  $m_j \equiv 0$  for all  $j$ , one  $m_j$  can be 1. The contribution of the first possibility to  $C_n$  is given by (37). In the second possibility, we make the change of variables,  $(t_1, t_2, \dots, t_{2n}) \rightarrow (s_1, s_2, \dots, s_{2n})$ ,

$$s_i = t_i, \quad i \leq 2j-1, \quad (38a)$$

$$s_i = t_i - t_{\text{H}}, \quad i > 2j-1. \quad (38b)$$

The Jacobian is apparently 1. The corresponding contribution to  $C_n$ , after integrating out  $s_{2j}$ , is ( $\theta = 2\pi\alpha$ )

$$\begin{aligned}
&n t_{\text{H}}^n \int_0^{t-t_{\text{H}}} ds_1 \cdots \int_{s_{2n-3}}^{t-t_{\text{H}}} ds_{2n-1} \left( \frac{1}{2} \right)^{n-1} e^{iE_b t_{\text{H}}/\hbar} \\
&= \frac{t_{\text{H}}^n (t-t_{\text{H}})^n}{(n-1)! 2^{n-1}} e^{i\theta}. \quad (39)
\end{aligned}$$

By (29), the corresponding contribution to  $\langle b|S(t)|b \rangle$  is then

$$-\gamma(t-t_{\text{H}})e^{-\gamma(t-t_{\text{H}})/2} e^{i\theta}. \quad (40)$$

In total, in the interval of  $t_{\text{H}} < t < 2t_{\text{H}}$ ,

$$S_{bb}(t) = e^{-\gamma t/2} - \gamma(t-t_{\text{H}})e^{-\gamma(t-t_{\text{H}})/2} e^{i\theta}. \quad (41)$$

Therefore, beyond  $t_{\text{H}}$ ,  $P_i$  is no longer purely exponential and is dependent on  $\alpha$ . Comparing (37) and (41), we see that  $P_i$  has a cusp at  $t = t_{\text{H}}$ .

Next consider  $2t_{\text{H}} < t < 3t_{\text{H}}$ . In this case, there are four types of contributing terms: (i)  $m_j = 0$  for all  $j$ ; (ii)  $m_j = 1$  for one  $j$ , while 0 for all other  $j$ ; (iii)  $m_j = 2$  for one  $j$  while 0 for all other  $j$ ; (iv)  $m_j = 1$  for two  $j$ , while 0 for all other  $j$ . The contributions of (i) and (ii) to  $\langle b|S(t)|b \rangle$  are given by (37) and (41), respectively. The contribution of (iii) is given by

$$\begin{aligned}
&\sum_{n=0}^{\infty} \left( -\frac{g^2}{\hbar^2} \right)^n t_{\text{H}}^n \left( \frac{1}{2} \right)^{n-1} e^{i2\theta} \frac{(t-2t_{\text{H}})^n}{n!} n \\
&= -\gamma(t-2t_{\text{H}})e^{-\gamma(t-2t_{\text{H}})/2} e^{i2\theta}. \quad (42)
\end{aligned}$$

The contribution of (iv) is

$$\begin{aligned}
&\sum_{n=0}^{\infty} \left( -\frac{g^2}{\hbar^2} \right)^n t_{\text{H}}^n \left( \frac{1}{2} \right)^{n-2} e^{i2\theta} \frac{(t-2t_{\text{H}})^n}{n!} \frac{n(n-1)}{2} \\
&= \frac{1}{2} \gamma^2 (t-2t_{\text{H}})^2 e^{-\gamma(t-2t_{\text{H}})/2} e^{i2\theta}. \quad (43)
\end{aligned}$$

In total, for  $2t_{\text{H}} < t < 3t_{\text{H}}$ ,

$$\begin{aligned}
&S_{bb}(t) = e^{-\gamma t/2} - \gamma(t-t_{\text{H}})e^{-\gamma(t-t_{\text{H}})/2} e^{i\theta} \\
&+ \left[ \frac{1}{2} \gamma^2 (t-2t_{\text{H}})^2 - \gamma(t-2t_{\text{H}}) \right] e^{-\gamma(t-2t_{\text{H}})/2} e^{i2\theta}. \quad (44)
\end{aligned}$$

Again,  $P_i$  shows a cusp at  $t = 2t_{\text{H}}$ .

Similarly, we can determine the expression of  $S_{bb}(t)$  for later intervals. For those interested, closed explicit expressions of  $S_{bb}(t)$  for an arbitrary interval can be found in Refs. 16 and 29. But it is not hard to persuade oneself that in each interval, there will be extra terms contributing to  $C_n$  than in the preceding interval, hence extra terms in  $S_{bb}(t)$ . Therefore,  $P_i$  shows a cusp each time  $t$  is an integral multiplier of the Heisenberg time  $t_{\text{H}}$ . In Fig. 3, the trajectories of  $P_i$  are shown for the same four values of  $\alpha$  as in Fig. 2. We see that when all orders of terms in (3) are taken into account, many features in Fig. 2, which contains only the first order term, are still preserved. For example, the periodic singularities persist and the period is unchanged, moreover, there is still no




FIG. 3: (Color online) The survival probability  $P_i$  on the discrete level  $|b\rangle$  in the exact solution. It shows cusps periodically and the period is the Heisenberg time  $t_H = 2\pi\hbar/\Delta$ . In the first period,  $P_i(t) = \exp(-\gamma t)$ , with  $\gamma = 2\pi g^2/\Delta\hbar$ . In this period, it is independent of the value of  $\alpha$ . But afterwards, it is sensitive to the value of  $\alpha$ . The parameters are chosen such that  $g/\Delta = 0.15$ .

$\alpha$ -dependence in the first interval  $0 < t \leq t_H$ .

## V. CONCLUSIONS AND DISCUSSIONS

We consider the model (13) as of great pedagogical value. First, the model allows a lucid derivation of Fermi's golden rule. The calculation can be carried out straightforwardly without invoking any approximation or argument. It illustrates the two key features of Fermi's golden rule very well, namely, the transition rate is proportional to the density of states and the coupling

squared. Second, it demonstrates in a vivid way how the rule might break down beyond the Heisenberg time, a point less emphasized in quantum mechanics textbooks. One reason might be that many authors have in mind a true continuum, for which the Heisenberg time is infinite and thus irrelevant. However, a quasi-continuum with finite level spacings or finite Heisenberg times, is a reality in many systems. This is the case, for example, in a molecule,<sup>27</sup> in a multi-mode optical cavity,<sup>17,18</sup> or in a one-dimensional tight binding model,<sup>28</sup> where the piecewise linear behavior of the transition probability was observed.

Interestingly, the transition dynamics of the model can actually be solved exactly by collecting all orders of terms in the Dyson series, which in many quantum mechanics textbooks is introduced formally but not put into full use. The rigorous exponential decay in the first interval is rare instead of common. The periodic cusps imply that the kinks in the first order perturbation approximation is not mere artifact—they are modified but not smeared out completely in the exact solution. Recently, cusps (periodic or not) have been found also in the quench dynamics of the transverse Ising model,<sup>30</sup> the tight binding model,<sup>31</sup> a generalized XY spin chain,<sup>32</sup> and some non-integrable models.<sup>33</sup> The mechanism varies from case to case. In view of these new progresses, why and when can the time evolution of some quantity be singular is now a problem worth consideration.

## Acknowledgments

The author is grateful to Hua-Tong Yang for his helpful comments. This work is supported by the Fujian Provincial Science Foundation under grant number 2016J05004.

\* Electronic address: wdlang06@163.com

<sup>1</sup> P. A. M. Dirac, "The quantum theory of the emission and absorption of radiation," Proc. R. Soc. Lond. A **114**, 243-265 (1927).

<sup>2</sup> E. Fermi, *Nuclear Physics* (University of Chicago Press, Chicago, 1950), page 142.

<sup>3</sup> E. Fermi, "Versuch einer Theorie der beta-Strahlen. I," Z. Physik **88**, 161-177 (1934).

<sup>4</sup> F. L. Wilson, "Fermi's theory of Beta decay," Am. J. Phys. **36**, 1150-1160 (1968).

<sup>5</sup> P. A. M. Dirac, *The Principles of Quantum Mechanics*, 4th ed. (Oxford University Press, Oxford, 1958).

<sup>6</sup> A. Messiah, *Quantum Mechanics* (Dover Publications, New York, 2014).

<sup>7</sup> L. D. Landau and E. M. Lifshitz, *Quantum Mechanics*, 3rd ed. (Elsevier, Singapore, 1965).

<sup>8</sup> C. Cohen-Tannoudji, B. Diu, and F. Laloe, *Quantum Mechanics* (Wiley-VCH, New York, 1990).

<sup>9</sup> S. Weinberg, *Lectures on Quantum Mechanics* (Cambridge University Press, New York, 2013).

<sup>10</sup> J. J. Sakurai, *Modern Quantum Mechanics* (Addison-Wesley, New York, 1994).

<sup>11</sup> J. Y. Zeng, *Quantum Mechanics*, 3rd ed. (Science Press, Beijing, 2000).

<sup>12</sup> E. Merzbacher, *Quantum Mechanics*, 2nd ed. (John Wiley & Sons, London, 1970).

<sup>13</sup> L. E. Ballentine, *Quantum Mechanics—A Modern Development* (World Scientific, Singapore, 2000).

<sup>14</sup> G. Baym, *Lectures on Quantum Mechanics* (Westview Press, New York, 1990).

<sup>15</sup> F. John, *Partial Differential Equations* (Springer-Verlag, New York, 1982).

<sup>16</sup> G. C. Stey and R. W. Gibberd, "Decay of quantum states in some exactly soluble models," Physica (Amsterdam) **60**, 1-26 (1972).

<sup>17</sup> J. Parker and C. R. Stroud, Jr., "Transient theory of cavity-modified spontaneous emission," Phys. Rev. A **35**, 4226-4237 (1987).

<sup>18</sup> H. Giessen, J. D. Berger, G. Mohs, P. Meystre, and S. F. Yelin, "Cavity-modified spontaneous emission: From Rabi

- oscillations to exponential decay,” *Phys. Rev. A* **53**, 2816-2821 (1996).
- <sup>19</sup> M. Ligare and S. Becker, “Simple soluble models of quantum damping applied to cavity quantum electrodynamics,” *Am. J. Phys.* **63**, 788 (1995).
- <sup>20</sup> M. Ligare and R. Oliveri, “The calculated photon: Visualization of a quantum field,” *Am. J. Phys.* **70**, 58-66 (2002).
- <sup>21</sup> G. Mussardo, *Statistical Field Theory* (Oxford University Press, Oxford, 2010).
- <sup>22</sup> E. Hecht, *Optics*, 4th ed. (Addison-Wesley, New York, 2002).
- <sup>23</sup> It is actually the other way around, i.e., the Fourier transform of the window function is the sinc function.
- <sup>24</sup> L. Rade and B. Westergren, *Mathematics Handbook for Science and Engineering*, 5th ed. (Springer-Verlag, Berlin, 2004).
- <sup>25</sup> W. Rudin, *Real and Complex Analysis*, 3rd ed. (McGraw-Hill, New York, 1987).
- <sup>26</sup> A function  $f(z)$  defined on the complex plane is said to be of exponential type if there exist constants  $A$  and  $\alpha$  such that  $|f(z)| \leq Ae^{\alpha|z|}$  in the limit of  $|z| \rightarrow \infty$ . Letting  $\beta$  stand for the infimum of all such  $\alpha$ , one then says that the function  $f$  is of exponential type  $\beta$ .
- <sup>27</sup> E. Kyrölä, “Photoexcitation of a quasi-continuum: connections to few-level dynamics,” *J. Opt. Soc. Am. B* **1**, 737-741 (1984).
- <sup>28</sup> J. M. Zhang and M. Haque, “Nonsmooth and level-resolved dynamics illustrated with a periodically driven tight-binding model,” *ScienceOpen Research* 2014 (DOI: 10.14293/S2199-1006.1.SOR-PHYS.A2CEM4.v1).
- <sup>29</sup> R. Lefebvre and J. Savolainen, “Memory functions and recurrences in intramolecular processes,” *J. Chem. Phys.* **60**, 2509-2515 (1974).
- <sup>30</sup> M. Heyl, A. Polkovnikov, and S. Kehrein, “Dynamical quantum phase transitions in the transverse-field Ising model,” *Phys. Rev. Lett.* **110**, 135704 (2013).
- <sup>31</sup> J. M. Zhang and H. T. Yang, “Cusps in the quenched dynamics of a Bloch state,” arXiv:1601.03569.
- <sup>32</sup> U. Divakaran, S. Sharma, and A. Dutta, “Tuning the presence of dynamical phase transitions in a generalized  $XY$  spin chain,” arXiv:1601.04851.
- <sup>33</sup> C. Karrasch and D. Schuricht, “Dynamical phase transitions after quenches in nonintegrable models,” *Phys. Rev. B* **87**, 195104 (2013).