

Epstein Barr virus - The cause of multiple sclerosis

Katarina Barukčić¹, Ilija Barukčić²

¹ Medical University Sofia, Bulgaria

Email: Katarina.Barukcic@gmx.de

² Internist, Horandstrasse, DE-26441 Jever, Germany

Email: Barukcic@t-online.de

Submitted to viXra Saturday, May 7, 2016. Accepted; Published

Copyright © 2016 by Katarina Barukčić, Ilija Barukčić .

Abstract

Although many studies have found a kind of a relationship between an Epstein–Barr virus (EBV) and the development of multiple sclerosis (MS), a fundamental aspect of this relationship remain uncertain. What is the cause of multiple sclerosis (MS)? In this study, we re-analysed the data as published by Wandinger et al. and were able to establish the new insight that *without* an Epstein–Barr virus (EBV) infection *no* development of multiple sclerosis (MS) is possible. Furthermore, we determined a highly significant causal relationship between Epstein–Barr virus (EBV) and multiple sclerosis. Altogether, Epstein–Barr virus (EBV) is the cause of multiple sclerosis (p-value 0,0004251570).

Keywords

Epstein Barr virus; Multiple sclerosis

1. Introduction

Multiple sclerosis (MS) is an unpredictable disease of the central nervous system which disrupts the communication between the brain and other parts of the body. Multiple sclerosis (MS) can range from relatively benign to somewhat disabling and devastating symptoms. Some of today approved drugs to treat multiple sclerosis include Novantrone (mitoxantrone), teriflunomide, dimethyl fumarate, copolymer I (Copaxone) and forms of beta interferon. Steroids are used to reduce the duration and severity of attacks in some patients suffering from Multiple sclerosis. Exercise and physical therapy can help to preserve remaining function. Various aids such as foot braces, canes, and walkers are of use to help patients to remain independent and mobile. Thus far, there is as yet no cure for multiple sclerosis while millions of people are suffering from this disease.

Epstein-Barr virus (EBV), a herpes virus, is a primary cause of infectious mononucleosis (IM) and associated with several malignancies including such as Hodgkin lymphoma, non-Hodgkin lymphoma, Burkitt lymphoma

and other. Epidemiological, molecular virology and other [1]-[6] studies have been able to establish EBV as a risk factor for the development of multiple sclerosis (MS) and provided some evidence that the pathogenesis of multiple sclerosis might involve a response to an EBV infection. Still, the cause of Multiple sclerosis is not identified.

2. Material and methods

2.1. Definitions

Definition. Bernoulli random variable

Let $t=+1, \dots, +N$ denote an individual Bernoulli trial each with *constant* success probability p . Let N denote the number of independent Bernoulli trials (the size of a random sample or of the population).

Definition. The 2x2 table

Let A_t denote a Bernoulli distributed random variable. Let $p(A_t)$ denote the probability of A_t . Let B_t denote a Bernoulli distributed random variable. Let $p(B_t)$ denote the probability of B_t . Let $p(a_t)=p(A_t \cap B_t)$ denote joint distribution of A_t and B_t . Let $p(b_t)=p(A_t \cap \underline{B}_t)$ denote joint distribution of A_t and \underline{B}_t . Let $p(c_t)=p(\underline{A}_t \cap B_t)$ denote joint distribution of \underline{A}_t and B_t . Let $p(d_t)=p(\underline{A}_t \cap \underline{B}_t)$ denote joint distribution of \underline{A}_t and \underline{B}_t . In general, it is $p(a_t)+p(b_t)+p(c_t)+p(d_t)=1$. Thus far, the relationships before are expressed in the 2×2 table (**Table**).

Table .The 2x2 Table

		Conditioned B_t		
		Yes	No	
Condition A_t	Yes	$p(a_t)$	$p(b_t)$	$p(a_t)+p(b_t)=p(A_t)$
	No	$p(c_t)$	$p(d_t)$	$p(c_t)+p(d_t)=p(\underline{A}_t)$
		$p(a_t)+p(c_t)=p(B_t)$	$p(b_t)+p(d_t)=p(\underline{B}_t)$	1

Thus far, let $A=N \times p(A_t)$ denote the expectation value. Let $\underline{A}=N-A=N \times (1-p(A_t))$ denote the expectation value. Let $B=N \times p(B_t)$ denote the expectation value. Let $\underline{B}=N-B=N \times (1-p(B_t))$ denote the expectation value. Let $a=N \times p(a_t)=N \times p(A_t \cap B_t)$ denote the expectation value. Let $b=N \times p(b_t)=N \times p(A_t \cap \underline{B}_t)$ denote the expectation value. Let $c=N \times p(c_t)=N \times p(\underline{A}_t \cap B_t)$ denote the expectation value. Let $d=N \times p(d_t)=N \times p(\underline{A}_t \cap \underline{B}_t)$ denote the expectation value. Let $N=a+b+c+d=N \times (p(a_t)+p(b_t)+p(c_t)+p(d_t))$ denote the size of the sample or the size of the population. Let $A=a+b$ denote the expectation value of the condition (i. e. a risk factor, the verum population, the exposed group). Let $\underline{A}=c+d$ denote the expectation value of the non-condition (i. e. the non-exposed group, the control population). Let $B=a+c$ denote the expectation value of the conditioned. Let $\underline{B}=b+d$ denote the expectation value of the not conditioned. Thus far, the relationships before are expressed in the 2×2 table (**Table**).

Table .The 2x2 Table

		Conditioned B		
		Yes	No	
Condition A	Yes	a	b	$a+b=A$
	No	c	d	$c+d=\underline{A}$
		$a+c=B$	$b+d=\underline{B}$	N

Definition. Risk Ratio or Relative Risk

Various quantitative techniques are used in Biostatistics to the describe and evaluate relationships among biologic and medical phenomena. Relative risk, defined by Fischer [7] as ψ , is an important [8], [9] statistical

*Special description of the title. (dispensable)

method used in in epidemiologic studies and clinical trials. Let $RR(A,B)$ denote the relative risk. Based on the 2 by 2 table above, the relative risk $RR(A,B)$ is defined as

$$RR(A,B) \equiv \frac{a/(a+b)}{c/(c+d)} \tag{1}$$

In epidemiology and statistics, relative risk (RR) is the ratio of *the probability of an event a occurring under conditions of being exposed to (a+b)*, the non-exposed to *the probability of c occurring under conditions of being exposed to (c+d)*, the non-exposed group. The relative risk (RR) is a widely used *measure of association* in epidemiology. A risk ratio $RR(A,B) < 1$ suggest that an exposure can be considered as being associated with a reduction in risk. A risk ratio $RR(A,B) > 1$ suggest that an exposure can be considered as being associated with an increase in risk.

Conditions

The following relationships are taken with friendly permission by Ilija Barukčić [10].

Definition. Conditio sine qua non relationship

Let $p(A_t \leftarrow B_t)$ denote [10] the extent to which a condition A is a *conditio sine qua non* of the conditioned B. The *conditio sine qua non* relationship is calculated as

$$p(A_t \leftarrow B_t) \equiv p(a_t) + p(b_t) + p(d_t) = \frac{N}{N} \times (p(a_t) + p(b_t) + p(d_t)) = \frac{a+b+d}{N} \equiv \frac{A+d}{N} \equiv \frac{a+B}{N} \tag{2}$$

The relationship before is expressed in the following 2×2 (Table).

Table . Conditio sine qua non.				
		Conditioned B		
		Yes	No	
Condition A	Yes	a	b	a+b= A
	No	c=0	d	c+d = <u>A</u>
		a+c = B	b+d = <u>B</u>	N

Definition. Anti conditio sine qua non relationship

Let $p(A_t > B_t)$ denote [10] the extent to which a condition A is not a *conditio sine qua non* of the conditioned B. The *anti conditio sine qua non* relationship is calculated as

$$p(A_t > B_t) \equiv p(c_t) = \frac{N}{N} (p(c_t)) \equiv \frac{N-a-b-d}{N} \equiv \frac{c}{N} \equiv 1 - p(A_t \leftarrow B_t) \tag{3}$$

The relationship before is expressed in the following 2x2 table (Table).

Table . Anti conditio sine qua non.				
		Conditioned B		
		Yes	No	
Condition A	Yes	a=0	b=0	a+b= A
	No	c	d=0	c+d = <u>A</u>
		a+c = B	b+d = <u>B</u>	N

Definition. Conditio per quam relationship

Let $p(A_t \rightarrow B_t)$ denote [10] the extent to which a condition A is a *conditio per quam* of the conditioned B. The *conditio per quam* is calculated as

$$p(A_t \rightarrow B_t) \equiv p(a_t) + p(c_t) + p(d_t) = \frac{N}{N} \times (p(a_t) + p(c_t) + p(d_t)) = \frac{a+c+d}{N} \equiv \frac{B+d}{N} \equiv \frac{a+A}{N} \quad (4)$$

The relationship before is expressed in the following 2x2 table (**Table**).

Table . Conditio per quam.

		Conditioned B		
		Yes	No	
Condition A	Yes	a	b=0	a+b= A
	No	c	d	c+d = <u>A</u>
		a+c = B	b+d = <u>B</u>	N

Definition. Anti conditio per quam relationship

Let $p(A_t > -B_t)$ denote [10] the extent to which a condition A is not a *conditio per quam* of the conditioned B. The anti *conditio per quam* relationship is calculated as

$$p(A_t > -B_t) \equiv p(b_t) = \frac{N}{N} \times (p(b_t)) \equiv \frac{b}{N} \equiv \frac{N-a-c-d}{N} \equiv 1 - p(A_t \rightarrow B_t) \quad (5)$$

The relationship before is expressed in the following 2×2 table (**Table**).

Table. Anti conditio per quam.

		Conditioned B		
		Yes	No	
Condition A	Yes	a=0	b	a+b= A
	No	c=0	d=0	c+d = <u>A</u>
		a+c = B	b+d = <u>B</u>	N

Definition. Conjunction A and B relationship

Let $p(A_t \cap B_t)$ denote [10] the extent to which a condition A is *conjugated* with the conditioned B. The conjunction is calculated as conjunction of the two events

$$p(A_t \cap B_t) \equiv p(a_t) = \frac{N}{N} \times (p(a_t)) = \frac{a}{N} \equiv \frac{N-b-c-d}{N} \equiv 1 - p(A_t \sqsupset B_t) \quad (6)$$

The relationship before is expressed in the following 2x2 table (**Table**).

Table . Conjugation

		Conditioned B		
		Yes	No	
Condition A	Yes	a	b=0	a+b= A
	No	c=0	d=0	c+d = <u>A</u>
		a+c = B	b+d = <u>B</u>	N

Definition. Exclusion relationship

Let $p(A_t \sqcap B_t)$ denote [10] the extent to which a condition A *excludes* the conditioned B *and vice versa*. The exclusion relationship (Sheffer stroke) named after Henry M. Sheffer is written as a vertical bar or an upwards arrow and calculated as

$$p(A_t \sqcap B_t) \equiv p(A_t | B_t) \equiv p(A_t \uparrow B_t) \equiv p(b_t) + p(c_t) + p(d_t) = \frac{N}{N} \times (p(b_t) + p(c_t) + p(d_t)) \quad (7)$$

or

$$p(A_t \sqcap B_t) \equiv p(A_t | B_t) \equiv p(A_t \uparrow B_t) \equiv \frac{b+c+d}{N} \equiv \frac{N-a}{N} \equiv 1 - p(A_t \cap B_t) \quad (8)$$

The relationship before is expressed in the following 2 x 2 table (**Table**).

Table.Exclusion.

		Conditioned B		
		Yes	No	
Condition A	Yes	a=0	b	a+b= A
	No	c	d	c+d = <u>A</u>
		a+c = B	b+d = <u>B</u>	N

Definition. Disjunction A or B relationship

Let $p(A_t \cup B_t)$ denote [10] the extent to which the condition A or the conditioned B are given. The inclusive disjunction also known as *alternation* is calculated as

$$p(A_t \cup B_t) \equiv p(a_t) + p(b_t) + p(c_t) = \frac{N}{N} \times (p(a_t) + p(b_t) + p(c_t)) \equiv 1 - p(A_t \sqcap B_t) \quad (9)$$

or

$$p(A_t \cup B_t) \equiv p(A_t) + p(B_t) - p(a_t) = \frac{a+b+c}{N} \equiv \frac{A+B-a}{N} \equiv 1 - p(A_t \sqcap B_t) \quad (10)$$

The relationship before is expressed in the following 2x2 table (**Table**).

Table . Disjunction

		Conditioned B		
		Yes	No	
Condition A	Yes	a	b	a+b= A
	No	c	d=0	c+d = <u>A</u>
		a+c = B	b+d = <u>B</u>	N

Definition. Neither A nor B relationship

Let $p(A_t \sqcup B_t)$ denote [10] the extent to which neither a condition A nor the conditioned B is given. The *neither A nor B* relationship was introduced by Charles Sanders Peirce and is known as Peirce's arrow and can be calculated as

$$p(A_t \sqcup B_t) \equiv p(A_t \downarrow B_t) \equiv p(d_t) = \frac{N}{N} \times (p(d_t)) = \frac{d}{N} \equiv \frac{N - a - b - c}{N} \equiv 1 - p(A_t \cap B_t) \quad (11)$$

The relationship before is expressed in the following 2 x 2 table (**Table**).

Table.Exclusion.

		Conditioned B		
		Yes	No	
Condition A	Yes	a=0	b=0	a+b= A
	No	c=0	d	c+d = <u>A</u>
		a+c = B	b+d = <u>B</u>	N

Definition. Equivalence of A and B relationship

Let $p(A_t \Leftrightarrow B_t)$ denote [10] the extent to which a condition A or the conditioned B are equivalent. The inclusive disjunction also known as alternation is calculated as

$$p(A_t \Leftrightarrow B_t) \equiv p(a_t) + p(d_t) = \frac{N}{N} \times (p(a_t) + p(d_t)) \equiv 1 - p(A_t \succ \Leftarrow B_t) \quad (12)$$

The relationship before is expressed in the following 2x2 table (**Table**).

Table . Disjunction

		Conditioned B		
		Yes	No	
Condition A	Yes	a	b=0	a+b= A
	No	c=0	d	c+d = <u>A</u>
		a+c = B	b+d = <u>B</u>	N

Definition. Either A or B relationship

Let $p(A_t \succ \Leftarrow B_t)$ denote [10] the extent to which *either* a condition A *or* the conditioned B is given. The *either A or B* relationship can be calculated as

$$p(A_t \succ \Leftarrow B_t) \equiv p(b_t) + p(c_t) = \frac{N}{N} \times (p(b_t) + p(c_t)) = \frac{b+c}{N} \equiv \frac{N - a - d}{N} \equiv 1 - p(A_t \Leftrightarrow B_t) \quad (13)$$

The relationship before is expressed in the following 2×2 table (**Table**).

Table.Exclusion.

		Conditioned B		
		Yes	No	
Condition A	Yes	a=0	b	a+b= A
	No	c	d=0	c+d = <u>A</u>
		a+c = B	b+d = <u>B</u>	N

2.2. Material

Patients and samples

Data and material for this re-analysis were published by Wandinger [11] et al., a study specifically designed to investigate the relation between multiple sclerosis (MS) and viral infections. Wandinger [11] et al. examined sera from a large cohort of 163 healthy control subjects (control group) and 108 patients with a diagnosis of clinically definite multiple sclerosis for the presence of human herpesviruses type 1 (HSV-1), HSV-2, cytomegalovirus (CMV) and EBV by the presence of IgG antibodies. In addition, other investigations (i. e. the detection of EBV DNA in all serum samples) were performed. Some of the data of Wandinger [11] et al. data about the prevalence of IgG antibodies in serum samples from multiple sclerosis (MS) patients and healthy control subjects are summarized in the table shown below (**Table**).

Table . Prevalence of IgG antibodies in MS patients and healthy control subjects xclusion.

Parameter	Multiple sclerosis (MS)	Healthy control subjects
anti-EBNA-1 IgG	108	147
Sample size	108	163

The data of the prevalence of IgG antibodies in serum samples from multiple sclerosis (MS) patients and healthy control subjects are viewed in the following 2×2 table (**Table**).

Table .EBV and multiple sclerosis (MS).

		Multiple sclerosis		
		yes	no	
EBV anti-EBNA-1 IgG	Yes	108	147	255
	No	0	16	16
		108	163	271

2.2. Methods

The chi-squared distribution

The properties of the chi-squared distribution were first investigated by Karl Pearson [12] in 1900. The chi-squared distribution is a widely used probability distributions in hypothesis testing, inferential statistics or in construction of confidence intervals.

Table . Chi square distribution for degree of freedom d. f.= 1.

Critical values of chi-square distribution		
p-Value	One sided	Two sided
0,1000000000	1,642374415	2,705543454
0,0500000000	2,705543454	3,841458821
0,0400000000	3,06490172	4,217884588
0,0300000000	3,537384596	4,709292247
0,0200000000	4,217884588	5,411894431
0,0100000000	5,411894431	6,634896601
0,0010000000	9,549535706	10,82756617
0,0001000000	13,83108362	15,13670523
0,0000100000	18,18929348	19,51142096
0,0000010000	22,59504266	23,92812698
0,0000001000	27,03311129	28,37398736
0,0000000100	31,49455797	32,84125335
0,0000000010	35,97368894	37,32489311
0,0000000001	40,46665791	41,8214562

In last consequence, the Chi Square with one degree of freedom is nothing but the distribution of a single normal deviate squared.

The binomial proportion confidence interval

The statistical significance of deviations from a theoretically expected distribution of observations can be tested by a *binomial test*. For large samples, the binomial distribution is well approximated by convenient Pearson's chi-squared test. The above relationships are grounded on the assumption, that the number of successes X out of a sample of n observations is equal to $X=N$. In general, let $df1_l$ denote the degrees of freedom 1 of the f -distribution for the lower confidence bound. Thus far, it is $df1_{lower} = 2(N-X+1)$. Under conditions where $N=X$ the proportion of success is $p(X/N)=1$, the is then $df1_{lower} = 2$. Let $df2_{lower}$ denote the degrees of freedom 2 of the f -distribution for the lower confidence bound. In particular, we obtain $df2_{lower} = 2 \times X$. Under conditions where $N=X$ the proportion of success is $p(X/N)=1$ and $df2_{lower} = 2 \times N$. The exact *one-sided* lower confidence interval with confidence level $1 - \alpha$ for the proportion of successes $p(X/N)=1$ can be calculated [13] as

$$P_{Lower} = \frac{N}{N + F_{(df1_{lower}, df2_{lower}, \alpha)}} \quad (14)$$

Example.

The F-Value for $X=N=271$ ($\alpha = 0.05$) is $F_{df1=2, df2=542, \alpha=0.05} = 3.01235141$. The exact *one-sided* lower confidence bound for the proportion of successes $p(X/N)=1$ can be calculated as

$$P_{Lower} = \frac{N}{N + F_{(df1_{lower}, df2_{lower}, \alpha)}} = \frac{271}{271 + 3,01235141} = 0,989006512 \quad (15)$$

In other words, we assume that the p in the population is greater or equal to $0,989006512$. Furthermore, the *one-sided* lower confidence interval with confidence level $1 - \alpha$ for the proportion of successes $p(X/N)=1$, reflects a significance level of i. e. $\alpha = 0.05$, and can be calculated for $N > 50$ approximately [13] as

$$P_{Lower} \approx 1 - \frac{3}{N} \quad (16)$$

A $100 \times (1 - \alpha)\%$ confidence interval consists of all those values $p(X/N)$ for which a test of the hypothesis

$p(X/N) = 1$ is not rejected at a significance level of $100 \times (\alpha)\%$.

Causal relationship k

The mathematical formula of the causal relationship k was used to determine the cause-effect relationship between Epstein-Barr virus (EBV) infections and multiple sclerosis (MS). According to Barukčić [14], the causal relationship k is calculated as

$$k(A_t, B_t) \equiv \frac{p(a_t) - p(A_t) \times p(B_t)}{\sqrt[2]{p(A_t) \times (1 - p(A_t)) \times p(B_t) \times (1 - p(B_t))}} \equiv \frac{(N \times a) - (A \times B)}{\sqrt[2]{A \times \underline{A} \times B \times \underline{B}}} \quad (17)$$

The relationship before is expressed in the following 2×2 table (Table).

		Effect B		
		Yes	No	
Cause A	Yes	a	b	a+b= A
	No	c	d	c+d= A
		a+c = B	b+d = B	N

Pearson's chi-squared test X^2

$$\chi^2 \equiv \frac{N \times ((a \times d) - (b \times c)) \times ((a \times d) - (b \times c))}{(a + b) \times (c + d) \times (a + c) \times (b + d)} \quad (18)$$

is used to evaluate how likely it is the observed causal relationship k arose by chance. The 2×2 contingency table is dichotomous while the statistical X^2 distribution is continuous. Thus far, Pearson's chi-square test tends to make results larger than they should be and is biased upwards on this account. This upwards bias of Pearson's chi-square test can be corrected by using Yates correction.

Scholium.

As a response to Yules association of two attributes Karl Pearson introduced *the mean square contingency* into statistics as

$$\phi^2 \equiv \frac{((a \times d) - (b \times c)) \times ((a \times d) - (b \times c))}{(a + b) \times (c + d) \times (a + c) \times (b + d)} \quad (19)$$

Still, Pearson failed to derive a mathematical formula of the causal relationship k and much more than this. Pearson himself exterminated any kind causation from statistics ultimately. Following Pearson, "We are now in a position, I think, to appreciate the scientific value of the word cause. Scientifically, cause ... is meaningless ..." [14]. According to Pearson, the words cause and effect belong strictly to the sphere of sense-impressions. Thus far, "there is ... no true cause and effect" [14]. The reader can hardly fail to have been impressed that Pearson himself denies any kind of causality. In the first place, there is no causation at all. "No phenomena are causal" [14]. Finally, "The wider view of the universe sees all phenomena as correlated, but not causally related." [14]. Consequently, "... there is association but not causation." [14]. We have now reached some very important conclusions about Pearson's account for causality. Due to Pearson, there is no causation at all. Pearson published a lot of mathematical formulas. But neither Pearson's correlation coefficient nor his mean square contingency can be regarded as the mathematical formula of the causal relationship k. In particular, Pearson failed to derive a mathematical formula of the causal relationship and to provide a self-consistent proof of a mathematical formula of the causal relationship.

Statistical analysis.

Data were analyzed using Microsoft Excel version 14.0.7166.5000 (32-Bit) software (Microsoft GmbH, Munich, Germany). The mathematical formula of the causal relationship k and the chi-square distribution were applied to determine the significance of causal relationship between EBV and multiple sclerosis (MS). A p value of < 0.05 was considered significant.

3. Results

3.1. Clinical characteristics

Wandinger [11] et al. examined 108 MS patients from the Department of Neurology (University of Lübeck School of Medicine). All patients were examined independently by two neurologists and had a diagnosis of clinically definite MS. Kurtzke's functional systems and Expanded Disability Status Scale (EDSS) were used to grade the physical disability.

3.2. EBV seropositivity

The viral status was classified by following serologic definitions. Wandinger [11] et al. defined *primary EBV infection* by positivity of anti-EA-IgG and/or anti-EA-IgM in the absence of anti-EBNA-1 antibodies. A *latent or past EBV infection* was defined by positivity of anti-EBNA-1 antibodies. A *reactivation of a latent EBV infection* was defined by EBNA-1-IgG-positive individuals by additional positive anti-EA-IgG and anti-EA-IgM or additional high anti-EA-IgM. The marker for *latent EBV infection* was defined by an anti-EBV nuclear antigen type 1 (anti-EBNA-1) immunoglobulin (Ig) G antibodies.

3.3. Epstein Barr virus (EBV) is a *conditio sine qua non* of multiple sclerosis (MS)

A hypothesis test is used to distinguish between the null hypothesis and the alternative hypothesis.

Theorem 1.

Null hypothesis: EBV is a condition sine qua non of multiple sclerosis (MS)

Alternative hypothesis: EBV is not a condition sine qua non of multiple sclerosis (MS).

Significance level (α) below which the null hypothesis will be rejected: 0.05

Proof by a statistical hypothesis test.

The data of the prevalence of IgG antibodies in serum samples from multiple sclerosis (MS) patients and healthy control subjects are viewed in the following 2×2 table (**Table**).

Table . Without EBV no multiple sclerosis (MS).

		Multiple sclerosis		
		yes	no	
EBV anti-EBNA-1 IgG	Yes	108	147	255
	No	0	16	16
		108	163	271

The proportion of successes $p(A_t < B_t)$ of the *conditio sine qua non relationship* in the sample or the test statistic can be calculated defined before as

$$p(A_t \leftarrow B_t) \equiv \frac{a+b+d}{N} \equiv \frac{A+d}{N} \equiv \frac{a+B}{N} \equiv \frac{108+147+16}{271} \equiv \frac{247}{271} \equiv 1 \quad (20)$$

The critical value p_{lower} calculated as

$$p_{\text{Lower}} \approx 1 - \frac{3}{N} = 1 - \frac{3}{247} = 0,988929889 \quad (21)$$

The critical value $p_{\text{lower}} = 0,989006512$ and is less than the proportion of successes $p(A_t \leftarrow B_t) = 1$ as obtained from the observations (the significance level $\alpha = 0.05$).

Conclusio.

We cannot reject the null hypothesis in favor of the alternative hypotheses.

Epstein Barr virus (EBV) is a *conditio sine qua non* of multiple sclerosis (MS).

In other words, *without* an infection with Epstein Barr virus (EBV) *no* development of multiple sclerosis (MS).

Quod erat demonstrandum.

3.4. Epstein Barr virus (EBV) is the cause of multiple sclerosis (MS)

Theorem 2.

Conditions.

Alpha level = 5%.

The two tailed critical Chi square value (d.f.=1) for alpha level 5% is 3,841458821.

Claims.

Null hypothesis (H_0): $k=0$. (No causal relationship).

There is no causal relationship between Epstein Barr virus (EBV) and multiple sclerosis (MS)

Alternative hypothesis (H_1): $k \neq 0$. (Causal relationship).

There is a significant causal relationship between Epstein Barr virus (EBV) and multiple sclerosis (MS)

Proof by two sided hypotheses test.

Based on the data of Wandinger et al.

Table .EBV and multiple sclerosis (MS).

		Multiple sclerosis		
		yes	no	
EBV anti-EBNA-1 IgG	Yes	108	147	255
	No	0	16	16
		108	163	271

we compute the causal relationship $k(\text{EBV,MS})_{\text{Obtained}}$ (our test statistic) as

$$k(\text{EBV,MS})_{\text{Obtained}} \equiv \frac{(N \times a) - (A \times B)}{\sqrt{A \times \underline{A} \times B \times \underline{B}}} = \frac{271 \times 108 - 255 \times 108}{\sqrt{108 \times 163 \times 255 \times 16}} = 0,2038956576 \quad (22)$$

Following Barukčić, the test statistics obtained is equivalent with a X^2 value of

$$\chi^2 \equiv k(\text{EBV, MS})_{\text{Obtained}} \times k(\text{EBV, MS})_{\text{Obtained}} \times N = 271 \times (0,2038956576)^2 = 11,2664020209 \quad (23)$$

A two tail Chi square of 11,2664020209 is equivalent to a p-value of 0,0004251570.

Conclusio.

The value of the test statistic (k obtained or Chi square calculated) is 11,2664020209 and exceeds the critical Chi square value of 3,841458821. Consequently, we reject the null hypothesis (H_0) and accept the alternative hypothesis (H_A).

There is a highly significant causal relationship between Epstein Barr virus (EBV) and multiple sclerosis ($k=+0,2038956576$, p-value 0,0004251570).

Quod erat demonstrandum.

4. Discussion

Today, the etiology of multiple sclerosis (MS) is largely unknown but multiple sclerosis (MS) is rare among individuals without serum EBV antibodies. Thus far, there is an accumulating literature for a role of Epstein-Barr virus (EBV) infections in the pathogenesis of multiple sclerosis (MS). Especially, several epidemiological studies suggested an association between infection with Epstein-Barr virus (EBV) and the occurrence of multiple sclerosis (MS) disease. In particular, a recent large prospective epidemiological study showed a relationship between an increase of serum antibody titres against EBV before onset of MS. Acherio et al. [16] conducted a prospective, nested case-control study of 62 439 women participating in the Nurses' Health Study to determine whether elevation in serum antibody titers to EBV precede the occurrence of multiple sclerosis (MS). Acherio et al. concluded that EBV is associated with the etiology of multiple sclerosis. Recently, Levin et al. [17] conducted a study among more than 3 million US military personnel and found a relationship between EBV infection and development of MS. Apart from these and other studies aiming at the aetiology of multiple sclerosis (MS), the cause of multiple sclerosis (MS) was still not identified.

We conducted a re-analysis of the study of Wandinger et al. to investigate the role between EBV infection and MS disease. Using the material obtained by the study of Wandinger et al., we questioned whether Epstein-Barr virus (EBV) is the cause or a cause of multiple sclerosis (MS). The study of Wandinger [11] et al. was properly constructed. In accordance with previous studies, Wandinger [11] et al. found an unexpectedly high seropositivity rate in MS patients for EBV compared with control subjects. Wandinger [11] et al. observed an association of the EBV with MS but failed to detect the true meaning of Epstein-Barr virus.

In addition, our study confirms *a conditio sine qua non relationship* between EBV infection and multiple sclerosis (MS). In other words, *without* an infection with Epstein-Barr virus (EBV) *no* development of multiple sclerosis (MS) (significance level alpha = 0.05). We observed a highly significant causal relationship between Epstein-Barr virus (EBV) and multiple sclerosis ($k=+0,203895658$, p value = 0,000425157). A particular aspect of our study is the identification of Epstein-Barr virus (EBV) as the cause of multiple sclerosis. Since *without* an infection by Epstein-Barr virus (EBV) *no* multiple sclerosis develops and due to the fact that there is a highly significant causal relationship between Epstein-Barr virus (EBV) and multiple sclerosis, we are allowed to deduce that Epstein-Barr virus (EBV) is not only a cause but the cause of multiple sclerosis (MS).

5. Conclusions

A particular aspect of our study is the identification of Epstein-Barr virus (EBV) as the cause of multiple sclerosis (MS). Finally, the cause of multiple sclerosis is identified.

References

- [1] Bray, P. F., Bloomer, L. C., Salmon, V. C., Bagley, M. H. and Larsen, P.D. (1983) Epstein-Barr virus infection and antibody synthesis in patients with multiple sclerosis. *Arch Neurol*, **40**, 406-408. <http://dx.doi.org/10.1001/archneur.1983.04050070036006>
- [2] Sumaya C. V., Myers L. W., Ellison G. W., Ench Y. (1985). Increased prevalence and titer of Epstein-Barr virus antibodies in patients with multiple sclerosis. *Annals of Neurology*, **17**, 371–377. <http://dx.doi.org/10.1002/ana.410170412>
- [3] Larsen, P. D., Bloomer, L. C., Bray, P. F. (1985) Epstein-Barr nuclear antigen and viral capsid antigen antibody titers in multiple sclerosis. *Neurology*, **35**, 435–438. <http://dx.doi.org/10.1212/WNL.35.3.435>
- [4] Operskalski, E. A., Visscher, B. R., Malmgren, R. M., Detels, R. (1989) A case-control study of multiple sclerosis. *Neurology*, **39**, 825–829. <http://dx.doi.org/10.1212/WNL.39.6.825>
- [5] Ascherio, A. and Munch, M. (2000) Epstein-Barr virus and multiple sclerosis. *Epidemiology*, **11**, 220–224.
- [6] Hernan, M. A., Zhang, S. M., Lipworth, L., Olek, M. J., Ascherio, A. (2001) Multiple sclerosis and age at infection with common viruses. *Epidemiology*, **12**, 301–306.
- [7] Fisher, R. A. (1935). The Logic of Inductive Inference. *Journal of the Royal Statistical Society*, **98**, 39-82. <http://dx.doi.org/10.2307/2342435>. *Ibid.*, p. 50.
- [8] Gart, J. G. (1962) Approximate Confidence Limits for the Relative Risk. *Journal of the Royal Statistical Society. Series B (Methodological)*, **24**, 454-463.
- [9] Gart, J. J. (1962) On the combination of relative risks. *Biometrics*, **18**, 601.
- [10] Barukčić, I. (1989) *Causality I. A theory of energy, time and space*. 5th Edition, 19th Revision 2011, Lulu: Morrisville, North Carolina. pp. 648.
- [11] Wandinger, K., Jabs, W., Siekhaus, A., Bubel, S., Trillenber, P., Wagner, H., Wessel, K., Kirchner, H., Hennig, H. (2000) Association between clinical disease activity and Epstein-Barr virus reactivation in MS. *Neurology*, **55**, 178-184. <http://dx.doi.org/10.1212/WNL.55.2.178>
- [12] Pearson, Karl (1900). On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. *Philosophical Magazine Series 5*, **50**, 157-175. <http://dx.doi.org/10.1080/14786440009463897>
- [13] Sachs, L. (1992). *Angewandte Statistik: Anwendung statistischer Methoden*. Siebte, völlig neue bearbeitete Auflage, Springer: Berlin, p. 439
- [14] Barukčić, I. (2016). The Mathematical Formula of the Causal Relationship k. *International Journal of Applied Physics and Mathematics*, **6**, 45-65. <http://dx.doi.org/10.17706/ijapm.2016.6.2.45-65>
- [15] Pearson, K. (1904). *Mathematical contributions to the theory of evolution. XIII. On the theory of contingency and its relation to association and normal correlation*. Dulau and Co.: London, pp. 426.
- [16] Ascherio, A., Munger, K. L., Lennette, E. T., Spiegelman, D., Hernán, M. A., Olek, M. J., Hankinson, S. E., Hunter, D. J.. (2001). Epstein-Barr virus antibodies and risk of multiple sclerosis: a prospective study. *Journal of the American Medical Association*, **286**, 3083-3088. <http://dx.doi.org/jama.286.24.3083>
- [17] Levin, L. I., Munger, K. L., Rubertone, M. V., Peck, C. A., Lennette, E. T., Spiegelman, D., Ascherio, A. (2003). Multiple Sclerosis and Epstein-Barr Virus. *Journal of the American Medical Association*, **289**, 1533-1536. <http://dx.doi.org/10.1001/jama.289.12.1533>