

Dynamic Universe Model explains the Variations of Gravitational Deflection Observations of Very-Long-Baseline Interferometry

Abstract

A new type of tensor mathematical formation, using 21000 linear equations for 133 bodies, Dynamic Universe Model is a singularity free N-body problem solution, based on Newtonian physics. It is different from Newtonian physics, as well as general relativity as they use differential and integral equations. It uses linear tensors without differential and integral equations, hence it gives unique solution. This approach saves many unsolved problems in present day physics. Its prediction in 2005 that there is no dark matter came true experimentally in Nov, 2013.

Here, using the same math setup but with different initial set of numerical values, (viz., 3d xyz initial positions, velocities of all the solar system, nearby stars, Galaxy center, nearby Galaxies etc.), this paper explains the wide range of values of gravitational deflection (bending) results of Very Long Baseline Interferometry (VLBI) observations. The observed VLBI results are in much higher range even after accounting the standard errors. These higher ranges in VLBI results are beyond explainable values of general relativistic predictions as well as with parameterized post-Newtonian (PPN) formalism. Dynamic Universe model's built in potential of considering simultaneous and dynamic gravitational effect of Sun, planets, local stars etc., makes these discrepancies comprehensible. For doing so, the capabilities of Dynamic Universe Model are extended into micro world i.e., the world of masses of light photons and radio wavelength photons, Neutrinos electrons and protons etc., (This definition is some thing like a subset of quantum physics, we will use this name further in this paper) for the extended by extending the original mathematical formulation for macro world viz., planets, stars, Galaxies etc. Here it is shown that, Dynamic Universe Model explains reasons for the variations in real world VLBI deflection (bending) observations.

Keywords: Dynamic universe Model; SITA programming; Micro particle world; VLBI Very Long Baseline Interferometry; Gravitational deflection (bending): PPN formalism, Tensors, singularity-free cosmology, Dark matter.

1. Introduction

1.1. Gravitational deflection (bending) Observations of Very Long Baseline Interferometry (VLBI)

In Very-long-baseline interferometry (VLBI), a type of astronomical interferometry in astronomy, a signal from an astronomical radio source, such as a quasar or a manmade satellite, is collected at multiple radio telescopes on Earth and analyzed for delays. From early 1960s, VLBI researchers / Physicists are calculating the gravitational deflection (bending) of radio waves near huge masses like Sun or Jupiter. These VLBI techniques are used all over the world, also in the field of Radio astronomical observations of quasars, Galaxies etc. There is a large variation in the observational Gravitational deflection (bending) results of VLBI. The statistical variations in their observed results in comparison with General Relativistic predictions are noticeable. This variation is clearly visible when the solar gravitational bending / deflection angle is plotted against Solar Elongation angle. Additionally many people have observed variations; some plotted these with respect to season, frequency etc.

Why is there such variation amongst different measurements? Basically what the scientific community is calculating is the gravitational influence of Sun only on the radio waves / light. As observed by physicists Shapiro et al and Heinkelmann et al etc., there are other influencing factors other than Sun and other planets are also to be accounted for. About these authors and many more will be discussed later in this paper. Being a singularity free N-body problem solution Dynamic Universe Model offers an answer for the above problem, as it can consider mutual gravitational effect of simultaneously and dynamically changing planets, stars, Milkyway center, other parts of Milkyway and other Galaxies etc.

Just considering the Sun alone is not correct. For example anybody may think when he is sitting on earth; he can neglect the Gravitational forces of Sun and Moon on him because of the strong gravitational force of earth here. But it is not true. He can see the high tides on seas and oceans on earth which are visual indication of gravitational effect of Sun and Moon on Earth. For solving such problems, Dynamic Universe Model can be used by solving a singularity free 3- body or 4 body problem solution.

1.2. What different authors say about VLBI (some main points)?

Shapiro et al (2004) gave one such typical large compilation of data. They gave a large number of observations, which were made all over the world to the tune of 2 million observational experiments in the line of VLBI by many Physicists from 1979 to 1999. The distribution of the solar elongation angles of the sources observed is shown in a graph in their paper in the page 121101-2. Overlaid on this plot is a solid line showing the relationship between solar elongation angle and predicted deflection angle (right-hand scale). That means PPN formalism predicts a variable deflection angle unlike General Relativity. Another important visualizing graph shown by Shapiro et al, in the same paper is a plot in page 121101-4 indicating ‘differences between global estimates of γ and GR’s prediction as a function of minimum source elevation angle ϵ_{\min} ; and minimum solar elongation angle Φ_{\min} ; for positive declination sources and negative declination sources’. This clearly indicates the variation in the Gravitational deflection (bending) angle of light / radio waves near a huge gravitational mass like Sun from the variation of γ also. That means the observed variations are not just purely dependent on ϕ , the solar elongation angle alone. There are some more additional factors, in other than γ and ϕ like positive declination sources and negative declination sources, etc.

Robert Heinkelmann and Harald Schuh (2009) in their paper have concluded: “Besides the Sun and the Earth, the gravitational time delays of Jupiter, Saturn, Venus, and the Moon will have to be considered in standard VLBI 2010 analyses”.

Let us examine the work done by other Physicists who reported this phenomenon in section 4 of this paper. Many of them felt the need for considering the gravitational effect of other Planets in the Solar system on the radio wave photons.

1.3. PPN and VLBI

Such variations formulated as parameterized post-Newtonian (PPN) formalism (see, Books by Misner et. al., (1973), and Will (1993), for example). The angle θ calculated PPN formalism by which an electromagnetic ray from a distant source is deflected by the Sun, and is given approximately by, “ $(1+\gamma) G M (1+\cos \phi) / bc^2$ ”. Where G is the universal gravitational constant, c is the speed of light in vacuum, M is the mass of the Sun, b is the minimum distance of path of ray / particle to the center of Sun, ϕ is the solar elongation angle, and γ is the PPN parameter for the space curvature to gravitational deflection. Here ϕ is the main factor for explaining the observed variations, introduced in this PPN formalism. However, the range of observed variations overshoots possible theoretical variations.

1.4. Using Dynamic Universe model for VLBI

Now let us see how to explain the variations in the Gravitational deflection (bending) angle as plotted against solar elongation angle Φ , using Dynamic Universe Model. For doing so, the capabilities of Dynamic Universe Model are extended into Micro world or the Photon / Particle zoo. Dynamic Universe model can calculate the simultaneous gravitational effect of many gravitating bodies like Sun, planets, local stars etc., while considering their dynamic movements. For this purpose, the original mathematical background of Dynamic Universe Model was extended into micro world. The required additional mathematics is in the mathematical section along with the original set. Using these extended capabilities into SITA programming, the setup of solar system was as on 01.01.2000@00.00:00 hrs. using Heliocentric ecliptic xyz values and try sending the radio photon from different directions i.e., in different solar elongation angle Φ_{\min} and trace the path of radio photon. Here 76 different xyz coordinates and different directions were taken for radio photons with the same status of solar system as on 01.01.2000@00.00:00 hrs. The only change from experiment to experiment is the initial position and direction of the photon. All these theoretical experiments were designed in such a way, the photon goes grazingly near Sun or at the minimum distance from the center of Sun at the moment of time as on 01.01.2000@00.00:00 hrs. precisely. That means all the Solar system setup was kept constant and changed the Solar Elongation Angle only, taking into account all the dynamic movements of planets and the their gravitational fields on the fast moving photon. Each of these computationally intensive theoretical experiments took a time 15 min at the lowest to 5 hours at the highest, on a recent HP Laptop, depending on the number of iterations. This paper was presented as a talk at COSPAR-12 (H0.2-0010-12). Now let us see what Dynamic Universe Model of Cosmology is...

2. Mathematical Background: Into the Micro world

Now let’s see how the Dynamic Universe Model extends into Micro world mathematically.

2.1. Additional mathematical formulation allowing for the Micro world:

In addition to previously published mathematical formulation (viz. eqn 1 to 31) for Dynamic Universe Model, (Snp.Gupta 2014), the following new equations are added to enable this model for operating into Micro world.

Combining both 3 and 25 (Newly introduced in this paper)

$$F_{\alpha} = \sum_{\substack{\alpha=1 \\ \alpha \neq \beta}}^N F_{\alpha\beta} - m_{\alpha} \nabla_{\alpha} \left[- \sum_{\substack{\beta=1 \\ \alpha \neq \beta}}^{N^{\gamma}} \frac{Gm_{\beta}^{\gamma}}{|x^{\gamma\beta} - x^{\gamma\alpha}|} - \sum_{\substack{\beta=1 \\ \alpha \neq \beta}}^{N^{\delta\gamma}} \frac{Gm_{\beta}^{\delta\gamma}}{|x^{\delta\gamma\beta} - x^{\delta\gamma\alpha}|} \right] \quad (32)$$

So ...

$$F_{\alpha} = \sum_{\substack{\alpha=1 \\ \alpha \neq \beta}}^N F_{\alpha\beta} + m_{\alpha} \nabla_{\alpha} \left[\sum_{\substack{\beta=1 \\ \alpha \neq \beta}}^{N^{\gamma}} \frac{Gm_{\beta}^{\gamma}}{|x^{\gamma\beta} - x^{\gamma\alpha}|} + \sum_{\substack{\beta=1 \\ \alpha \neq \beta}}^{N^{\delta\gamma}} \frac{Gm_{\beta}^{\delta\gamma}}{|x^{\delta\gamma\beta} - x^{\delta\gamma\alpha}|} \right] \quad (33)$$

\sum and ∇

Interchanging

operations:

$$F_{\alpha} = \sum_{\substack{\alpha=1 \\ \alpha \neq \beta}}^N F_{\alpha\beta} + m_{\alpha} \left[\sum_{\substack{\beta=1 \\ \alpha \neq \beta}}^{N^{\gamma}} \nabla_{\alpha} \frac{Gm_{\beta}^{\gamma}}{|x^{\gamma\beta} - x^{\gamma\alpha}|} + \sum_{\substack{\beta=1 \\ \alpha \neq \beta}}^{N^{\delta\gamma}} \nabla_{\alpha} \frac{Gm_{\beta}^{\delta\gamma}}{|x^{\delta\gamma\beta} - x^{\delta\gamma\alpha}|} \right] \quad (34)$$

Or

$$F_{\alpha} = \sum_{\substack{\alpha=1 \\ \alpha \neq \beta}}^N F_{\alpha\beta} + \sum_{\substack{\beta=1 \\ \alpha \neq \beta}}^{N^{\gamma}} \nabla_{\alpha} \frac{Gm_{\alpha} m_{\beta}^{\gamma}}{|x^{\gamma\beta} - x^{\gamma\alpha}|} + \sum_{\substack{\beta=1 \\ \alpha \neq \beta}}^{N^{\delta\gamma}} \nabla_{\alpha} \frac{Gm_{\alpha} m_{\beta}^{\delta\gamma}}{|x^{\delta\gamma\beta} - x^{\delta\gamma\alpha}|} \quad (35)$$

The above equation 35 means the force on α th point mass will be a sum of three components i.e., the summation of attraction forces due to point masses from its own system, ensemble and aggregate. This is the key result that can be applied to point masses or subatomic particles or can be combined in any fashion together. The equations 34 & 35 are important, simple & straightforward results.

The theoretical Circular velocities are different to that of observed, the missing mass (Dark matter) arises due to Calculation error, there is no other reason and dark matter does not exist in reality (SNP Gupta (2005b)). In a present paper, about "there is no dark matter" it was shown that concepts like *relative constant Mass, variable mass and missing mass* etc., are not required. And the details of earlier publications, in books as well as papers are available in the same paper. One can refer to the same paper for main foundations and a general introduction for Dynamic Universe Model (SNP Gupta (2014)). One can see the references in this paper for the details..

Let us apply the above equations to some practical situations like radio waves that are going grazingly near Sun from an artificial satellite and are coming to Earth (VLBI). Let us use SITA algorithm for calculation.

3. Initial Values

3.1. Initial value of Mass of Photon:

Here in this paper the radio photon mass was taken as $1.1 \text{ e-}52 \text{ kg}$ which is much smaller than Neutrino mass = $1.5 \text{ ev} / c^2 = 2.685 \text{ e-}36 \text{ kg}$. Or Radio wave photon mass = $9.019 \text{ e-}46 \text{ kg}$ etc. The behaviour shown by such a mass will be experienced by higher mass also. Every mass behave in the similar way.

3.2. Initial Values table and other tables

Full set of initial values are shown in Table 1 web version. '**Vak Table 1 and 2 PLB Initial values Full set.xls**', Tab: Table1. ... The summary of all the 76 experiments conducted and reported in this paper can be seen in Table 2 in 13 columns.

Table 1. HELIO CENTRIC ECLIPTIC XYZ VALUES of positions as on 01.01.2000@00.00:00 hrs in Metre					
Sl.No.	Name	Mass (kg)	xecliptic	yecliptic	zecliptic
1	Photon (The initial Position vary with experiment)	1.10E-52	200000000	696000000	650947504.6
2	Mercury	3.30E+23	50644179263	8540296134	-3949485753
3	Venus	4.87E+24	69657878862	82614198079	-2889306238
4	Earth	5.97E+24	-29565785818	1.44E+11	-2869446.398
5	Mars	6.42E+23	-3275068912	-2.18E+11	-4484946284
6	Jupiter	1.90E+27	4.09E+11	-6.46E+11	-6473185584
7	Saturn	5.68E+26	-1.36E+12	3.40E+11	48167461412
8	Uranus	8.68E+25	2.98E+12	-4.32E+11	-40141525477
9	Neptune	1.02E+26	3.61E+12	-2.67E+12	-28350290138
10	Pluto	1.27E+22	69315882273	-4.70E+12	4.83E+11
11	Moon	7.35E+22	-29191657344	1.44E+11	16609650.17
12	SUN	1.99E+30	0	0	0
13	near star	3.98E+29	-3.07E+16	-2.48E+16	5.99E+15
14	near star	1.89E+30	-1.70E+16	-4.50E+13	3.79E+16
15	near star	2.19E+30	-1.72E+16	-1.53E+14	3.79E+16
16	near star	7.95E+29	-1.86E+15	1.64E+15	-5.61E+16
17	near star	8.95E+29	9.03E+15	-7.13E+15	-7.78E+16

76 theoretical experiments were conducted in Dynamic Universe Model with various initial conditions all with 0.1 sec time step and with number of iterations ranging from 20 to 3000. For all these experiments the starting coordinates are heliocentric ecliptic xyz values in our solar system as on 01.01.2000@00.00:00 hrs with masses in Kg and distances are in Metre. The execution times were ranging from 20 minutes to 24 hours on this modern

laptop (Hp make, with Intel core i5 processor, 8Gb Ram, 650Gb HDD, NVIDIA display Hp pavilion dv6).

Full set of initial values are shown in Table 1 web version. This Table 1 gives the initial values of all the point masses, their respective masses in Kg, and their x y z Cartesian positions. Please refer the Table 1 in the web with name: 'Table 1.xls'. (**34445-116835-1-SP.xls**) ... The summary of all the 76 experiments conducted and reported in this paper can be seen in Table 2 in 13 columns. The various filenames, Time step values, Number of iterations, initial position, and final position of the Photon, General relativistic deflection (bending) angle, Calculated Solar elongation angle in degrees, SITA Deflection etc., are mentioned in the web with name: 'Table 2.xls'. (**34445-116836-1-SP.xls**) This table 2 is the summary and condensed from the Table 3, where there are 71 columns. So the finer differences between each experiment can be found from main Table 3 easily. This Table 3 is uploaded into web along with main file. File name is '**Vak Table 3 Results 71 columns.xls**' (**34445-116837-1-SP.xls**). There are 76 data records of final iteration data for each file in 71 columns in a single Excel sheet. The header rows for all the data is in the 3rd and 4th rows. As a result of SITA calculations, there is an assortment of xyz positions, velocities and accelerations in all iterations at each time step for every one of the 133 point masses. It is possible to track any one of these values for every time step throughout the experiment as needed. The Table 4 contains all the individual iteration output for each iteration of the 76 experiments. This data is in an XLS file and uploaded in Web. There are 19681 lines in total in these files in 55 columns. There will be additional blank rows before the next file data. Each data set varies in number of data rows depending on number of iterations. Each row represents a set of data for that time step. Number of data rows from file to file varies from 40 to 3000. Refer file name: **Vak Table 4 Consolidation of Iteration Results.xls**. This file is being very big; it was divided into three parts as it was very lengthy viz.. (**34445-116839-1-SP.xls**), (**34445-116840-1-SP.xls**), (**34445-116841-1-SP.xls**) Any further additional details can be obtained by contacting the author. All these tables are available in the webpage which can be down loaded. These **1. Table 1.xls**, **2. Table 2.xls**, **3. Vak Table 3 Results 71 columns.xls** and **4. Vak Table 4. Consolidation of Iteration Results.xls** are attached with this paper. (See the **34445 series numbers in bold for the attached files here**)

4. VLBI Discussion:

Here in this paper the capabilities of Dynamic Universe Model are extended into Micro world or the Photon / Particle zoo. Then the question comes how to test this solution. There are many papers on VLBI which discussed about variation of solar deflection (bending) angle, and let us see some of the views expressed by different authors.

4.1. Shapiro et al (2004), in their paper 'Measurement of Solar Gravitational Deflection of VLBI data of Radio waves, shown their data set as consisting of measurements from observations in about 2500 _24-hour sessions ("experiments") spanning the years 1979–1999. The experiments that they used in this study involved 87VLBI sites and 541 radio sources, yielding a total of more than 1.7×10^6 ionosphere-corrected group-delay measurements. Their overall observations clearly the variation in the Gravitational deflection angle of light near a huge gravitational mass like Sun with respect to solar elongation angle.

4.2. The authors Ojars J. Sovers et al (1998), also mentioned in their results about the variation, their 83 page paper is very good for beginners on the VLBI. They gave a detailed description of different experiments, a

good history and the experiments that gave higher accuracies for angular positioning for point-like extragalactic radio sources at the sub-milliarcsecond (nanoradian) level.

4.3. The authors E. Fomalont et al in 2009 observed position changes with respect to session and Frequency in their paper. *One can infer the reason for the observed position changes is the effect of Universal Gravitational force (Let's call this as UGF). This UGF varies as the configuration of the experimental set up varies from session to session. Here configuration means the positions of Earth, Sun, the observing satellite and the occulting planet etc., at the time of observation.*

4.4. In other words, Ed Fomalont et al in 2010 reported a larger error which cannot be explained by conventional methods.....In their own words as explanation for their 'figure three'... "The Residual Differential Separation between Cassini and J1127+0555: The separation between the two sources, after removal of the best-fit offset, velocity and acceleration from the measured offsets. The predicted gravitational deflection (bending) of Cassini and J1127+0555 has been removed, so the difference should be zero if the GR prediction is correct. The left plot is the east-west difference; the right plot is the north-south difference. The position difference estimates were obtained for the three 2.5-hour intervals on each day, and the error are the one-sigma errors expected from signal-to-noise considerations alone" Additionally in the text also... "If we further remove the radial and aberration deflection prediction by GR, then the resultant relative position of Cassini with respect to J1127 + 0555 should be zero. This position difference with the above adjustments is shown in the paper. The departure of the residuals from zero has an error of about 0.01 mas E/W, and 0.02 mas N/S (this resolution is twice as poor). Since the aberrational deflection is 0.07 mas at closest encounter on February 10, this experiment may be a more accurate measure of the aberration deflection than that of the 2002 Jupiter experiment Fomalont & Kopeikin in 2003 ." *Ed Fomalont is telling about large errors.*

4.5. D. E. Lebach et al, in 1995 , estimated γ from each of the single-day data subsets that included one days' data from before the solar occultation of quasar 3C279 and one from after. These pair wise estimates ranged from $\gamma = 0.9440 \pm 0.0350$ to $\gamma = 1.0214 \pm 0.0338$ (SSEs : statistical standard error); six of these estimates deviated from $\gamma = 1$ by more than 1.5 SSEs, and only one ($\gamma = 0.9903 \pm 0.0035$), which is based on 2 and 23 GHz data from 4 and 11 October, deviated by $\gamma = 1$ by more than 2.5 SSEs. *The overall estimates of γ by Lebach et al (1995) [4] from different data subsets may not show statistically unusual deviation. But what they are showing us are the data subsets at the extreme limits where deviation from the calculated value using GR is very high. Here also the reason is the effect of UGF varies on the radio photon, which depends on the configuration of universe at the time of observation.*

4.6. The conclusion of Robert Heinkelmann and Harald Schuh (2009) that gravitation of other planets (6 bodies viz., Sun, Earth, Jupiter, Saturn, Venus, and the Moon) will have to be considered in VLBI is an important factor for consideration." Scientific community know that 3-body problem is not solvable for the last 500 years in the conventional methods. Dynamic Universe model is a singularity free 3-body or 7 body or general n-body problem solution. *Dynamic universe model considers Newtonian gravitational effect of 133 bodies (or of any arbitrary N bodies) without using General Relativity. Hence such problems can be solved easily.*

4.7. The authors ‘S. B. Lambert and C. Le Poncin-Lafitte’ in 2009 also concluded... “we wish to mention that, although current VLBI appears to not be competitive with spacecraft systems for relativistic experiments, the huge number of VLBI measurements, in all directions and at a large number of epochs, constitutes an interesting potential for testing other theories than the PPN formalism. “Here Lambert & Le Poncin-Lafitte [9] are inviting other types of theories...”

4.8 There are a large number of papers / literature available on VLBI, giving similar conclusions by other Physicists, which I could not mention here.

5. Resulting Graph:




Fig 1. shows the gravitational deflection (bending) angles calculated from the 76 theoretical experiments, using Dynamic Universe Model’s SITA software. Here the setup of solar system was taken as was on 01.01.2000@00.00:00 hrs for all these experiments, but the incoming radio photon direction was different in all these experiments. Some deflection angles (the angle between incoming and outgoing ray of radio wave) from SITA experiments were higher than the GR prediction and others have different values. It may please be noted that GR predicts only one deflection angle for all the incident angles of the ray and does not depend on solar system configuration at that time. PPN formalism also cannot explain fully about these variations (see fig given by Shapiro et al (2004), for details see the text of paper).

6. Discussion on Results:

Many physicists require some other additional factors to be considered, like simultaneous and dynamical gravitation effects of other planets. Now let us summarize what the other physicists concluded on VLBI. Shapiro et al showed the variation in the Gravitational deflection angle with respect to solar elongation angle. Fomalont et al in 2009 observed position changes with respect to session and Frequency. Fomalont et al in 2010 reported larger errors which cannot be explained by conventional methods and difference should be zero if

the GR prediction is correct. Lebach et al, in 1995, in their study of the occultation of quasar 3C279 concluded that the error gone up by more than 2.5 times the statistical standard error. Robert Heinkelmann and Harald Schuh (2009) in their paper have concluded: “Besides the Sun and the Earth, the gravitational time delays of Jupiter, Saturn, Venus, and the Moon will have to be considered in standard VLBI2010 analyses. Lambert and Le Poncin-Lafitte’ in 2009, also concluded... “We wish to mention that, although current VLBI appears to not be competitive with spacecraft systems for relativistic experiments, the huge number of VLBI measurements, in all directions and at a large number of epochs, constitutes an interesting potential for testing other theories than the PPN formalism.

The resulting Gravitational deflection (bending) angles from Dynamic Universe model’s SITA software are in line with the thinking of present day VLBI physicists as we saw in the paper. The results are plotted and shown in a graph in Fig 1. This graph shows the Gravitational deflection angle of Radio wave / particles near SUN, which was calculated using Dynamic Universe Model and SITA algorithms plotted against the solar elongation angle (angle between the Sun and the source as viewed from Earth). For explaining such variation, even after taking the help of PPN formalism General relativity could not provide satisfactory explanation. Here it is not necessary. All these variations in angle, some of which are more than GR predictions and some are less than spread over solar elongation angle, are direct calculations of Dynamic Universe Model using Universal Gravitational Force calculated for the setup of Universe at that instant.

7. Conclusions:

Dynamic Universe model gives better results compared to GR. GR gives only one value irrespective of positions of solar system and angle of incidence. The observed deflection values are sometimes more and sometimes less than GR predicted values.

In this paper, a set of theoretical simulations were done, in which, the setup of solar system was taken as on 01.01.2000@00.00:00 hrs, and Heliocentric ecliptic xyz values were used for all the theoretical experiments. The only difference between experiments is the direction of the ray of radio photon in the starting (angle of incidence of the ray). In other words the radio photon was sent in different solar elongation angles (Φ_{min}). And Dynamic Universe model is used for tracing the path of this radio photon. Here 76 different xyz coordinates and different directions were taken for radio photons with the same status of solar system as on 01.01.2000@00.00:00 hrs.

All these final deflection angles are different in value. These results indicate that the structure of our universe is continuously changing dynamically, and depending on Universal Gravitation Force (UGF) on the radio photon in that path, the trajectory of the ray radio wave is different. The UGF on the particles of the ray is based positions and distances of Planets, stars, Galaxies etc., relative to radio photon and varies the dynamically. In addition, it is observed all the point masses follow their own fixed trajectory irrespective of its own mass. That means by changing only value of mass under consideration, the trajectory followed by it will not change by keeping all the other things constant.

Hence, it can be concluded that the gravitational deflection near SUN will depend not only on incident angle of the incoming ray but also dynamic positions of the planets at that time and various other mass positions in the Universe and Universal Gravitation Force. For explaining, the Very Long Baseline Interferometry (VLBI) gravitational deflection (bending) observations, Dynamic Universe Model can be successfully used. This become possible as the capabilities of Dynamic Universe Model could be extended into micro world in the mathematical section, by extending the original mathematical formulation. It can be safely concluded that Dynamic Universe

Model is better over PPN formalism for explaining variations in VLBI solar deflection observations. By taking actual values of positions, and velocities planets and stars etc., we can calculate the real values of deflections.

7. Acknowledgements:

I gracefully acknowledge the guidance given by the Lord of Puri for extending the Mathematics. I gracefully acknowledge the efforts taken by Mrs. Sheetal of SAP (USA) for making grammar corrections in this document.

8. References:

- Akerib, D. S., Araujo, H. M., Bai, X., Bailey, A. J., Balajthy, J., Bedikian, S., ... Zhang, C. (2013). First results from the LUX dark matter experiment at the Sanford Underground Research Facility. Retrieved from <http://arxiv.org/abs/1310.8214>
- Akerib, D. S., Bernardo, E., Bernstein, A., Bradley, A., Byram, D., ... Solovov, V. N. (2013). Technical results from the surface run of the LUX dark matter experiment. *Astroparticle Physics*, 45, 34-43. <http://dx.doi.org/10.1016/j.astropartphys.2013.02.001>
- Colless M.M., Dalton G.B., Maddox S.J., Sutherland W.J., Norberg P., Cole S.M., Bland-Hawthorn J., Bridges T.J., Cannon R.D., Collins C.A., Couch W.J., Cross N., Deeley K., De Propriis R., Driver S.P., Efstathiou G., Ellis R.S., Frenk C.S., Glazebrook K., Jackson C.A., Lahav O., Lewis I.J., Lumsden S., Madgwick D.S., Peacock J.A., Peterson B.A., Price I.A., Seaborne M., Taylor K., 2001, MNRAS, 328, 1039 [ADS | astro-ph/0106498] See 2dFGRS publications <http://www.mso.anu.edu.au/2dFGRS/>
- Cruz, Martínez-González, Vielva & Cayón (2005), "Detection of a non-Gaussian Spot in WMAP", MNRAS 356 29-40 [astro-ph/0405341]
- Fomalont, E. S. Kopeikin, G. Lanyi, J. Benson, (2009) paper name 'Progress in Measurements of the Gravitational Bending of Radio Waves Using the VLBA' in section 4.3, and Table 4 <http://arxiv.org/abs/0904.3992v1>
- Fomalont, Ed Sergei Kopeikin, Dayton Jones, Mareki Honma & Oleg Titov in their paper 'Recent VLBA/VERA/IVS Tests of General Relativity' in Figure 3. (2010) <http://cienciaescolar.net/proyectos/wp-content/uploads/2010/01/0912.3421v1.pdf> ... Book 'Relativity in Fundamental Astronomy (IAU S261): Dynamics, Reference Frames' ... Proceedings of IAU Symposium No.261, 2009. Page 286
- Fomalont, Ed , Sergei Kopeikin [2003], (arXiv:astro-ph/0302294) , "The Measurement of the Light Deflection from Jupiter: Experimental Results" DOI: 10.1086/378785, [2003ApJ...598..704F](https://doi.org/10.1086/378785)
- Fairall, A. P., Palumbo, G. G. C., Vettolani, G., Kauffman, G., Jones, A., & Baiesi- 1990MNRAS.247P..21F. See in Wikipedia "The Large scale structure of cosmos" http://en.wikipedia.org/wiki/Large-scale_structure_of_the_cosmos H.P. Robertson, Kinematics and world Structure III , The Astrophysical Journal, May 1936, vol 83 pp 257.
- Gott et al., 2005, ApJ, 624, 463 <http://www.journals.uchicago.edu/ApJ/journal/issues/ApJ/v624n2/59364/59364.html> <http://www.astro.princeton.edu/universe/ms.pdf> See for various maps of Universe and Fig 8 at: <http://www.astro.princeton.edu/universe/>
- Lawrence Rudnick, Shea Brown, Liliya R. Williams, Extragalactic Radio Sources and the WMAP Cold Spot, *Astrophysics journal* and arXiv:0704.0908v2 [astro-ph],

- Lebach, D. E. B. E. Corey, I. I. Shapiro, M. I. Ratner, J. C. Webber, A. E. E. Rogers, J. L. Davis, and T. A. Herring, in their paper ‘Measurement of the Solar Gravitational Deflection of Radio Waves Using Very-Long-Baseline Interferometry’ in Phys. Rev. Lett. 75, 1439–1442 (1995),
- Misner, C.W. K. S. Thorne, and J. A. Wheeler, Gravitation (Freeman, San Francisco, 1973), Chap. 39.
- Robert Heinkelmann and Harald Schuh in Relativity in Fundamental Astronomy Proceedings IAU Symposium No. 261, 2009, edited by S. A. Klioner, P. K. Seidelman & M. H. Soffel, c International Astronomical Union 2010 doi:10.1017/S1743921309990524 title of paper ‘Very long baseline interferometry: accuracy limits and relativistic tests’, Pages 286-290
- NY Times (2013). Dark-matter-experiment-has-found-nothing-scientists-say-proudly. Retrieved from http://www.nytimes.com/2013/10/31/science/space/dark-matter-experiment-has-found-nothing-scientists-say-proudly.html?_r=0
- Ojars J. Sovers, John L. Fanselow, Christopher S. Jacobs, Jet Propulsion Laboratory, ‘Astrometry and geodesy with radio interferometry: experiments, models, results’ Rev. Mod. Phys. 70, 1393 – Published 1 October 1998, DOI: <http://dx.doi.org/10.1103/RevModPhys.70.1393>, <http://arxiv.org/abs/astro-ph/9712238v1>
- S.Samurovic et al; 0811.0698v1 Arxiv, ‘Mond vs Newtonian dynamics GC’, A&A accepted Nov 5, 2008. (For mass calculations)
- SNP. Gupta web the following very big tables, whose names are given below, needed along with main paper.
1. Table 1.xls
 2. Table 2.xls
 3. Vak Table 3 Results 71 columns.xls
 4. Vak Table 4 Consolidation of Iteration Results.xls
- These files are available at the web page for all the above full tables:
<https://skydrive.live.com/#cid=485CC4B593A12043&id=485CC4B593A12043%213522>
- SNP. Gupta “Dynamic Universe Model’s Prediction “No Dark Matter” in the Universe Came True!” Applied Physics Research; Vol. 6, No. 2; 2014, Page 8, ISSN 1916-9639 E-ISSN 1916-9647, Published by Canadian Center of Science and Education URL: <http://dx.doi.org/10.5539/apr.v6n2p8>
doi:10.5539/apr.v6n2p8
- SNP. Gupta. (2005a). Dynamic Universe Model of cosmology: Missing mass in Galaxy. Presented in 7th Astronomical conf by HEL.A.S., Kefallinia, Greece 8-11, Sept, 2005.
- SNP. Gupta. (2005b). On Missing mass, “DYNAMIC UNIVERSE MODEL of cosmology: Missing mass in Galaxy” Presented at OMEG05 Origin of Matter and Evolution of Galaxies, Tokyo university, Tokyo, JAPAN.
- SNP. Gupta. (2007). “Dynamic Universe Model of cosmology: Missing mass in Galaxy” presented in PATHWAYS THROUGH AN ECLECTIC UNIVERSE conf held at Tenerife, Spain, in 23rd to 27th 2007 April
- SNP. Gupta. (2010). *Dynamic Universe Model: A singularity-free N-body problem solution*. Germany: VDM Verlag Dr. Müller.
- SNP. Gupta. (2011a). *Dynamic Universe Model: SITA singularity free software*. Germany: VDM Verlag Dr. Müller.
- SNP. Gupta. (2011b). *Dynamic Universe Model: SITA software simplified*. Germany: VDM Verlag Dr. Müller.
- SNP. Gupta. (2012a). *Dynamic Universe Model: Blue Shifted Galaxies Prediction*. Germany: LAP Publisher.
- SNP. Gupta. (2012b). Singularity free N-body simulations DUMOC No-dark matter. COSPAR13, Retrieved

from <https://skydrive.live.com/?id=485CC4B593A12043!135&cid=485cc4b593a12043> “H0.1-0023-12 POSTER room sun-sat July 15 1600 A1 size--- Singularity free N-body simulations DUMOC No-dark matter.bmp”

SNP. Gupta. (2013). Introduction to Dynamic Universe Model. *International Journal of Scientific Research and Reviews*, 2(1), 203-226

SNP. Gupta. (2014), *Applied Physics Research* ISSN 1916-9639 (Print) ISSN 1916-9647 (Online) DOI: 10.5539/apr.v6n2p8, [Vol 6, No 2 \(2014\)](#) Title of paper: “Dynamic Universe Model’s Prediction “No Dark Matter” in the Universe Came True!”

S. S. Shapiro, J. L. Davis, D. E. Lebach, and J. S. Gregory ‘Measurement of the Solar Gravitational Deflection of Radio Waves using Geodetic Very-Long-Baseline Interferometry Data, 1979–1999’ See Fig. 1. PHYSICAL REVIEW LETTERS, VOLUME 92, NUMBER 12, 26 MARCH 2004 121101-1 to 121101- 4.

UC Davis. Large Underground Xenon at UC Davis, JOURNAL: “Physics”, Retrieved from <http://lux.physics.ucdavis.edu/> 30 Oct 2013 <http://dx.doi.org/10.1088/1748-0221/8/10/C10003>

Will, C.M. Theory and Experiment in Gravitational Physics (Cambridge University Press, Cambridge, England, 1993).

Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.

This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).

.....