

On the properties of generalized multiplicative coupled fibonacci sequence of r^{th} order

Research Article
A. D. Godase^{1,*}, M. B. Dhakne²
¹*Department of Mathematics, V. P. College, Vaijapur, India*
²*Department of Mathematics, Dr. B. A. M. University, Aurangabad, India*
Received 02 February 2015; accepted (in revised version) 28 February 2015

Abstract: Coupled Fibonacci sequences of lower order have been generalized in number of ways. In this paper the Multiplicative Coupled Fibonacci Sequence has been generalized for r^{th} order with some new interesting properties.

MSC: 11B39 • 11B37 • 11B99

Keywords: Fibonacci sequence • Coupled Fibonacci sequence • Recurrence relation

© 2015 IJAAMM all rights reserved.

1. Introduction

Definition 1.1.

The Multiplicative Coupled Fibonacci Sequence of 2^{nd} order is defined as, Let $\{X_i\}_{i=0}^{i=\infty}$ and $\{Y_i\}_{i=0}^{i=\infty}$ be two infinite sequences and four arbitrary real numbers a, b, c, d are given. The Multiplicative Coupled Fibonacci Sequence of 2^{nd} order is generated by the following four different ways:

First scheme:

$$\begin{aligned} X_{n+2} &= X_{n+1} \cdot X_n, n \geq 0 \\ Y_{n+2} &= Y_{n+1} \cdot Y_n, n \geq 0 \end{aligned}$$

Second scheme:

$$\begin{aligned} X_{n+2} &= Y_{n+1} \cdot X_n, n \geq 0 \\ Y_{n+2} &= X_{n+1} \cdot Y_n, n \geq 0 \end{aligned}$$

Third scheme:

$$\begin{aligned} X_{n+2} &= X_{n+1} \cdot Y_n, n \geq 0 \\ Y_{n+2} &= Y_{n+1} \cdot X_n, n \geq 0 \end{aligned}$$

Fourth scheme:

$$\begin{aligned} X_{n+2} &= Y_{n+1} \cdot Y_n, n \geq 0 \\ Y_{n+2} &= X_{n+1} \cdot X_n, n \geq 0 \end{aligned}$$

* Corresponding author.

E-mail address: ashokgodse2012@gmail.com

Definition 1.2.

The Multiplicative Coupled Fibonacci Sequence of 3^{rd} order is defined as, Let $\{X_i\}_{i=0}^{i=\infty}$ and $\{Y_i\}_{i=0}^{i=\infty}$ be two infinite sequences and six arbitrary real numbers a, b, c, d, e, f are given. The Multiplicative Coupled Fibonacci Sequence of 3^{rd} order is generated by the following eight different ways:

First scheme:

$$\begin{aligned} X_{n+3} &= Y_{n+2} \cdot Y_{n+1} \cdot Y_n, n \geq 0 \\ Y_{n+3} &= X_{n+2} \cdot X_{n+1} \cdot X_n, n \geq 0 \end{aligned}$$

Second scheme:

$$\begin{aligned} X_{n+3} &= X_{n+2} \cdot X_{n+1} \cdot X_n, n \geq 0 \\ Y_{n+3} &= Y_{n+2} \cdot Y_{n+1} \cdot Y_n, n \geq 0 \end{aligned}$$

Third scheme:

$$\begin{aligned} X_{n+3} &= Y_{n+2} \cdot Y_{n+1} \cdot X_n, n \geq 0 \\ Y_{n+3} &= X_{n+2} \cdot X_{n+1} \cdot Y_n, n \geq 0 \end{aligned}$$

Fourth scheme:

$$\begin{aligned} X_{n+3} &= Y_{n+2} \cdot X_{n+1} \cdot Y_n, n \geq 0 \\ Y_{n+3} &= X_{n+2} \cdot Y_{n+1} \cdot X_n, n \geq 0 \end{aligned}$$

Fifth scheme:

$$\begin{aligned} X_{n+3} &= Y_{n+2} \cdot X_{n+1} \cdot X_n, n \geq 0 \\ Y_{n+3} &= X_{n+2} \cdot Y_{n+1} \cdot Y_n, n \geq 0 \end{aligned}$$

Sixth scheme:

$$\begin{aligned} X_{n+3} &= X_{n+2} \cdot X_{n+1} \cdot Y_n, n \geq 0 \\ Y_{n+3} &= Y_{n+2} \cdot Y_{n+1} \cdot X_n, n \geq 0 \end{aligned}$$

Seventh scheme:

$$\begin{aligned} X_{n+3} &= X_{n+2} \cdot Y_{n+1} \cdot Y_n, n \geq 0 \\ Y_{n+3} &= Y_{n+2} \cdot X_{n+1} \cdot X_n, n \geq 0 \end{aligned}$$

Eighth scheme:

$$\begin{aligned} X_{n+3} &= X_{n+2} \cdot Y_{n+1} \cdot X_n, n \geq 0 \\ Y_{n+3} &= Y_{n+2} \cdot X_{n+1} \cdot Y_n, n \geq 0 \end{aligned}$$

In recent years many authors have been generalized Coupled Fibonacci sequences of lower order in number of ways. In this paper the Multiplicative Coupled Fibonacci Sequence has been generalized for r^{th} order with some new interesting properties.

2. Multiplicative Coupled Fibonacci Sequence of r^{th} order

Definition 2.1.

The Multiplicative Coupled Fibonacci Sequence of r^{rd} order is defined as, Let $\{X_i\}_{i=0}^{i=\infty}$ and $\{Y_i\}_{i=0}^{i=\infty}$ be two infinite sequences and $2r$ arbitrary real numbers $x_0, x_1, x_2, x_3, \dots, x_{r-1}$ and $y_0, y_1, y_2, y_3, \dots, y_{r-1}$ are given. The Multiplicative Coupled Fibonacci Sequence of r^{th} order is generated by the following 2^r different ways:

First scheme:

$$\begin{aligned} X_{n+r} &= Y_{n+r-1} \cdot Y_{n+r-2} \cdot Y_{n+r-3} \cdots Y_n, n \geq 0 \\ Y_{n+r} &= X_{n+r-1} \cdot X_{n+r-2} \cdot X_{n+r-3} \cdots X_n, n \geq 0 \end{aligned}$$

⋮

$(2^r)^{th}$ scheme:

$$\begin{aligned} X_{n+r} &= X_{n+r-1} \cdot X_{n+r-2} \cdot X_{n+r-3} \cdots X_n, n \geq 0 \\ Y_{n+r} &= Y_{n+r-1} \cdot Y_{n+r-2} \cdot Y_{n+r-3} \cdots Y_n, n \geq 0 \end{aligned}$$

are given below:

Table 1. First few terms of the sequences under 2^{rth} scheme

n	X_{n+r}	Y_{n+r}
0	$y_0 \cdot y_1 \cdot y_2 \cdot y_3 \cdots \cdot y_{r-1}$	$x_0 \cdot x_1 \cdot x_2 \cdot x_3 \cdots \cdot x_{r-1}$
1	$x_0 \cdot x_1 \cdot x_2 \cdot x_3 \cdots \cdot x_{r-1} \cdot y_1 \cdot y_2 \cdot y_3 \cdots \cdot y_{r-1}$	$x_1 \cdot x_2 \cdot x_3 \cdots \cdot x_{r-1} \cdot y_0 \cdot y_1 \cdot y_2 \cdot y_3 \cdots \cdot y_{r-1}$
2	$x_0^2 \cdot x_1^2 \cdot x_2^2 \cdot x_3^2 \cdots \cdot x_{r-1}^2 \cdot y_0 \cdot y_1 \cdot y_2^2 \cdot y_3^2 \cdots \cdot y_{r-1}^2$	$x_0 \cdot x_1 \cdot x_2^2 \cdot x_3^2 \cdots \cdot x_{r-1}^2 \cdot y_0 \cdot y_1^2 \cdot y_2^2 \cdot y_3^2 \cdots \cdot y_{r-1}^2$
3	$x_0^3 \cdot x_1^3 \cdot x_2^4 \cdot x_3^4 \cdots \cdot x_{r-1}^4 \cdot y_0^2 \cdot y_1^3 \cdot y_2^3 \cdot y_3^4 \cdots \cdot y_{r-1}^4$	$x_0^2 \cdot x_1^3 \cdot x_2^3 \cdot x_3^4 \cdots \cdot x_{r-1}^4 \cdot y_0^2 \cdot y_1^3 \cdot y_2^4 \cdot y_3^4 \cdots \cdot y_{r-1}^4$
4	$x_0^4 \cdot x_1^5 \cdot x_2^7 \cdot x_3^8 \cdots \cdot x_{r-1}^8 \cdot y_0^4 \cdot y_1^6 \cdot y_2^7 \cdot y_3^7 \cdot y_4^8 \cdots \cdot y_{r-1}^8$	$x_0^4 \cdot x_1^5 \cdot x_2^7 \cdot x_3^8 \cdots \cdot x_{r-1}^8 \cdot y_0^4 \cdot y_1^6 \cdot y_2^7 \cdot y_3^8 \cdots \cdot y_{r-1}^8$

3. Main Results

In this section many of the fabulous properties of generalized multiplicative coupled Fibonacci sequence of r^{th} order under 2^{rth} scheme are established.

Theorem 3.1.

For every integer $n \geq 0, r \geq 0$

$$X_{n(r+1)} \cdot Y_0 = Y_{n(r+1)} \cdot X_0 \quad (1)$$

Proof. If $n = 0$, then result is true because

$$X_0 \cdot Y_0 = Y_0 \cdot X_0$$

Assume that the result is true for some integer $n \geq 1$

Now,

$$\begin{aligned} X_{n+r} &= Y_{n+r-1} \cdot Y_{n+r-2} \cdot Y_{n+r-3} \cdots Y_n \\ Y_{n+r} &= X_{n+r-1} \cdot X_{n+r-2} \cdot X_{n+r-3} \cdots X_n, \end{aligned}$$

Using Induction Method,

For, $n + 1$

$$\begin{aligned} X_{(n+1)(r+1)} \cdot Y_0 &= [Y_{n(r+1)+r} \cdot Y_{n(r+1)+r-1} \cdot Y_{n(r+1)+r-2} \cdots Y_{n(r+1)+1}] \cdot Y_0 \\ &= [X_{n(r+1)+(r-1)} \cdot X_{n(r+1)+(r-2)} \cdot X_{n(r+1)+(r-3)} \cdots X_{n(r+1)}] \cdot [Y_{n(r+1)+r-1} \cdot Y_{n(r+1)+r-2} \cdots Y_{n(r+1)+1}] \cdot Y_0 \\ &= [X_{n(r+1)+(r-1)} \cdot X_{n(r+1)+(r-2)} \cdot X_{n(r+1)+(r-3)} \cdots X_{n(r+1)+1}] \cdot [Y_{n(r+1)+r-1} \cdot Y_{n(r+1)+r-2} \cdots Y_{n(r+1)+1}] [X_{n(r+1)} \cdot Y_0] \end{aligned}$$

Using induction hypothesis,

$$\begin{aligned} &= [X_{n(r+1)+(r-1)} \cdot X_{n(r+1)+(r-2)} \cdot X_{n(r+1)+(r-3)} \cdots X_{n(r+1)+1}] \cdot [Y_{n(r+1)+r-1} \cdot Y_{n(r+1)+r-2} \cdots Y_{n(r+1)+1}] [Y_{n(r+1)} \cdot X_0] \\ &= [X_{n(r+1)+(r-1)} \cdot X_{n(r+1)+(r-2)} \cdot X_{n(r+1)+(r-3)} \cdots X_{n(r+1)+1}] \cdot [X_{n(r+1)+r} \cdot X_0] \\ &= [Y_{n(r+1)+r+1} \cdot X_0] \\ &= [Y_{(n+1)(r+1)} \cdot X_0] \end{aligned}$$

Hence result is true for $n + 1$. □

Theorem 3.2.

For every integer $n \geq 0, r \geq 0$

$$X_{n(r+1)+1} \cdot Y_1 = Y_{n(r+1)+1} \cdot X_1 \quad (2)$$

Proof. : If $n = 0$, then result is true because

$$X_1 \cdot Y_1 = Y_1 \cdot X_1$$

Assume that the result is true for some integer $n \geq 1$

Now,

$$\begin{aligned} X_{n+r} &= Y_{n+r-1} \cdot Y_{n+r-2} \cdot Y_{n+r-3} \cdots Y_n \\ Y_{n+r} &= X_{n+r-1} \cdot X_{n+r-2} \cdot X_{n+r-3} \cdots X_n, \end{aligned}$$

Using Induction Method,
For, $n + 1$

$$\begin{aligned} X_{(n+1)(r+1)+1} \cdot Y_1 &= X_{n(r+1)+(r+2)} \cdot Y_1 \\ &= [Y_{n(r+1)+r+1} \cdot Y_{n(r+1)+r} \cdot Y_{n(r+1)+r-1} \cdots Y_{n(r+1)+2}] \cdot Y_1 \\ &= [X_{n(r+1)+r} \cdot X_{n(r+1)+(r-1)} \cdot X_{n(r+1)+(r-2)} \cdots X_{n(r+1)+1}] \cdot [Y_{n(r+1)+r} \cdot Y_{n(r+1)+r-1} \cdots Y_{n(r+1)+2}] \cdot Y_1 \\ &= [X_{n(r+1)+r} \cdot X_{n(r+1)+(r-1)} \cdot X_{n(r+1)+(r-2)} \cdots X_{n(r+1)+2}] \cdot [Y_{n(r+1)+r} \cdot Y_{n(r+1)+r-1} \cdots Y_{n(r+1)+2}] [X_{n(r+1)+1} \cdot Y_1] \end{aligned}$$

Using induction hypothesis,

$$\begin{aligned} &= [X_{n(r+1)+r} \cdot X_{n(r+1)+(r-1)} \cdot X_{n(r+1)+(r-2)} \cdots X_{n(r+1)+2}] \cdot [Y_{n(r+1)+r} \cdot Y_{n(r+1)+r-1} \cdots Y_{n(r+1)+2}] [Y_{n(r+1)+1} \cdot X_1] \\ &= [X_{n(r+1)+r} \cdot X_{n(r+1)+(r-1)} \cdot X_{n(r+1)+(r-2)} \cdots X_{n(r+1)+2}] \cdot [X_{n(r+1)+(r+1)} \cdot X_1] \\ &= [Y_{n(r+1)+(r+1)+1} \cdot X_1] \\ &= [Y_{(n+1)(r+1)+1} \cdot X_1] \end{aligned}$$

Hence result is true for $n + 1$. □

In the similar way, it could be proved for the following results by induction method.

Theorem 3.3.

:For every integer $n \geq 0, r \geq 0$

$$X_{n(r+1)+2} \cdot Y_2 = Y_{n(r+1)+2} \cdot X_2 \quad (3)$$

Theorem 3.4.

:For every integer $n \geq 0, r \geq 0$

$$X_{n(r+1)+3} \cdot Y_3 = Y_{n(r+1)+3} \cdot X_3 \quad (4)$$

Theorem 3.5.

:For every integer $n \geq 0, r \geq 0$ and $m \geq 0$

$$X_{n(r+1)+m} \cdot Y_m = Y_{n(r+1)+m} \cdot X_m \quad (5)$$

Proof. : If $n = 0$, then result is true because

$$X_m \cdot Y_m = Y_m \cdot X_m$$

Assume that the result is true for some integer $n \geq 1$

Now,

$$\begin{aligned} X_{n+r} &= Y_{n+r-1} \cdot Y_{n+r-2} \cdot Y_{n+r-3} \cdots Y_n \\ Y_{n+r} &= X_{n+r-1} \cdot X_{n+r-2} \cdot X_{n+r-3} \cdots X_n, \end{aligned}$$

Using Induction Method,
For, $n + 1$

$$\begin{aligned} X_{(n+1)(r+1)+m} \cdot Y_m &= X_{n(r+1)+(r+m+1)} \cdot Y_m \\ &= [Y_{n(r+1)+r+m} \cdot Y_{n(r+1)+r+m-1} \cdot Y_{n(r+1)+r+m-2} \cdots Y_{n(r+1)+m+1}] \cdot Y_m \\ &= [X_{n(r+1)+r+m-1} \cdot X_{n(r+1)+(r+m-2)} \cdot X_{n(r+1)+(r+m-3)} \cdots X_{n(r+1)+m}] \\ &\quad \cdot [Y_{n(r+1)+r+m-1} \cdot Y_{n(r+1)+r+m-2} \cdots Y_{n(r+1)+m+1}] \cdot Y_m \\ &= [X_{n(r+1)+r+m-1} \cdot X_{n(r+1)+(r+m-2)} \cdot X_{n(r+1)+(r+m-3)} \cdots X_{n(r+1)+m+1}] \\ &\quad \cdot [Y_{n(r+1)+r+m-1} \cdot Y_{n(r+1)+r+m-2} \cdots Y_{n(r+1)+m+1}] [X_{n(r+1)+m} \cdot Y_m] \end{aligned}$$

Using induction hypothesis,

$$\begin{aligned}
 &= [X_{n(r+1)+r+m-1} \cdot X_{n(r+1)+(r+m-2)} \cdot X_{n(r+1)+(r+m-3)} \cdots X_{n(r+1)+m+1}] \\
 &\cdot [Y_{n(r+1)+r+m-1} \cdot Y_{n(r+1)+r+m-2} \cdots Y_{n(r+1)+m+1}] [Y_{n(r+1)+m} \cdot X_m] \\
 &= [X_{n(r+1)+r+m-1} \cdot X_{n(r+1)+(r+m-2)} \cdot X_{n(r+1)+(r+m-3)} \cdots X_{n(r+1)+m+1}] [X_{n(r+1)+(r+m)} \cdot X_m] \\
 &= [Y_{n(r+1)+(r+m)+1} \cdot X_m] \\
 &= [Y_{(n+1)(r+1)+m} \cdot X_m]
 \end{aligned}$$

Hence result is true for $n + 1$. □

Theorem 3.6.

For every integer $n \geq 0, r \geq 0$

$$\begin{aligned}
 1. \quad \prod_{i=1}^{i=n} X_{ri+1} &= \prod_{i=1}^{i=rn} Y_i \\
 2. \quad \prod_{i=1}^{i=n} Y_{ri+1} &= \prod_{i=1}^{i=rn} X_i
 \end{aligned}$$

Proof. : If $n = 1$, then result is true because

$$X_{r+1} = Y_r \cdot Y_{r-1} \cdot Y_{r-2} \cdots Y_1$$

Assume that the result is true for some integer $n \geq 1$

Now,

$$\begin{aligned}
 X_{n+r} &= Y_{n+r-1} \cdot Y_{n+r-2} \cdot Y_{n+r-3} \cdots Y_n \\
 Y_{n+r} &= X_{n+r-1} \cdot X_{n+r-2} \cdot X_{n+r-3} \cdots X_n,
 \end{aligned}$$

Using Induction Method,

For, $n + 1$

$$\prod_{i=1}^{i=n+1} X_{ri+1} = \prod_{i=1}^{i=n} X_{ri+1} \cdot X_{r(n+1)+1}$$

Using induction hypothesis,

$$\begin{aligned}
 &= \prod_{i=1}^{i=rn} Y_i \cdot X_{rn+r+1} \\
 &= \prod_{i=1}^{i=rn} Y_i \cdot Y_{rn+r} \cdot Y_{rn+r-1} \cdot Y_{rn+r-2} \cdots Y_{rn+1} \\
 &= \prod_{i=1}^{i=rn+r} Y_i
 \end{aligned}$$

Hence result is true for $n + 1$. □

In the similar way (2) can be proved using induction method.

4. Conclusions

The identities of generalized multiplicative coupled Fibonacci sequence of r^{th} order are described here, the ideas can be extended for generalized multiplicative coupled Fibonacci sequence of r^{th} order with negative integers.

5. Acknowledgments

Authors are thankful to reviewers for their valuable comments in improvement of the paper.

References

- [1] K. Atanassov,L. Atanassov,D. Sasselov A New Perspective to the Generalization of the Fibonacci Sequence,*The Fibonacci Quarterly* 23(1) (1985) 21-28.
- [2] Koshy T.,Fibonacci and Lucas numbers with applications, Wiley-Intersection Pub.,2001.
- [3] K. T. Atanassov Remark on a New Direction of Generalized Fibonacci Sequence, *The Fibonacci Quarterly* 33(3) (1995) 249-250.
- [4] A. F. Horadam. A Generalized Fibonacci Sequence, *American Mathematical Monthly* 68(5) (1961) 455-459.
- [5] M. Singh , O. Shikhwal, S. Jain Coupled Fibonacci Sequences of Fifth order and Some Identities, *International Journal in Mathematical Analysis* 4(25) (2010) 1247-1254.
- [6] A. D. Godase, M. B. Dhakne, Fundamental Properties of Mutiplicative Coupled Fibonacci Sequences of Fourth Order Under Two Specific Schemes, *International Journal of Mathematical Archive* 4(6) (2013) 74-81.
- [7] A. D. Godase., M. B. Dhakne, Recurrent Formulas of The Generalized Fibonacci Sequences of Fifth Order, *International Journal of Mathematical Archive* 4(6) (2013) 61-67
- [8] A. D. Godase, M. B. Dhakne, On the properties of k-Fibonacci and k-Lucas numbers, *International Journal of Advances in Applied Mathematics and Mechanics* 2 (2014) 100-106.
- [9] Sangeeta Garg, On a new sequence of q-Baskakov-Szasz-Stancu operators, *International Journal of Advances in Applied Mathematics and Mechanics* 1(4) (2014) 74-82.
- [10] Animesh Gupta, Fixed point for generalized contractions on G- metric spaces, *International Journal of Advances in Applied Mathematics and Mechanics* 2(2) (2014) 39-51.