

Magnetic Tunnel Junctions

For the last two decades, magnetic tunnel junctions (MTJs) have played a central role in spintronic devices such as read heads of hard disk drives and nonvolatile magnetoresistive random access memories (MRAMs), and researchers are constantly working to improve their performance. [24]

*Magnetic materials are a vital ingredient in the components that store information in computers and mobile phones. Now, A*STAR researchers have developed a material that could help these magnetic-based memory devices to store and retrieve data faster while using less power. [23]*

A team of researchers with members from institutions in Germany and Israel has developed a way to launch plasmons with controlled amounts of angular momentum using spiral-like structures fashioned into a smooth layer of gold plate. [22]

Work at the New York Genome Centre represents a big step towards DNA-based information storage. Andrew Masterson reports. [21]

At Caltech, a group of researchers led by Assistant Professor of Bioengineering Lulu Qian is working to create circuits using not the usual silicon transistors but strands of DNA. [20]

Researchers have introduced a new type of "super-resolution" microscopy and used it to discover the precise walking mechanism behind tiny structures made of DNA that could find biomedical and industrial applications. [19]

Genes tell cells what to do—for example, when to repair DNA mistakes or when to die—and can be turned on or off like a light switch. Knowing which genes are switched on, or expressed, is important for the treatment and monitoring of disease. Now, for the first time, Caltech scientists have developed a simple way to visualize gene expression in cells deep inside the body using a common imaging technology. [18]

Researchers at The University of Manchester have discovered that a potential new drug reduces the number of brain cells destroyed by stroke and then helps to repair the damage. [17]

Researchers at the University of Connecticut have uncovered new information about how particles behave in our bloodstream, an important advancement that could help pharmaceutical scientists develop more effective cancer drugs. [16]

For the past 15 years, the big data techniques pioneered by NASA's Jet Propulsion Laboratory in Pasadena, California, have been revolutionizing biomedical research. On Sept. 6, 2016, JPL and the National Cancer Institute (NCI), part of the National Institutes of Health, renewed a research partnership through 2021, extending the development of data science that originated in space exploration and is now supporting new cancer discoveries. [15]

IBM scientists have developed a new lab-on-a-chip technology that can, for the first time, separate biological particles at the nanoscale and could enable physicians to detect diseases such as cancer before symptoms appear. [14]

Scientists work toward storing digital information in DNA. [13]

Leiden theoretical physicists have proven that DNA mechanics, in addition to genetic information in DNA, determines who we are. Helmut Schiessel and his group simulated many DNA sequences and found a correlation between mechanical cues and the way DNA is folded. They have published their results in PLoS One. [12]

We model the electron clouds of nucleic acids in DNA as a chain of coupled quantum harmonic oscillators with dipole-dipole interaction between nearest neighbours resulting in a van der Waals type bonding. [11]

Scientists have discovered a secret second code hiding within DNA which instructs cells on how genes are controlled. The amazing discovery is expected to open new doors to the diagnosis and treatment of diseases, according to a new study. [10]

There is also connection between statistical physics and evolutionary biology, since the arrow of time is working in the biological evolution also.

From the standpoint of physics, there is one essential difference between living things and inanimate clumps of carbon atoms: The former tend to be much better at capturing energy from their environment and dissipating that energy as heat. [8]

This paper contains the review of quantum entanglement investigations in living systems, and in the quantum mechanically modeled photoactive prebiotic kernel systems. [7]

The human body is a constant flux of thousands of chemical/biological interactions and processes connecting molecules, cells, organs, and fluids, throughout the brain, body, and nervous system. Up until recently it was thought that all these interactions operated in a linear sequence, passing on information much like a runner passing the baton to the next runner. However,

the latest findings in quantum biology and biophysics have discovered that there is in fact a tremendous degree of coherence within all living systems.

The accelerating electrons explain not only the Maxwell Equations and the Special Relativity, but the Heisenberg Uncertainty Relation, the Wave-Particle Duality and the electron's spin also, building the Bridge between the Classical and Quantum Theories.

The Planck Distribution Law of the electromagnetic oscillators explains the electron/proton mass rate and the Weak and Strong Interactions by the diffraction patterns. The Weak Interaction changes the diffraction patterns by moving the electric charge from one side to the other side of the diffraction pattern, which violates the CP and Time reversal symmetry.

The diffraction patterns and the locality of the self-maintaining electromagnetic potential explains also the Quantum Entanglement, giving it as a natural part of the Relativistic Quantum Theory and making possible to understand the Quantum Biology.

Contents

Preface.....	5
Spintronic technology advances with newly designed magnetic tunnel junctions	5
A novel 'soft' magnetic material could enable faster computer memory	7
Researchers launch plasmons with controlled amounts of angular momentum	7
One day, more than your genes will be stored in DNA	8
Computing with biochemical circuits made easy	9
Super-resolution system reveals mechanics of tiny 'DNA walker'	11
Visualizing gene expression with MRI.....	12
New drug limits and then repairs brain damage in stroke.....	13
When push comes to shove: Size matters for particles in our bloodstream	14
Fighting cancer with space research	16
IBM lab-on-a-chip breakthrough aims to help physicians detect cancer	18
Sorting bioparticles at the nanoscale.....	19
Scientists work toward storing digital information in DNA.....	19
Second layer of information in DNA confirmed	22
Mechanical cues	23
Simulation.....	23
Quantum entanglement between the electron clouds of nucleic acids in DNA.....	23
Scientists discover secret code hidden within human DNA	24

This Physicist Has a Groundbreaking Idea about Why Life Exists.....	25
Photoactive Prebiotic Systems	28
Significance Statement	28
Figure legend	29
Quantum Biology.....	30
Quantum Consciousness	30
Information – Entropy Theory of Physics	31
Information – Entropy Theory of Life	31
Creating quantum technology.....	32
Quantum Entanglement.....	32
The Bridge	32
Accelerating charges	32
Relativistic effect	33
Heisenberg Uncertainty Relation.....	33
Wave – Particle Duality	33
Atomic model	33
The Relativistic Bridge.....	33
The weak interaction	34
The General Weak Interaction.....	35
Fermions and Bosons.....	35
Van Der Waals force	35
Electromagnetic inertia and mass.....	36
Electromagnetic Induction	36
Relativistic change of mass.....	36
The frequency dependence of mass	36
Electron – Proton mass rate	36
Gravity from the point of view of quantum physics	36
The Gravitational force	36
The Higgs boson	37
Higgs mechanism and Quantum Gravity.....	38
What is the Spin?	38
The Graviton	38
Conclusions	39
References	39

Author: George Rajna

Preface

Scientists have discovered a secret second code hiding within DNA which instructs cells on how genes are controlled. The amazing discovery is expected to open new doors to the diagnosis and treatment of diseases, according to a new study. Ever since the genetic code was deciphered over 40 years ago, scientists have believed that it only described how proteins are made. However, the revelation made by the research team led by John Stamatoyannopoulos of the University of Washington indicates that genomes use the genetic code to write two separate languages. [10]

Jeremy England, a 31-year-old assistant professor at the Massachusetts Institute of Technology, has derived a mathematical formula that he believes explains this capacity. The formula, based on established physics, indicates that when a group of atoms is driven by an external source of energy (like the sun or chemical fuel) and surrounded by a heat bath (like the ocean or atmosphere), it will often gradually restructure itself in order to dissipate increasingly more energy. This could mean that under certain conditions, matter inexorably acquires the key physical attribute associated with life. [8]

We define our modeled self-assembled supramolecular photoactive centers, composed of one or more sensitizer molecules, precursors of fatty acids and a number of water molecules, as a photoactive prebiotic kernel system. [7]

The human body is a constant flux of thousands of chemical/biological interactions and processes connecting molecules, cells, organs, and fluids, throughout the brain, body, and nervous system. Up until recently it was thought that all these interactions operated in a linear sequence, passing on information much like a runner passing the baton to the next runner. However, the latest findings in quantum biology and biophysics have discovered that there is in fact a tremendous degree of coherence within all living systems. [5]

Quantum entanglement is a physical phenomenon that occurs when pairs or groups of particles are generated or interact in ways such that the quantum state of each particle cannot be described independently – instead, a quantum state may be given for the system as a whole. [4]

I think that we have a simple bridge between the classical and quantum mechanics by understanding the Heisenberg Uncertainty Relations. It makes clear that the particles are not point like but have a dx and dp uncertainty.

Spintronic technology advances with newly designed magnetic tunnel junctions

For the last two decades, magnetic tunnel junctions (MTJs) have played a central role in spintronic devices such as read heads of hard disk drives and nonvolatile magnetoresistive random access memories (MRAMs), and researchers are constantly working to improve their performance. One of

the most prominent achievements that accelerated the technology's practical applications was the realization of giant tunnel magnetoresistance (TMR) ratios by using rock-salt type MgO crystalline barrier. Now, in an article appearing in this week's issue of Applied Physics Letters, a Japanese team of researchers has succeeded in applying MgGa₂O₄ to a tunnel barrier, the core part of an MTJ, as an alternative material to more conventional insulators such as MgO and MgAl₂O₄.

An MTJ has a laminated structure consisting of a nanoscale insulating layer, called a tunnel barrier, sandwiched between two magnetic layers. One of the most important performance indexes of an MTJ is the tunnel magnetoresistance ratio (TMR ratio), the magnitude of resistance change. Magnesium oxide (MgO) is commonly used as a tunnel barrier since a large TMR ratio can be easily obtained.

"In order to widen the application range of MTJs further, we wanted to greatly tune the MTJ properties by replacing the tunnel barrier material," said Hiroaki Sukegawa, a scientist at the National Institute for Materials Science in Japan. "Particularly, for many MTJ applications, we need to have a large TMR ratio and low device resistance and for that we chose a tunnel barrier material with a low band gap."

The team selected semiconducting MgGa₂O₄, which has a band gap much lower than the conventional MgO insulator, and used existing technology to make an ultrathin MgAl₂O₄ layer to achieve the parameters they were looking for.

The biggest challenge was obtaining a high quality MgGa₂O₄ layer with defect-free interfaces since that is essential to achieving a large TMR ratio.

"We first attempted an oxidation method using an Mg-Ga alloy layer for the MgGa₂O₄ layer preparation however, this process also caused significant oxidation on the surface of the magnetic layer under the Mg-Ga, and the resulting fabricated structure did not function as an MTJ device," Sukegawa said. Inspired by their recent work on a high quality MgAl₂O₄ fabrication, the team then tried a direct sputtering method; the MgGa₂O₄ layer was formed by radio frequency sputtering from a high-density MgGa₂O₄ sintered target to reduce the interfacial over-oxidation.

This new method was very effective in producing a high quality MgGa₂O₄ tunnel barrier with extremely sharp and defect-free interfaces. It was a pleasant and unexpected surprise.

"We did not expect that we could construct an MTJ showing a large TMR ratio using MgGa₂O₄ in such a short time since there were few tunnel barrier materials capable of providing the large TMR ratio at room temperature that we were looking for," Sukegawa said..

This work demonstrates that, contrary to past understanding, MTJ tunnel barriers can be "designed." It was believed that tuning the physical parameters of the tunnel barrier while maintaining large TMR ratios was nearly impossible. These results strongly indicate that various physical properties of the tunnel barrier can be designed by selecting the composition of spinel based barrier materials as necessary while achieving efficient spin-dependent transport (i.e. large TMR ratio).

While there is still more work to be done to achieve larger TMR ratios, these results open the possibility of using "tunnel barrier design" with various spinel oxides to create new spintronic applications. [24]

A novel 'soft' magnetic material could enable faster computer memory

Magnetic materials are a vital ingredient in the components that store information in computers and mobile phones. Now, A*STAR researchers have developed a material that could help these magnetic-based memory devices to store and retrieve data faster while using less power.

Memory devices work when a small magnetic field is applied to the storage medium to align atomic-level magnets known as spins. This spin alignment, or magnetization, in one region of the magnetic material can represent one 'bit' of information, which can be 'read' back again using a magnet. Scientists are trying to improve the performance of magnetic memories by reducing both the energy required to change the magnetization and unwanted noise.

One approach is to use a magnetic material with a property known as negative magnetocrystalline anisotropy. This means that less energy is required to align the spins in one direction than another, and so the material is generally easier to magnetize and demagnetize. This low coercivity is useful because this so-called 'soft' magnetic material can guide a magnetic field onto the storage layer, thus lowering the intensity of field that must be applied to alter the 'hard' material's magnetization.

Tiejun Zhou and co-workers from the A*STAR Data Storage Institute found a way to further reduce the coercivity of a soft material called cobalt iridium by adding rhodium.

The team created their magnetic material with a technique known as direct current magnetron sputtering. Cobalt, iridium and rhodium were simultaneously ejected from separate solid sources in a vacuum chamber and deposited on a silicon substrate. By changing the power supplied to each of the sources, the researchers could control the composition of the final material, increasing the amount of rhodium at the expense of iridium. Measurements of the magnetic properties of CoIr-Rh films demonstrated that the introduction of this rhodium reduced the coercivity and the damping constant by more than a half of that of unmodified cobalt iridium.

"When used in a device, such negative magnetocrystalline anisotropy materials enable higher frequency operation at lower driving current and the creation of a higher in-plane alternating-current magnetic field for effective assisted switching, and higher stability against stray fields and temperature fluctuations," explains Zhou. The team demonstrated this improved performance in a memory device called a spin torque oscillator.

The results show that CoIr-Rh could help to develop commercial low-energy magnetic storage. "By fine tuning the composition, we can continuously improve the properties of magnetic materials to meet the criteria required for industry-level applications," says Zhou. [23]

Researchers launch plasmons with controlled amounts of angular momentum

A team of researchers with members from institutions in Germany and Israel has developed a way to launch plasmons with controlled amounts of angular momentum using spiral-like structures fashioned into a smooth layer of gold plate. In their paper published in the journal *Science*, the

group describes their technique and why they believe it could one day provide the basis for a new type of storage device.

Prior work had shown that it was possible to use orbital angular momentum of photons as a way to store data in the air (because of their different phase advances—each could be used to represent an individual character), which in practical use would be optical fibers. In this new effort, the researchers have found a way to do much the same thing, only on a chip, which allows for it to be confined, making it much easier to port to an actual application.

To accomplish this feat, the researchers turned to plasmonics, which are oscillations in quantum particles or quasiparticles. In this case, it was the oscillation of electrons that occurs when photons strike a metal surface. The idea, the researchers noted, was that plasmons allow for confining light, which meant that it was possible to make its wavelength conform to a metal surface. They found they were able to kick off plasmon waves with angular momentum by etching spirals into a gold plate—that allowed for monitoring its motion using an electron microscope.

Using it, they were able to see that firing photons at the spiral in the plates caused electrons to be ejected, which was an indirect way to "see" plasmons, which were controlled by the angular momentum of photons. This meant it was possible to use the initial wave advances to represent characters within the confines of a computer chip. That day is still a long way off, of course—the work was more a proof of concept than a finished product. But if a way can be found to mass produce chips using the technique, end users could see a huge increase in the amount of data storage. The work also suggests, the team notes, that it might be possible to do some things with light that have never been possible before. [22]

One day, more than your genes will be stored in DNA

As anyone old enough to remember video tapes will testify, the problem with information storage is that after a while the machines needed to retrieve the data no longer exist.

A pair of New York scientists, however, are working to resolve the issue by adapting a storage medium that has been around for billions of years: DNA.

DNA is, of course, an exceptionally versatile and reliable information storage system. Using just four nucleobases – cytosine, guanine, adenine and thymine – it effectively encapsulates, and replicates, the building and operating instructions for every living thing on earth.

A complete set on chromosomes in a single human cell has been estimated to contain 1.5 Gigabytes worth of data. That data is replicated in every non-sex cell in the body, but if it wasn't – if every cell was able to encode different information – the average human's total DNA would easily store 150 trillion Gigabytes.

Scientists Yaniv Erlich and Dina Zielinski of the New York Genome Centre have picked up on earlier experimental work on the use of artificially generated DNA as a storage mechanism. Using a coding strategy they dub the DNA Fountain they report successfully using genetic material to store and retrieve a full computer operating system, a movie, and some other files.

In a paper lodged on the pre-print site biorxiv, research awaiting peer review, Erlich and Zielinski note that two bits of information is the maximum storage capacity for a single nucleobase and its support molecules – collectively called a nucleotide. However, once a few biochemical constraints and the need to free up a bit of space to allow indexing are taken into account the maximum storage achieved turned out to be just over 1.8 bits.

The DNA Fountain works by breaking down binary code – the code that stores data on computers and digital discs – into short lengths, and then transfers the information to DNA nucleotides using a computer program that is already widely used to prevent drop-outs in mobile phone and broadcast environments.

The information, once ensconced in the lab-generated DNA, was easily retrieved using existing gene-sequencing technology, after which it was transformed back into usable binary. Erlich and Zielinski successfully recovered all of their data without a single error.

Although DNA has a half-life of some 500-odd years, it still represents a much more robust and archival storage medium than anything else so far invented. (Think about scratched vinyl albums, and now-useless Jaz drive cartridges, for instance.) Furthermore, the scientists note, the storage capacity of DNA is phenomenal. One gram of the stuff could easily store 215 Petabytes of data. A single Petabyte comprises 1,000,000,000,000 bytes. That's a lot of movies. [21]

Computing with biochemical circuits made easy

Electronic circuits are found in almost everything from smartphones to spacecraft and are useful in a variety of computational problems from simple addition to determining the trajectories of interplanetary satellites. At Caltech, a group of researchers led by Assistant Professor of Bioengineering Lulu Qian is working to create circuits using not the usual silicon transistors but strands of DNA.

The Qian group has made the technology of DNA circuits accessible to even novice researchers—including undergraduate students—using a software tool they developed called the Seesaw Compiler. Now, they have experimentally demonstrated that the tool can be used to quickly design DNA circuits that can then be built out of cheap "unpurified" DNA strands, following a systematic wet-lab procedure devised by Qian and colleagues.

A paper describing the work appears in the February 23 issue of Nature Communications.

Although DNA is best known as the molecule that encodes the genetic information of living things, they are also useful chemical building blocks. This is because the smaller molecules that make up a strand of DNA, called nucleotides, bind together only with very specific rules—an A nucleotide binds to a T, and a C nucleotide binds to a G. A strand of DNA is a sequence of nucleotides and can become a double strand if it binds with a sequence of complementary nucleotides.

DNA circuits are good at collecting information within a biochemical environment, processing the information locally and controlling the behavior of individual molecules. Circuits built out of DNA strands instead of silicon transistors can be used in completely different ways than electronic circuits. "A DNA circuit could add 'smarts' to chemicals, medicines, or materials by making their

functions responsive to the changes in their environments," Qian says. "Importantly, these adaptive functions can be programmed by humans."

To build a DNA circuit that can, for example, compute the square root of a number between 0 and 16, researchers first have to carefully design a mixture of single and partially double-stranded DNA that can chemically recognize a set of DNA strands whose concentrations represent the value of the original number. Mixing these together triggers a cascade of zipping and unzipping reactions, each reaction releasing a specific DNA strand upon binding. Once the reactions are complete, the identities of the resulting DNA strands reveal the answer to the problem.

With the Seesaw Compiler, a researcher could tell a computer the desired function to be calculated and the computer would design the DNA sequences and mixtures needed. However, it was not clear how well these automatically designed DNA sequences and mixtures would work for building DNA circuits with new functions; for example, computing the rules that govern how a cell evolves by sensing neighboring cells.

"Constructing a circuit made of DNA has thus far been difficult for those who are not in this research area, because every circuit with a new function requires DNA strands with new sequences and there are no off-the-shelf DNA circuit components that can be purchased," says Chris Thachuk, senior postdoctoral scholar in computing and mathematical sciences and second author on the paper. "Our circuit-design software is a step toward enabling researchers to just type in what they want to do or compute and having the software figure out all the DNA strands needed to perform the computation, together with simulations to predict the DNA circuit's behavior in a test tube. Even though these DNA strands are still not off-the-shelf products, we have now shown that they do work well for new circuits with user-designed functions."

"In the 1950s, only a few research labs that understood the physics of transistors could build early versions of electronic circuits and control their functions," says Qian. "But today many software tools are available that use simple and human-friendly languages to design complex electronic circuits embedded in smart machines. Our software is kind of like that: it translates simple and human-friendly descriptions of computation to the design of complex DNA circuits."

The Seesaw Compiler was put to the test in 2015 in a unique course at Caltech, taught by Qian and called "Design and Construction of Programmable Molecular Systems" (BE/CS 196 ab). "How do you evaluate the accessibility of a new technology? You give the technology to someone who is intellectually capable but has minimal prior background," Qian says.

"The students in this class were undergrads and first-year graduate students majoring in computer science and bioengineering," says Anupama Thubagere, a graduate student in biology and bioengineering and first author on the paper. "I started working with them as a head teaching assistant and together we soon discovered that using the Seesaw Compiler to design a DNA circuit was easy for everyone."

However, building the designed circuit in the wet lab was not so simple. Thus, with continued efforts after the class, the group set out to develop a systematic wet-lab procedure that could guide researchers—even novices like undergraduate students—through the process of building DNA

circuits. "Fortunately, we found a general solution to every challenge that we encountered, now making it easy for everyone to build their own DNA circuits," Thubagere says.

The group showed that it was possible to use cheap, "unpurified" DNA strands in these circuits using the new process. This was only possible because steps in the systematic wet-lab procedure were designed to compensate for the lower synthesis quality of the DNA strands.

"We hope that this work will convince more computer scientists and researchers from other fields to join our community in developing increasingly powerful molecular machines and to explore a much wider range of applications that will eventually lead to the transformation of technology that has been promised by the invention of molecular computers," Qian says.

The paper is titled, "Compiler-aided systematic construction of large-scale DNA strand displacement circuits using unpurified components." [20]

Super-resolution system reveals mechanics of tiny 'DNA walker'

Researchers have introduced a new type of "super-resolution" microscopy and used it to discover the precise walking mechanism behind tiny structures made of DNA that could find biomedical and industrial applications.

The researchers also demonstrated how the "DNA walker" is able to release an anticancer drug, representing a potential new biomedical technology, said Jong Hyun Choi, an associate professor of mechanical engineering at Purdue University.

Synthetic nanomotors and walkers are intricately designed systems that draw chemical energy from the environment and convert it into mechanical motion. However, because they are too small to be observed using conventional light microscopes, researchers have been unable to learn the precise steps involved in the walking mechanisms, knowledge essential to perfecting the technology.

"If you cannot resolve or monitor these walkers in action, you will be unable to understand their mechanical operation," Choi said.

He led a Purdue team that has solved this problem by developing a super-resolution microscopy system designed to study the DNA walkers. The new findings appeared in the journal *Science Advances* on Jan. 20.

Researchers around the world are creating synthetic motors based on DNA and RNA, the genetic materials in cells that consist of a sequence of four chemical bases: adenine, guanine, cytosine and thymine. The designs are inspired by natural biological motors that have evolved to perform specific tasks critical to the function of cells.

The Purdue researchers have designed a DNA walking system consisting of an enzymatic core and two arms. The walker travels along a carbon-nanotube track "decorated" with strands of RNA. The enzymatic core cleaves off segments of these RNA strands as the walker continuously moves forward, binding to and harvesting energy from the RNA. The walker moves in a six-step cycle that repeats as long as there is RNA fuel.

A fluorescent nanoparticle is attached to one arm of the DNA walker, causing it to glow when exposed to light in the visible part of the spectrum. The carbon-nanotube track also fluoresces when exposed to light in a portion of the near-infrared spectrum. Because the new super-resolution microscopy system operates in both the visible and near-infrared spectra, it is possible to track the walking mechanism.

The super-resolution technology allows researchers to resolve structural features far smaller than the wavelength of visible light, which is normally difficult using conventional microscopes because of the Abbe diffraction limit, established by physicist Ernst Abbe in 1873. The limit is about 250 nanometers, which is large compared to the tiny walkers, measuring about 5 nanometers long.

As the DNA walker is exposed to laser light, the nanoparticle and nanotube flash on and off randomly. These flashes are captured as numerous fluorescing dots in thousands of imaging frames. This collection of points is then used to reconstruct the precise motion of the walker, which moves in a six-step cycle that involves cleaving portions of the RNA strand and harvesting its energy before moving on to the next strand.

Findings revealed three primary steps dominate this walking mechanism.

"So, if you can control these three steps within this walking cycle then you can really study and better control these walkers," Choi said. "You can speed them up, you can make them stop and move in different directions."

Whereas it previously would have taken 20 hours or longer to study a complete walking cycle, the new approach speeds the process to roughly one minute. [19]

Visualizing gene expression with MRI

Genes tell cells what to do—for example, when to repair DNA mistakes or when to die—and can be turned on or off like a light switch. Knowing which genes are switched on, or expressed, is important for the treatment and monitoring of disease. Now, for the first time, Caltech scientists have developed a simple way to visualize gene expression in cells deep inside the body using a common imaging technology.

Researchers in the laboratory of Mikhail Shapiro, assistant professor of chemical engineering and Heritage Medical Research Institute Investigator, have invented a new method to link magnetic resonance imaging (MRI) signals to gene expression in cells—including tumor cells—in living tissues. The technique, which eventually could be used in humans, would allow gene expression to be monitored non-invasively, requiring no surgical procedures such as biopsies.

The work appears in the December 23 online edition of the journal *Nature Communications*.

In MRI, hydrogen atoms in the body—atoms that are mostly contained in water molecules and fat—are excited using a magnetic field. The excited atoms, in turn, emit signals that can be used to create images of the brain, muscle, and other tissues, which can be distinguished based on the local physical and chemical environment of the water molecules. While this technique is widely used, it usually provides only anatomical snapshots of tissues or physiological functions such as blood flow rather than observations of the activity of specific cells.

"We thought that if we could link signals from water molecules to the expression of genes of interest, we could change the way the cell looks under MRI," says Arnab Mukherjee, a postdoctoral scholar in chemical engineering at Caltech and co-lead author on the paper.

The group turned to a protein that naturally occurs in humans, called aquaporin. Aquaporin sits within the membrane that envelops cells and acts as a gatekeeper for water molecules, allowing them to move in and out of the cell. Shapiro's team realized that increasing the number of aquaporins on a given cell made it stand out in MRI images acquired using a common clinical technique called diffusion-weighted imaging, which is sensitive to the movement of water molecules. They then linked aquaporin to genes of interest, making it what scientists call a reporter gene. This means that when a gene of interest is turned on, the cell will overexpress aquaporin, making the cell look darker under diffusion-weighted MRI.

The researchers showed that this technique was successful in monitoring gene expression in a brain tumor in mice. After implanting the tumor, they gave the mice a drug to trigger the tumor cells to express the aquaporin reporter gene, which made the tumor look darker in MRI images.

"Overexpression of aquaporin has no negative impact on cells because it is exclusive to water and simply allows the molecules to go back and forth across the cell membrane," Shapiro says. Under normal physiological conditions the number of water molecules entering and exiting an aquaporin-expressing cell is the same, so that the total amount of water in each cell does not change. "Aquaporin is a very convenient way to genetically change the way that cells look under MRI."

Though the work was done in mice, it has the potential for clinical translation, according to Shapiro. Aquaporin is a naturally occurring gene and will not cause an immune reaction. Previously developed reporter genes for MRI have been much more limited in their capabilities, requiring the use of specific metals that are not always available in some tissues.

"An effective reporter gene for MRI is a 'holy grail' in biomedical imaging because it would allow cellular function to be observed non-invasively," says Shapiro. "Aquaporins are a new way to think about this problem. It is remarkable that simply allowing water molecules to more easily get into and out of cells in a tissue gives us the ability to remotely see those cells in the middle of the body."

The paper is titled "Non-invasive imaging using reporter genes altering cellular water permeability." [18]

New drug limits and then repairs brain damage in stroke

Researchers at The University of Manchester have discovered that a potential new drug reduces the number of brain cells destroyed by stroke and then helps to repair the damage.

A reduction in blood flow to the brain caused by stroke is a major cause of death and disability, and there are few effective treatments.

A team of scientists at The University of Manchester has now found that a potential new stroke drug not only works in rodents by limiting the death of existing brain cells but also by promoting the birth of new neurones (so-called neurogenesis).

This finding provides further support for the development of this anti-inflammatory drug, interleukin-1 receptor antagonist (IL-1Ra in short), as a new treatment for stroke. The drug is already licensed for use in humans for some conditions, including rheumatoid arthritis. Several early stage clinical trials in stroke with IL-1Ra have already been completed in Manchester, though it is not yet licensed for this condition.

In the research, published in the biomedical journal *Brain, Behavior and Immunity*, the researchers show that in rodents with a stroke there is not only reduced brain damage early on after the stroke, but several days later increased numbers of new neurones, when treated with the anti-inflammatory drug IL-1Ra.

Previous attempts to find a drug to prevent brain damage after stroke have proved unsuccessful and this new research offers the possibility of a new treatment.

Importantly, the use of IL-1Ra might be better than other failed drugs in stroke as it not only limits the initial damage to brain cells, but also helps the brain repair itself long-term through the generation of new brain cells.

These new cells are thought to help restore function to areas of the brain damaged by the stroke. Earlier work by the same group showed that treatment with IL-1Ra does indeed help rodents regain motor skills that were initially lost after a stroke. Early stage clinical trials in stroke patients also suggest that IL-1Ra could be beneficial.

The current research is led by Professor Stuart Allan, who commented: "The results lend further strong support to the use of IL-1Ra in the treatment of stroke, however further large trials are necessary."

The paper, 'Reparative effects of interleukin-1 receptor antagonist in young and aged/co-morbid rodents after cerebral ischemia', was published in the journal *Brain, Behavior and Immunity*. [17]

When push comes to shove: Size matters for particles in our bloodstream

Researchers at the University of Connecticut have uncovered new information about how particles behave in our bloodstream, an important advancement that could help pharmaceutical scientists develop more effective cancer drugs.

Making sure cancer medications reach the leaky blood vessels surrounding most tumor sites is one of the critical aspects of treatment and drug delivery. While surface chemistry, molecular interactions, and other factors come into play once drug-carrying particles arrive at a tumor, therapeutic medication doesn't do very much good if it never reaches its intended target.

Anson Ma, an assistant professor of chemical and biomolecular engineering at UConn, used a microfluidic channel device to observe, track, and measure how individual particles behaved in a simulated blood vessel.

The research team's goal: to learn more about the physics influencing a particle's behavior as it travels in our blood and to determine which particle size might be the most effective for delivering

drugs to their targets. The team's experimental findings mark the first time such quantitative data has been gathered. The study was published Oct. 4 in the Biophysical Journal.

"Even before particles reach a target site, you have to worry about what is going to happen with them after they get injected into the bloodstream," Ma says. "Are they going to clump together? How are they going to move around? Are they going to get swept away and flushed out of our bodies?"

Using a high-powered fluorescence microscope in UConn's Complex Fluids Lab, Ma was able to observe particles being carried along in the simulated blood vessel in what could be described as a vascular Running of the Bulls. Red blood cells race through the middle of the channel as the particles - highlighted under the fluorescent light - get carried along in the rush, bumping and bouncing off the blood cells until they are pushed to open spaces - called the cell-free layer - along the vessel's walls.

What Ma found was that larger particles - the optimum size appeared to be about 2 microns - were most likely to get pushed to the cell-free layer where their chances of carrying medication into a tumor site are greatest. The research team also determined that 2 microns was the largest size that should be used if particles are going to have any chance of going through the leaky blood vessel walls into the tumor site.

"When it comes to using particles for the delivery of cancer drugs, size matters," Ma says. "When you have a bigger particle, the chance of it bumping into blood cells is much higher, there are a lot more collisions, and they tend to get pushed to the blood vessel walls."

The results were somewhat surprising. In preparing their hypothesis, the research team estimated that smaller particles were probably the most effective since they would move the most in collisions with blood cells, much like what happens when a small ball bounces off a larger one. But just the opposite proved true. The smaller particles appeared to skirt through the mass of moving blood cells and were less likely to experience the "trampoline" effect and get bounced to the cell-free layer, says Ma.

The research was funded by the National Science Foundation's Early-concept Grants for Exploratory Research or EAGER program, which supports exploratory work in its early stages on untested, but potentially transformative, research ideas or approaches.

Knowing how particles behave in our circulatory system should help improve targeted drug delivery, Ma says, which in turn will further reduce the toxic side effects caused by potent cancer drugs missing their target and impacting the body's healthy tissue.

Measuring how different sized particles move in the bloodstream may also be beneficial in bioimaging, where scientists and doctors want to keep particles circulating in the bloodstream long enough for imaging to occur. In that case, smaller particles would be better, says Ma.

Moving forward, Ma would like to explore other aspects of particle flow in our circulatory system such as how particles behave when they pass through a constricted area, such as from a blood vessel to a capillary. Capillaries are only about 7 microns in diameter. The average human hair is 100

microns. Ma says he would like to know how that constricted space might impact particle flow or the ability of particles to accumulate near the vessel walls.

"We have all of this complex geometry in our bodies," says Ma. "Most people just assume there is no impact when a particle moves from a bigger channel to a smaller channel because they haven't quantified it. Our plan is to do some experiments to look at this more carefully, building on the work that we just published." [16]

Fighting cancer with space research

Every day, NASA spacecraft beam down hundreds of petabytes of data, all of which has to be codified, stored and distributed to scientists across the globe. Increasingly, artificial intelligence is helping to "read" this data as well, highlighting similarities between datasets that scientists might miss.

For the past 15 years, the big data techniques pioneered by NASA's Jet Propulsion Laboratory in Pasadena, California, have been revolutionizing biomedical research. On Sept. 6, 2016, JPL and the National Cancer Institute (NCI), part of the National Institutes of Health, renewed a research partnership through 2021, extending the development of data science that originated in space exploration and is now supporting new cancer discoveries.

The NCI-supported Early Detection Research Network (EDRN) is a consortium of biomedical investigators who share anonymized data on cancer biomarkers, chemical or genetic signatures related to specific cancers. Their goal is to pool all their research data into a single, searchable network, with the goal of translating their collective work into techniques for early diagnosis of cancer or cancer risk.

In the time they've worked together, JPL and EDRN's efforts have led to the discovery of six new Food and Drug Administration-approved cancer biomarkers and nine biomarkers approved for use in Clinical Laboratory Improvement Amendments labs. The FDA has approved each of these biomarkers for use in cancer research and diagnosis. These agency-approved biomarkers have been used in more than 1 million patient diagnostic tests worldwide.

"After the founding of EDRN in 2000, the network needed expertise to take data from multiple studies on cancer biomarkers and create a single, searchable network of research findings for scientists," said Sudhir Srivastava, chief of NCI's Cancer Biomarkers Research Group and head of EDRN. JPL had decades of experience doing similar work for NASA, where spacecraft transmit hundreds of petabytes of data to be coded, stored and distributed to scientists across the globe.

Dan Crichton, the head of JPL's Center for Data Science and Technology, a joint initiative with Caltech in Pasadena, California, helped establish a JPL-based informatics center dedicated to supporting EDRN's big data efforts. In the renewed partnership, JPL is expanding its data science efforts to research and applying technologies for additional NCI-funded programs. Those programs include EDRN, the Consortium for Molecular and Cellular Characterization of Screen-Detected Lesions, and the Informatics Technology for Cancer Research initiative.

"From a NASA standpoint, there are significant opportunities to develop new data science capabilities that can support both the mission of exploring space and cancer research using common

methodological approaches," Crichton said. "We have a great opportunity to perfect those techniques and grow JPL's data science technologies, while serving our nation.

Crichton said JPL has led the way when it comes to taking data from raw observations to scientific conclusions. One example: JPL often deals with measurements from a variety of sensors—say, cameras and mass spectrometers. Both can be used to study a star, planet or similar target object. But it takes special software to recognize that readings from very different instruments relate to one another.

There's a similar problem in cancer research, where readings from different biomedical tests or instruments require correlation with one another. For that to happen, data have to be standardized, and algorithms must be "taught" to know what they're looking for.

Since the time of its founding, EDRN's major challenge has been access. Research centers all over the United States had large numbers of biomarker specimens, but each had its own way of labeling, storing and sharing their datasets. Ten sites may have high-quality specimens for study, but if their common data elements—age of patient, cancer type and other characteristics - aren't listed uniformly, they can't be studied as a whole.

"We didn't know if they were early-stage or late-stage specimens, or if any level of treatment had been tried," Srivastava said. "And JPL told us, 'We do this type of thing all the time! That's how we manage our Planetary Data System.'"

As the network has developed, it has added members from dozens of institutions, including Dartmouth College's Geisel School of Medicine; Harvard Medical School's Massachusetts General Hospital; Stanford's NIST Genome-Scale Measurements Group; University of Texas' MD Anderson Cancer Center; and numerous others.

Christos Patriotis, program director at NCI's Cancer Biomarkers Research Group, said the network's members now include international researchers from the U.K., China, Japan, Australia, Israel and Chile.

"The more we expand, the more data we integrate," Patriotis said. "Instead of being silos, now our partners can integrate their findings. Each system can speak to the others."

As JPL and NCI's collaboration advances, next steps include image recognition technology, such as helping EDRN archive images of cancer specimens. Those images could be analyzed by computer vision, which is currently used to spot similarities in star clusters and other astrophysics research.

In the near future, Crichton said, machine learning algorithms could compare a CT scan with an archive of similar images, searching for early signs of cancer based on a patient's age, ethnic background and other demographics.

"As we develop more automated methods for detecting and classifying features in images, we see great opportunities for enhancing data discovery," Crichton said.

"We have examples where algorithms for detection of features in astronomy images have been transferred to biology and vice-versa." [15]

IBM lab-on-a-chip breakthrough aims to help physicians detect cancer

IBM scientists have developed a new lab-on-a-chip technology that can, for the first time, separate biological particles at the nanoscale and could enable physicians to detect diseases such as cancer before symptoms appear.

As reported today in the journal *Nature Nanotechnology*, the IBM team's results show size-based separation of bioparticles down to 20 nanometers (nm) in diameter, a scale that gives access to important particles such as DNA, viruses and exosomes. Once separated, these particles can potentially be analyzed by physicians to reveal signs of disease even before patients experience any physical symptoms and when the outcome from treatment is most positive. Until now, the smallest bioparticle that could be separated by size with on-chip technologies was about 50 times or larger, for example, separation of circulating tumor cells from other biological components.

IBM is collaborating with a team from the Icahn School of Medicine at Mount Sinai to continue development of this lab-on-a-chip technology and plans to test it on prostate cancer, the most common cancer in men in the U.S.

In the era of precision medicine, exosomes are increasingly being viewed as useful biomarkers for the diagnosis and prognosis of malignant tumors. Exosomes are released in easily accessible bodily fluids such as blood, saliva or urine. They represent a precious biomedical tool as they can be used in the context of less invasive liquid biopsies to reveal the origin and nature of a cancer.

The IBM team targeted exosomes with their device as existing technologies face challenges for separating and purifying exosomes in liquid biopsies. Exosomes range in size from 20-140nm and contain information about the health of the originating cell that they are shed from. A determination of the size, surface proteins and nucleic acid cargo carried by exosomes can give essential information about the presence and state of developing cancer and other diseases.

IBM's results show they could separate and detect particles as small as 20 nm from smaller particles, that exosomes of size 100 nm and larger could be separated from smaller exosomes, and that separation can take place in spite of diffusion, a hallmark of particle dynamics at these small scales. With Mt. Sinai, the team plans to confirm their device is able to pick up exosomes with cancer-specific biomarkers from patient liquid biopsies.

"The ability to sort and enrich biomarkers at the nanoscale in chip-based technologies opens the door to understanding diseases such as cancer as well as viruses like the flu or Zika," said Gustavo Stolovitzky, Program Director of Translational Systems Biology and Nanobiotechnology at IBM Research. "Our lab-on-a-chip device could offer a simple, noninvasive and affordable option to potentially detect and monitor a disease even at its earliest stages, long before physical symptoms manifest. This extra amount of time allows physicians to make more informed decisions and when the prognosis for treatment options is most positive."

With the ability to sort bioparticles at the nanoscale, Mt. Sinai hopes that IBM's technology can provide a new method to eavesdrop on the messages carried by exosomes for cell-to-cell communications. This can elucidate important questions about the biology of diseases as well as pave the way to noninvasive and eventually affordable point-of-care diagnostic tools. Monitoring

this intercellular conversation more regularly could allow medical experts to track an individual's state of health or progression of a disease.

"When we are ahead of the disease we usually can address it well; but if the disease is ahead of us, the journey is usually much more difficult. One of the important developments that we are attempting in this collaboration is to have the basic grounds to identify exosome signatures that can be there very early on before symptoms appear or before a disease becomes worse," said Dr. Carlos Cordon-Cardo, Professor and Chairman for the Mount Sinai Health System Department of Pathology. "By bringing together Mount Sinai's domain expertise in cancer and pathology with IBM's systems biology experience and its latest nanoscale separation technology, the hope is to look for specific, sensitive biomarkers in exosomes that represent a new frontier to offering clues that might hold the answer to whether a person has cancer or how to treat it."

Sorting bioparticles at the nanoscale

Lab-on-a-chip technologies have become an incredibly helpful diagnostic tool for physicians as they can be significantly faster, portable, easy to use and require less sample volume to help detect diseases. The goal is to shrink down to a single silicon chip all of the processes necessary to analyze a disease that would normally be carried out in a full-scale biochemistry lab.

Using a technology called nanoscale deterministic lateral displacement, or nano-DLD, IBM scientists Dr. Joshua Smith and Dr. Benjamin Wunsch led development of a lab-on-a-chip technology that allows a liquid sample to be passed, in continuous flow, through a silicon chip containing an asymmetric pillar array. This array allows the system to sort a microscopic waterfall of nanoparticles, separating particles by size down to tens of nanometers resolution. IBM has already scaled down the chip size to 2cm by 2cm, while continuing development to increase the device density to improve functionality and throughput.

Much like how a road through a small tunnel only allows smaller cars to pass while forcing bigger trucks to detour around, nano-DLD uses a set of pillars to deflect larger particles while allowing smaller particles to flow through the gaps of the pillar array unabated, effectively separating this particle "traffic" by size while not disrupting flow. Interestingly, IBM scientists noticed that nano-DLD arrays can also split a mixture of many different particle sizes into a spread of streams, much like a prism splits white light into different colors. The continuous flow nature of this technology circumvents stop-and-go batch processing typical of conventional separation techniques.

Leveraging IBM's vast semiconductor expertise with its growing capabilities in experimental biology, IBM scientists used manufacturable silicon processes to produce the nano-DLD arrays for their lab-on-a-chip device. As part of its on-going strategy, IBM researchers are working to increase the diversity of bioparticles that can be separated with their device, and improving the precision and specificity for real-world clinical applications. [14]

Scientists work toward storing digital information in DNA

Her computer, Karin Strauss says, contains her "digital attic"—a place where she stores that published math paper she wrote in high school, and computer science schoolwork from college.

She'd like to preserve the stuff "as long as I live, at least," says Strauss, 37. But computers must be replaced every few years, and each time she must copy the information over, "which is a little bit of a headache."

It would be much better, she says, if she could store it in DNA—the stuff our genes are made of.

Strauss, who works at Microsoft Research in Redmond, Washington, is working to make that sci-fi fantasy a reality.

She and other scientists are not focused in finding ways to stow high school projects or snapshots or other things an average person might accumulate, at least for now. Rather, they aim to help companies and institutions archive huge amounts of data for decades or centuries, at a time when the world is generating digital data faster than it can store it.

To understand her quest, it helps to know how companies, governments and other institutions store data now: For long-term storage it's typically disks or a specialized kind of tape, wound up in cartridges about three inches on a side and less than an inch thick. A single cartridge containing about half a mile of tape can hold the equivalent of about 46 million books of 200 pages apiece, and three times that much if the data lends itself to being compressed.

A tape cartridge can store data for about 30 years under ideal conditions, says Matt Starr, chief technology officer of Spectra Logic, which sells data-storage devices. But a more practical limit is 10 to 15 years, he says.

It's not that the data will disappear from the tape. A bigger problem is familiar to anybody who has come across an old eight-track tape or floppy disk and realized he no longer has a machine to play it. Technology moves on, and data can't be retrieved if the means to read it is no longer available, Starr says.

So for that and other reasons, long-term archiving requires repeatedly copying the data to new technologies.

Into this world comes the notion of DNA storage. DNA is by its essence an information-storing molecule; the genes we pass from generation to generation transmit the blueprints for creating the human body. That information is stored in strings of what's often called the four-letter DNA code. That really refers to sequences of four building blocks—abbreviated as A, C, T and G—found in the DNA molecule. Specific sequences give the body directions for creating particular proteins.

Digital devices, on the other hand, store information in a two-letter code that produces strings of ones and zeroes. A capital "A," for example, is 01000001.

Converting digital information to DNA involves translating between the two codes. In one lab, for example, a capital A can become ATATG. The idea is once that transformation is made, strings of DNA can be custom-made to carry the new code, and hence the information that code contains.

One selling point is durability. Scientists can recover and read DNA sequences from fossils of Neanderthals and even older life forms. So as a storage medium, "it could last thousands and thousands of years," says Luis Ceze of the University of Washington, who works with Microsoft on DNA data storage.

Advocates also stress that DNA crams information into very little space. Almost every cell of your body carries about six feet of it; that adds up to billions of miles in a single person. In terms of information storage, that compactness could mean storing all the publicly accessible data on the internet in a space the size of a shoebox, Ceze says.

In fact, all the digital information in the world might be stored in a load of whitish, powdery DNA that fits in space the size of a large van, says Nick Goldman of the European Bioinformatics Institute in Hinxton, England.

What's more, advocates say, DNA storage would avoid the problem of having to repeatedly copy stored information into new formats as the technology for reading it becomes outmoded.

"There's always going to be someone in the business of making a DNA reader because of the health care applications," Goldman says. "It's always something we're going to want to do quickly and inexpensively."

Getting the information into DNA takes some doing. Once scientists have converted the digital code into the 4-letter DNA code, they have to custom-make DNA.

For some recent research Strauss and Ceze worked on, that involved creating about 10 million short strings of DNA.

Twist Bioscience of San Francisco used a machine to create the strings letter by letter, like snapping together Lego pieces to build a tower. The machine can build up to 1.6 million strings at a time.

Each string carried just a fragment of information from a digital file, plus a chemical tag to indicate what file the information came from.

To read a file, scientists use the tags to assemble the relevant strings. A standard lab machine can then reveal the sequence of DNA letters in each string.

Nobody is talking about replacing hard drives in consumer computers with DNA. For one thing, it takes too long to read the stored information. That's never going to be accomplished in seconds, says Ewan Birney, who works on DNA storage with Goldman at the bioinformatics institute.

But for valuable material like corporate records in long-term storage, "if it's worth it, you'll wait," says Goldman, who with Birney is talking to investors about setting up a company to offer DNA storage.

Sri Kosuri of the University of California Los Angeles, who has worked on DNA information storage but now largely moved on to other pursuits, says one challenge for making the technology practical is making it much cheaper.

Scientists custom-build fairly short strings DNA now for research, but scaling up enough to handle information storage in bulk would require a "mind-boggling" leap in output, Kosuri says. With current technology, that would be hugely expensive, he says.

George Church, a prominent Harvard genetics expert, agrees that cost is a big issue. But "I'm pretty optimistic it can be brought down" dramatically in a decade or less, says Church, who is in the process of starting a company to offer DNA storage methods.

For all the interest in the topic, it's worth noting that so far the amount of information that researchers have stored in DNA is relatively tiny.

Earlier this month, Microsoft announced that a team including Strauss and Ceze had stored a record 200 megabytes. The information included 100 books—one, fittingly, was "Great Expectations"—along with a brief video and many documents. But it was still less than 5 percent the capacity of an ordinary DVD.

Yet it's about nine times the mark reported just last month by Church, who says the announcement shows "how fast the field is moving."

Meanwhile, people involved with archiving digital data say their field views DNA as a possibility for the future, but not a cure-all.

"It's a very interesting and promising approach to the storage problem, but the storage problem is really only a very small part of digital preservation," says Cal Lee, a professor at the University of North Carolina's School of Information and Library Science.

It's true that society will probably always have devices to read DNA, so that gets around the problem of obsolete readers, he says. But that's not enough.

"If you just read the ones and zeroes, you don't know how to interpret it," Lee says.

For example, is that string a picture, text, a sound clip or a video? Do you still have the software to make sense of it?

What's more, the people in charge of keeping digital information want to check on it periodically to make sure it's still intact, and "I don't know how viable that is with DNA," says Euan Cochrane, digital preservation manager at the Yale University Library. It may mean fewer such check-ups, he says.

Cochrane, who describes his job as keeping information accessible "10 years to forever," says DNA looks interesting if its cost can be reduced and scientists find ways to more quickly store and recover information.

Starr says his data-storage device company hasn't taken a detailed look at DNA technology because it's too far in the future.

There are "always things out on the horizon that could store data for a very long time," he says. But the challenge of turning those ideas into a practical product "really trims the field down pretty quickly." [13]

Second layer of information in DNA confirmed

Leiden theoretical physicists have proven that DNA mechanics, in addition to genetic information in DNA, determines who we are. Helmut Schiessel and his group simulated many DNA sequences and

found a correlation between mechanical cues and the way DNA is folded. They have published their results in PLoS One.

When James Watson and Francis Crick identified the structure of DNA molecules in 1953, they revealed that DNA information determines who we are. The sequence of the letters G, A, T and C in the famous double helix determines what proteins are made by our cells. If you have brown eyes, for example, this is because a series of letters in your DNA encodes for proteins that build brown eyes. Each cell contains the exact same letter sequence, and yet every organ behaves differently. How is this possible?

Mechanical cues

Since the mid 1980s, it has been hypothesized that there is a second layer of information on top of the genetic code consisting of DNA mechanical properties.

Each of our cells contains two meters of DNA molecules, and these molecules need to be wrapped up tightly to fit inside a single cell. The way in which DNA is folded determines how the letters are read out, and therefore which proteins are actually made. In each organ, only relevant parts of the genetic information are read. The theory suggests that mechanical cues within the DNA structures determine how preferentially DNA folds.


Simulation

For the first time, Leiden physicist Helmut Schiessel and his research group provide strong evidence that this second layer of information indeed exists. With their computer code, they have simulated the folding of DNA strands with randomly assigned mechanical cues. It turns out that these cues indeed determine how the DNA molecule is folded into so-called nucleosomes. Schiessel found correlations between the mechanics and the actual folding structure in the genome of two organisms—baker's yeast and fission yeast. This finding reveals evolutionary changes in DNA—mutations—that have two very different effects: The letter sequence encoding for a specific protein can change, or the mechanics of the DNA structure can change, resulting in different packaging and levels of DNA accessibility, and therefore differing frequency of production of that protein. [12]

Quantum entanglement between the electron clouds of nucleic acids in DNA

We model the electron clouds of nucleic acids in DNA as a chain of coupled quantum harmonic oscillators with dipole-dipole interaction between nearest neighbours resulting in a van der Waals type bonding. Crucial parameters in our model are the distances between the acids and the coupling between them, which we estimate from numerical simulations. We show that for realistic parameters nearest neighbour entanglement is present even at room temperature. We quantify the amount of entanglement in terms of negativity and single base von Neumann entropy. We find that the strength of the single base von Neumann entropy depends on the neighbouring sites, thus questioning the notion of treating single bases as logically independent units. We derive an analytical expression for the binding energy of the coupled chain in terms of entanglement and show the connection between entanglement and correlation energy, a quantity commonly used in quantum chemistry. [11]

Scientists discover secret code hidden within human DNA



This undated handout illustration shows the DNA double helix (AFP Photo) This undated handout illustration shows the DNA double helix (AFP Photo)

Scientists have discovered a secret second code hiding within DNA which instructs cells on how genes are controlled. The amazing discovery is expected to open new doors to the diagnosis and treatment of diseases, according to a new study.

Ever since the genetic code was deciphered over 40 years ago, scientists have believed that it only described how proteins are made. However, the revelation made by the research team led by John Stamatoyannopoulos of the University of Washington indicates that genomes use the genetic code to write two separate languages.

“For over 40 years we have assumed that DNA changes affecting the genetic code solely impact how proteins are made,” said Stamatoyannopoulos, according to the press release. “Now we know that this basic assumption about reading the human genome missed half of the picture.”

Scientists discovered that the second language instructs the cells on how genes are controlled, according to findings published in *Science* magazine on Friday. The study is part of the Encyclopedia of DNA Elements Project, also known as ENCODE.

DNA (Deoxyribonucleic acid) is a nucleic acid that is the main constituent of the chromosomes of all organisms, except some viruses. DNA is self-replicating, plays a central role in protein synthesis, and is responsible for the transmission of hereditary characteristics from parents to offspring.

The second language remained hidden for so long because one language is written on top of the other, scientists said.

Scientists already knew that the genetic code uses a 64-letter alphabet called codons. The research team discovered that some of the codons can have two meanings – one related to proteins, the other to gene control. Those codons were given the name ‘duons.’

And it’s those duons that are expected to change the way physicians interpret human genomes, and give clues for the treatments of diseases.

“The fact that the genetic code can simultaneously write two kinds of information means that many DNA changes that appear to alter protein sequences may actually cause disease by disrupting gene control programs or even both mechanisms simultaneously,” said Stamatoyannopoulos.


Speaking about the discovery, Stamatoyannopoulos said that the “new findings highlight that DNA is an incredibly powerful information storage device, which nature has fully exploited in unexpected ways.” [10]

This Physicist Has a Groundbreaking Idea about Why Life Exists

“You start with a random clump of atoms, and if you shine light on it for long enough, it should not be so surprising that you get a plant,” England said.

England’s theory is meant to underlie, rather than replace, Darwin’s theory of evolution by natural selection, which provides a powerful description of life at the level of genes and populations. “I am certainly not saying that Darwinian ideas are wrong,” he explained. “On the contrary, I am just saying that from the perspective of the physics, you might call Darwinian evolution a special case of a more general phenomenon.”

At the heart of England’s idea is the second law of thermodynamics, also known as the law of increasing entropy or the “arrow of time.” Hot things cool down, gas diffuses through air, eggs scramble but never spontaneously unscramble; in short, energy tends to disperse or spread out as time progresses. Entropy is a measure of this tendency, quantifying how dispersed the energy is among the particles in a system, and how diffuse those particles are throughout space. It increases as a simple matter of probability: There are more ways for energy to be spread out than for it to be concentrated.



A computer simulation by Jeremy England and colleagues shows a system of particles confined inside a viscous fluid in which the turquoise particles are driven by an oscillating force. Over time (from top to bottom), the force triggers the formation of more bonds among the particles.


Thus, as particles in a system move around and interact, they will, through sheer chance, tend to adopt configurations in which the energy is spread out. Eventually, the system arrives at a state of maximum entropy called “thermodynamic equilibrium,” in which energy is uniformly distributed. A cup of coffee and the room it sits in become the same temperature, for example.

Although entropy must increase over time in an isolated or “closed” system, an “open” system can keep its entropy low — that is, divide energy unevenly among its atoms — by greatly increasing the entropy of its surroundings. In his influential 1944 monograph “What Is Life?” the eminent quantum physicist Erwin Schrödinger argued that this is what living things must do. A plant, for example, absorbs extremely energetic sunlight, uses it to build sugars, and ejects infrared light, a much less

concentrated form of energy. The overall entropy of the universe increases during photosynthesis as the sunlight dissipates, even as the plant prevents itself from decaying by maintaining an orderly internal structure.

Self-replication (or reproduction, in biological terms), the process that drives the evolution of life on Earth, is one such mechanism by which a system might dissipate an increasing amount of energy over time.

As England put it, "A great way of dissipating more is to make more copies of yourself."



Self-Replicating Sphere Clusters: According to new research at Harvard, coating the surfaces of microspheres can cause them to spontaneously assemble into a chosen structure, such as a polytetrahedron (red), which then triggers nearby spheres into forming an identical structure.

Scientists have already observed self-replication in nonliving systems. According to new research led by Philip Marcus of the University of California, Berkeley, and reported in *Physical Review Letters* in August, vortices in turbulent fluids spontaneously replicate themselves by drawing energy from shear in the surrounding fluid. And in a paper in *Proceedings of the National Academy of Sciences*, Michael Brenner, a professor of applied mathematics and physics at Harvard, and his collaborators present theoretical models and simulations of microstructures that self-replicate. These clusters of specially coated microspheres dissipate energy by roping nearby spheres into forming identical clusters. "This connects very much to what Jeremy is saying," Brenner said. [8]

Photoactive Prebiotic Systems

We propose that life first emerged in the form of such minimal photoactive prebiotic kernel systems and later in the process of evolution these photoactive prebiotic kernel systems would have produced fatty acids and covered themselves with fatty acid envelopes to become the minimal cells of the Fatty Acid World. Specifically, we model self-assembling of photoactive prebiotic systems with observed quantum entanglement phenomena. We address the idea that quantum entanglement was important in the first stages of origins of life and evolution of the biospheres because simultaneously excite two prebiotic kernels in the system by appearance of two additional quantum entangled excited states, leading to faster growth and self-replication of minimal living cells. The quantum mechanically modeled possibility of synthesizing artificial self-reproducing quantum entangled prebiotic kernel systems and minimal cells also impacts the possibility of the most probable path of emergence of photocells on the Earth or elsewhere. We also examine the quantum entangled logic gates discovered in the modeled systems composed of two prebiotic kernels. Such logic gates may have application in the destruction of cancer cells or becoming building blocks of new forms of artificial cells including magnetically active ones.

Significance Statement

Our investigated self-assembly of molecules towards supramolecular bioorganic and minimal cellular systems depends on the quantum mechanics laws which induce hydrogen and Van der Waals bindings (Tamulis A, Grigalavicius, M, *Orig Life Evol Biosph* 41:51-71, 2011).

In the work presented here, quantum entanglement takes the form of a quantum superposition of the active components in synthesized self-assembling and self-replicating living systems. When a quantum calculation of an entangled system is made that causes one photoactive biomolecule of such a pair to take on a definite value (e.g., electron density transfer or electron spin density transfer), the other member of this entangled pair will be found to have taken the appropriately correlated value (e.g., electron density transfer or electron spin density transfer). In our simulations, the separation distance of supramolecular bio systems changes took place during geometry optimization procedures, which mimic real-world intermolecular interaction processes.


Our discovered phenomenon of the quantum entanglement in the prebiotic systems enhance the photosynthesis in the proposed systems because simultaneously excite two prebiotic kernels in the system by appearance of two additional quantum entangled excited states (Tamulis A, Grigalavicius M, Baltrusaitis J, *Orig Life Evol Biosph* 43:49-66, 2013; Tamulis A, Grigalavicius M, Krisciukaitis S (2014), *J Comput Theor Nanos*, 11, 1597-1608, 2014; Tamulis A, Grigalavicius M, 8:117-140, 2014.). We can propose that quantum entanglement enhanced the emergence of photosynthetic prebiotic kernels and accelerated the evolution of photosynthetic life because of additional absorbed light energy, leading to faster growth and self-replication of minimal living cells.

We can state that: Livings are self-assembled and self-replicating wet and warm stochastically moving supramolecular systems where quantum entanglement can be continuously generated and destroyed by non-equilibrium effects in an environment where no static entanglement exists; quantum entanglement involve the biomolecule inside one living or between other neighboring livings.

This warm quantum coherence is basic for the explanation of DNA stability and for the understanding of brain magnetic orientation during migration in more than 50 species of birds, fishes and insects. Exists experimental evidence for quantum-coherent is used for more efficient light-harvesting in plant photosynthesis. Quantum entanglement exists in supramolecules determining the sense of smell and in the brain neurons microtubules due to quantum vibrations.

In the work presented here, we started to design and quantum mechanical investigations of the molecular logical devices which are useful for construction of nano medicine biorobots against the molecular diseases such a cancer tumors, and against the new kinds of synthesized microorganisms and nano guns.

Figure legend



You can see in the enclosed figure the quantum entanglement phenomenon in the closely self-assembled two synthesized protocell system due to the photo excited electron charge transfer from one protocell to another that leads to closer self-assembly and exchange of energy and information.

Visualization of the electron charge tunneling associated with the 6th (467.3 nm) excited state. The transition is mainly from squaraine molecule of the first protocell situated in the bottom of this bi cellular system to precursor of fatty acid (pFA) molecule of the second subsystem (in the top) and little from the 1,4-bis(N,N-dimethylamino)naphthalene molecule (in the top-right) to the same pFA molecule of the second subsystem (in the top). The electron cloud hole is indicated by the dark blue color while the transferred electron cloud location is designated by the gray color.

As a result, these nonlinear quantum interactions compressed the overall molecular system resulting in a smaller gap between the HOMO and LUMO electron energy levels which allows enhanced tunneling of photo excited electrons from the sensitizer squaraine and (1,4-bis(N,N-dimethylamino)naphthalene) to the pFA molecule resulting in its cleavage. The new fatty acid joins the existing minimal cell thus increasing it in size. After reaching some critical size, the minimal cell should divide (i.e. self-replicate) into two separate smaller minimal cells. [7]

Quantum Biology

Researchers have long suspected that something unusual is afoot in photosynthesis. Particles of light called photons, streaming down from the Sun; arrive randomly at the chlorophyll molecules and other light-absorbing 'antenna' pigments that cluster inside the cells of every leaf, and within every photosynthetic bacterium. But once the photons' energy is deposited, it doesn't stay random. Somehow, it gets channeled into a steady flow towards the cell's photosynthetic reaction centre, which can then use it at maximum efficiency to convert carbon dioxide into sugars. Quantum coherence in photosynthesis seems to be beneficial to the organisms using it. But did their ability to exploit quantum effects evolve through natural selection? Or is quantum coherence just an accidental side effect of the way certain molecules are structured? [6]

Quantum Consciousness

Extensive scientific investigation has found that a form of quantum coherence operates within living biological systems through what is known as biological excitations and biophoton emission. What this means is that metabolic energy is stored as a form of electromechanical and electromagnetic excitations. These coherent excitations are considered responsible for generating and maintaining long-range order via the transformation of energy and very weak electromagnetic signals. After nearly twenty years of experimental research, Fritz-Albert Popp put forward the hypothesis that biophotons are emitted from a coherent electrodynamics field within the living system.

What this means is that each living cell is giving off, or resonating, a biophoton field of coherent energy. If each cell is emitting this field, then the whole living system is, in effect, a resonating field-a ubiquitous nonlocal field. And since biophotons are the entities through which the living system communicates, there is near-instantaneous intercommunication throughout. And this, claims Popp, is the basis for coherent biological organization -- referred to as quantum coherence. This discovery

led Popp to state that the capacity for evolution rests not on aggressive struggle and rivalry but on the capacity for communication and cooperation. In this sense the built-in capacity for species evolution is not based on the individual but rather living systems that are interlinked within a coherent whole: Living systems are thus neither the subjects alone, nor objects isolated, but both subjects and objects in a mutually communicating universe of meaning. . . . Just as the cells in an organism take on different tasks for the whole, different populations enfold information not only for themselves, but for all other organisms, expanding the consciousness of the whole, while at the same time becoming more and more aware of this collective consciousness.

Biophysicist Mae-Wan Ho describes how the living organism, including the human body, is coordinated throughout and is "coherent beyond our wildest dreams." It appears that every part of our body is "in communication with every other part through a dynamic, tunable, responsive, liquid crystalline medium that pervades the whole body, from organs and tissues to the interior of every cell."

What this tells us is that the medium of our bodies is a form of liquid crystal, an ideal transmitter of communication, resonance, and coherence. These relatively new developments in biophysics have discovered that all biological organisms are constituted of a liquid crystalline medium. Further, DNA is a liquid-crystal, lattice-type structure (which some refer to as a liquid crystal gel), whereby body cells are involved in a holographic instantaneous communication via the emitting of biophotons (a source based on light). This implies that all living biological organisms continuously emit radiations of light that form a field of coherence and communication. Moreover, biophysics has discovered that living organisms are permeated by quantum wave forms. [5]

Information – Entropy Theory of Physics

Viewing the confined gas where the statistical entropy not needs the information addition is not the only physical system. There are for example quantum mechanical systems where the information is a very important qualification. The perturbation theory needs higher order calculations in QED or QCD giving more information on the system as in the chess games happens, where the entropy is not enough to describe the state of the matter. The variation calculation of chess is the same as the perturbation calculation of physics to gain information, where the numbers of particles are small for statistical entropy to describe the system. The role of the Feynman graphs are the same as the chess variations of a given position that is the depth of the variations tree, the Information is the same as the order of the Feynman graphs giving the Information of the micro system. [9]

Information – Entropy Theory of Life

There is also connection between statistical physics and evolutionary biology, since the arrow of time is working in the biological evolution also.

The Fluctuation Theorem says that there is a probability that entropy will flow in a direction opposite to that dictated by the Second Law of Thermodynamics. In this case the Information is growing that

is the matter formulas are emerging from the chaos. So the Weak Interaction has two directions, samples for one direction is the Neutron decay, and Hydrogen fusion is the opposite direction. The living biological systems have also entropy lowering and information growing direction by building more complicated or entangled molecules, governed by the quantum mechanics and the general weak interaction. On the other hand there is the arrow of time; the entropy growing is lowering the information by dissipating these entangled or otherwise connected biomolecules, aging the living systems.

Creating quantum technology

Another area of potential application is in quantum computing. The long-standing goal of the physicists and engineers working in this area is to manipulate data encoded in quantum bits (qubits) of information, such as the spin-up and spin-down states of an electron or of an atomic nucleus. Qubits can exist in both states at once, thus permitting the simultaneous exploration of all possible answers to the computation that they encode. In principle, this would give quantum computers the power to find the best solution far more quickly than today's computers can — but only if the qubits can maintain their coherence, without the noise of the surrounding environment, such as the jostling of neighboring atoms, destroying the synchrony of the waves. [6]

Quantum Entanglement

Measurements of physical properties such as position, momentum, spin, polarization, etc. performed on entangled particles are found to be appropriately correlated. For example, if a pair of particles is generated in such a way that their total spin is known to be zero, and one particle is found to have clockwise spin on a certain axis, then the spin of the other particle, measured on the same axis, will be found to be counterclockwise. Because of the nature of quantum measurement, however, this behavior gives rise to effects that can appear paradoxical: any measurement of a property of a particle can be seen as acting on that particle (e.g. by collapsing a number of superimposed states); and in the case of entangled particles, such action must be on the entangled system as a whole. It thus appears that one particle of an entangled pair "knows" what measurement has been performed on the other, and with what outcome, even though there is no known means for such information to be communicated between the particles, which at the time of measurement may be separated by arbitrarily large distances. [4]

The Bridge

The accelerating electrons explain not only the Maxwell Equations and the Special Relativity, but the Heisenberg Uncertainty Relation, the wave particle duality and the electron's spin also, building the bridge between the Classical and Quantum Theories. [1]

Accelerating charges

The moving charges are self maintain the electromagnetic field locally, causing their movement and this is the result of their acceleration under the force of this field. In the classical physics the charges will distributed along the electric current so that the electric potential lowering along the current, by

linearly increasing the way they take every next time period because this accelerated motion. The same thing happens on the atomic scale giving a Δp impulse difference and a Δx way difference between the different part of the not point like particles.

Relativistic effect

Another bridge between the classical and quantum mechanics in the realm of relativity is that the charge distribution is lowering in the reference frame of the accelerating charges linearly: $ds/dt = at$ (time coordinate), but in the reference frame of the current it is parabolic: $s = a/2 t^2$ (geometric coordinate).

Heisenberg Uncertainty Relation

In the atomic scale the Heisenberg uncertainty relation gives the same result, since the moving electron in the atom accelerating in the electric field of the proton, causing a charge distribution on Δx position difference and with a Δp momentum difference such a way that they product is about the half Planck reduced constant. For the proton this Δx much less in the nucleon, than in the orbit of the electron in the atom, the Δp is much higher because of the greater proton mass.

This means that the electron and proton are not point like particles, but has a real charge distribution.

Wave – Particle Duality

The accelerating electrons explains the wave – particle duality of the electrons and photons, since the elementary charges are distributed on Δx position with Δp impulse and creating a wave packet of the electron. The photon gives the electromagnetic particle of the mediating force of the electrons electromagnetic field with the same distribution of wavelengths.

Atomic model

The constantly accelerating electron in the Hydrogen atom is moving on the equipotential line of the proton and it's kinetic and potential energy will be constant. Its energy will change only when it is changing its way to another equipotential line with another value of potential energy or getting free with enough kinetic energy. This means that the Rutherford-Bohr atomic model is right and only that changing acceleration of the electric charge causes radiation, not the steady acceleration. The steady acceleration of the charges only creates a centric parabolic steady electric field around the charge, the magnetic field. This gives the magnetic moment of the atoms, summing up the proton and electron magnetic moments caused by their circular motions and spins.

The Relativistic Bridge

Commonly accepted idea that the relativistic effect on the particle physics it is the fermions' spin - another unresolved problem in the classical concepts. If the electric charges can move only with accelerated motions in the self maintaining electromagnetic field, once upon a time they would

reach the velocity of the electromagnetic field. The resolution of this problem is the spinning particle, constantly accelerating and not reaching the velocity of light because the acceleration is radial. One origin of the Quantum Physics is the Planck Distribution Law of the electromagnetic oscillators, giving equal intensity for 2 different wavelengths on any temperature. Any of these two wavelengths will give equal intensity diffraction patterns, building different asymmetric constructions, for example proton - electron structures (atoms), molecules, etc. Since the particles are centers of diffraction patterns they also have particle – wave duality as the electromagnetic waves have. [2]

The weak interaction

The weak interaction transforms an electric charge in the diffraction pattern from one side to the other side, causing an electric dipole momentum change, which violates the CP and time reversal symmetry. The Electroweak Interaction shows that the Weak Interaction is basically electromagnetic in nature. The arrow of time shows the entropy grows by changing the temperature dependent diffraction patterns of the electromagnetic oscillators.

Another important issue of the quark model is when one quark changes its flavor such that a linear oscillation transforms into plane oscillation or vice versa, changing the charge value with 1 or -1. This kind of change in the oscillation mode requires not only parity change, but also charge and time changes (CPT symmetry) resulting a right handed anti-neutrino or a left handed neutrino.

The right handed anti-neutrino and the left handed neutrino exist only because changing back the quark flavor could happen only in reverse, because they are different geometrical constructions, the u is 2 dimensional and positively charged and the d is 1 dimensional and negatively charged. It needs also a time reversal, because anti particle (anti neutrino) is involved.

The neutrino is a 1/2 spin creator particle to make equal the spins of the weak interaction, for example neutron decay to 2 fermions, every particle is fermions with $\frac{1}{2}$ spin. The weak interaction changes the entropy since more or less particles will give more or less freedom of movement. The entropy change is a result of temperature change and breaks the equality of oscillator diffraction intensity of the Maxwell–Boltzmann statistics. This way it changes the time coordinate measure and makes possible a different time dilation as of the special relativity.

The limit of the velocity of particles as the speed of light appropriate only for electrical charged particles, since the accelerated charges are self maintaining locally the accelerating electric force. The neutrinos are CP symmetry breaking particles compensated by time in the CPT symmetry, that is the time coordinate not works as in the electromagnetic interactions, consequently the speed of neutrinos is not limited by the speed of light.

The weak interaction T-asymmetry is in conjunction with the T-asymmetry of the second law of thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes the weak interaction, for example the Hydrogen fusion.

Probably because it is a spin creating movement changing linear oscillation to 2 dimensional oscillation by changing d to u quark and creating anti neutrino going back in time relative to the

proton and electron created from the neutron, it seems that the anti neutrino fastest then the velocity of the photons created also in this weak interaction?

A quark flavor changing shows that it is a reflection changes movement and the CP- and T- symmetry breaking!!! This flavor changing oscillation could prove that it could be also on higher level such as atoms, molecules, probably big biological significant molecules and responsible on the aging of the life.

Important to mention that the weak interaction is always contains particles and antiparticles, where the neutrinos (antineutrinos) present the opposite side. It means by Feynman's interpretation that these particles present the backward time and probably because this they seem to move faster than the speed of light in the reference frame of the other side.

Finally since the weak interaction is an electric dipole change with $\frac{1}{2}$ spin creating; it is limited by the velocity of the electromagnetic wave, so the neutrino's velocity cannot exceed the velocity of light.

The General Weak Interaction

The Weak Interactions T-asymmetry is in conjunction with the T-asymmetry of the Second Law of Thermodynamics, meaning that locally lowering entropy (on extremely high temperature) causes for example the Hydrogen fusion. The arrow of time by the Second Law of Thermodynamics shows the increasing entropy and decreasing information by the Weak Interaction, changing the temperature dependent diffraction patterns. A good example of this is the neutron decay, creating more particles with less known information about them.

The neutrino oscillation of the Weak Interaction shows that it is a general electric dipole change and it is possible to any other temperature dependent entropy and information changing diffraction pattern of atoms, molecules and even complicated biological living structures.

We can generalize the weak interaction on all of the decaying matter constructions, even on the biological too. This gives the limited lifetime for the biological constructions also by the arrow of time. There should be a new research space of the Quantum Information Science the 'general neutrino oscillation' for the greater than subatomic matter structures as an electric dipole change. There is also connection between statistical physics and evolutionary biology, since the arrow of time is working in the biological evolution also.

The Fluctuation Theorem says that there is a probability that entropy will flow in a direction opposite to that dictated by the Second Law of Thermodynamics. In this case the Information is growing that is the matter formulas are emerging from the chaos. So the Weak Interaction has two directions, samples for one direction is the Neutron decay, and Hydrogen fusion is the opposite direction.

Fermions and Bosons

The fermions are the diffraction patterns of the bosons such a way that they are both sides of the same thing.

Van Der Waals force

Named after the Dutch scientist Johannes Diderik van der Waals – who first proposed it in 1873 to explain the behaviour of gases – it is a very weak force that only becomes relevant when atoms and molecules are very close together. Fluctuations in the electronic cloud of an atom mean that it will

have an instantaneous dipole moment. This can induce a dipole moment in a nearby atom, the result being an attractive dipole–dipole interaction.

Electromagnetic inertia and mass

Electromagnetic Induction

Since the magnetic induction creates a negative electric field as a result of the changing acceleration, it works as an electromagnetic inertia, causing an electromagnetic mass. [1]

Relativistic change of mass

The increasing mass of the electric charges the result of the increasing inductive electric force acting against the accelerating force. The decreasing mass of the decreasing acceleration is the result of the inductive electric force acting against the decreasing force. This is the relativistic mass change explanation, especially importantly explaining the mass reduction in case of velocity decrease.

The frequency dependence of mass

Since $E = h\nu$ and $E = mc^2$, $m = h\nu / c^2$ that is the m depends only on the ν frequency. It means that the mass of the proton and electron are electromagnetic and the result of the electromagnetic induction, caused by the changing acceleration of the spinning and moving charge! It could be that the m_0 inertial mass is the result of the spin, since this is the only accelerating motion of the electric charge. Since the accelerating motion has different frequency for the electron in the atom and the proton, they masses are different, also as the wavelengths on both sides of the diffraction pattern, giving equal intensity of radiation.

Electron – Proton mass rate

The Planck distribution law explains the different frequencies of the proton and electron, giving equal intensity to different lambda wavelengths! Also since the particles are diffraction patterns they have some closeness to each other – can be seen as a gravitational force. [2]

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

Gravity from the point of view of quantum physics

The Gravitational force

The gravitational attractive force is basically a magnetic force.

The same electric charges can attract one another by the magnetic force if they are moving parallel in the same direction. Since the electrically neutral matter is composed of negative and positive

charges they need 2 photons to mediate this attractive force, one per charges. The Big Bang caused parallel moving of the matter gives this magnetic force, experienced as gravitational force.

Since graviton is a tensor field, it has spin = 2, could be 2 photons with spin = 1 together.

You can think about photons as virtual electron – positron pairs, obtaining the necessary virtual mass for gravity.

The mass as seen before a result of the diffraction, for example the proton – electron mass ratio $M_p=1840 M_e$. In order to move one of these diffraction maximum (electron or proton) we need to intervene into the diffraction pattern with a force appropriate to the intensity of this diffraction maximum, means its intensity or mass.

The Big Bang caused acceleration created radial currents of the matter, and since the matter is composed of negative and positive charges, these currents are creating magnetic field and attracting forces between the parallel moving electric currents. This is the gravitational force experienced by the matter, and also the mass is result of the electromagnetic forces between the charged particles. The positive and negative charged currents attracts each other or by the magnetic forces or by the much stronger electrostatic forces!?

The gravitational force attracting the matter, causing concentration of the matter in a small space and leaving much space with low matter concentration: dark matter and energy.

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

The Higgs boson

By March 2013, the particle had been proven to behave, interact and decay in many of the expected ways predicted by the Standard Model, and was also tentatively confirmed to have + parity and zero spin, two fundamental criteria of a Higgs boson, making it also the first known scalar particle to be discovered in nature, although a number of other properties were not fully proven and some partial results do not yet precisely match those expected; in some cases data is also still awaited or being analyzed.

Since the Higgs boson is necessary to the W and Z bosons, the dipole change of the Weak interaction and the change in the magnetic effect caused gravitation must be conducted. The Wien law is also important to explain the Weak interaction, since it describes the T_{max} change and the diffraction patterns change. [2]

Higgs mechanism and Quantum Gravity

The magnetic induction creates a negative electric field, causing an electromagnetic inertia. Probably it is the mysterious Higgs field giving mass to the charged particles? We can think about the photon as an electron-positron pair, they have mass. The neutral particles are built from negative and positive charges, for example the neutron, decaying to proton and electron. The wave – particle duality makes sure that the particles are oscillating and creating magnetic induction as an inertial mass, explaining also the relativistic mass change. Higher frequency creates stronger magnetic induction, smaller frequency results lesser magnetic induction. It seems to me that the magnetic induction is the secret of the Higgs field.

In particle physics, the Higgs mechanism is a kind of mass generation mechanism, a process that gives mass to elementary particles. According to this theory, particles gain mass by interacting with the Higgs field that permeates all space. More precisely, the Higgs mechanism endows gauge bosons in a gauge theory with mass through absorption of Nambu–Goldstone bosons arising in spontaneous symmetry breaking.

The simplest implementation of the mechanism adds an extra Higgs field to the gauge theory. The spontaneous symmetry breaking of the underlying local symmetry triggers conversion of components of this Higgs field to Goldstone bosons which interact with (at least some of) the other fields in the theory, so as to produce mass terms for (at least some of) the gauge bosons. This mechanism may also leave behind elementary scalar (spin-0) particles, known as Higgs bosons.

In the Standard Model, the phrase "Higgs mechanism" refers specifically to the generation of masses for the W^\pm , and Z weak gauge bosons through electroweak symmetry breaking. The Large Hadron Collider at CERN announced results consistent with the Higgs particle on July 4, 2012 but stressed that further testing is needed to confirm the Standard Model.

What is the Spin?

So we know already that the new particle has spin zero or spin two and we could tell which one if we could detect the polarizations of the photons produced. Unfortunately this is difficult and neither ATLAS nor CMS are able to measure polarizations. The only direct and sure way to confirm that the particle is indeed a scalar is to plot the angular distribution of the photons in the rest frame of the centre of mass. A spin zero particles like the Higgs carries no directional information away from the original collision so the distribution will be even in all directions. This test will be possible when a much larger number of events have been observed. In the mean time we can settle for less certain indirect indicators.

The Graviton

In physics, the graviton is a hypothetical elementary particle that mediates the force of gravitation in the framework of quantum field theory. If it exists, the graviton is expected to be massless (because the gravitational force appears to have unlimited range) and must be a spin-2 boson. The spin follows from the fact that the source of gravitation is the stress-energy tensor, a second-rank tensor (compared to electromagnetism's spin-1 photon, the source of which is the four-current, a first-rank tensor). Additionally, it can be shown that any massless spin-2 field would give rise to a force indistinguishable from gravitation, because a massless spin-2 field must couple to (interact with) the stress-energy tensor in the same way that the gravitational field does. This result suggests that, if a massless spin-2 particle is discovered, it must be the graviton, so that the only experimental verification needed for the graviton may simply be the discovery of a massless spin-2 particle. [3]

Conclusions

“The fact that the genetic code can simultaneously write two kinds of information means that many DNA changes that appear to alter protein sequences may actually cause disease by disrupting gene control programs or even both mechanisms simultaneously,” said Stamatoyannopoulos.

Speaking about the discovery, Stamatoyannopoulos said that the “new findings highlight that DNA is an incredibly powerful information storage device, which nature has fully exploited in unexpected ways.” [10]

There is also connection between statistical physics and evolutionary biology, since the arrow of time is working in the biological evolution also.

Prentiss, who runs an experimental biophysics lab at Harvard, says England’s theory could be tested by comparing cells with different mutations and looking for a correlation between the amount of energy the cells dissipate and their replication rates. [8]

Exists experimental evidence for quantum-coherent is used for more efficient light-harvesting in plant photosynthesis. Quantum entanglement exists in supramolecules determining the sense of smell and in the brain neurons microtubules due to quantum vibrations.

In the work presented here, we started to design and quantum mechanical investigations of the molecular logical devices which are useful for construction of nano medicine biorobots against the molecular diseases such a cancer tumors, and against the new kinds of synthesized microorganisms and nano guns. [7]

One of the most important conclusions is that the electric charges are moving in an accelerated way and even if their velocity is constant, they have an intrinsic acceleration anyway, the so called spin, since they need at least an intrinsic acceleration to make possible they movement .

The accelerated charges self-maintaining potential shows the locality of the relativity, working on the quantum level also. [1]

The bridge between the classical and quantum theory is based on this intrinsic acceleration of the spin, explaining also the Heisenberg Uncertainty Principle. The particle – wave duality of the electric charges and the photon makes certain that they are both sides of the same thing.

The Secret of Quantum Entanglement that the particles are diffraction patterns of the electromagnetic waves and this way their quantum states every time is the result of the quantum state of the intermediate electromagnetic waves. [2]

These relatively new developments in biophysics have discovered that all biological organisms are constituted of a liquid crystalline medium. Further, DNA is a liquid-crystal, lattice-type structure (which some refer to as a liquid crystal gel), whereby body cells are involved in a holographic instantaneous communication via the emitting of biophotons (a source based on light). This implies that all living biological organisms continuously emit radiations of light that form a field of coherence and communication. Moreover, biophysics has discovered that living organisms are permeated by quantum wave forms. [5]

Basing the gravitational force on the accelerating Universe caused magnetic force and the Planck Distribution Law of the electromagnetic waves caused diffraction gives us the basis to build a Unified Theory of the physical interactions also.

References

[1] The Magnetic field of the Electric current and the Magnetic induction

http://academia.edu/3833335/The_Magnetic_field_of_the_Electric_current

[2] 3 Dimensional String Theory

http://academia.edu/3834454/3_Dimensional_String_Theory

[3] Graviton Production By Two Photon and Electron-Photon Processes In Kaluza-Klein Theories With Large Extra Dimensions

<http://arxiv.org/abs/hep-ph/9909392>

[4] Quantum Entanglement

http://en.wikipedia.org/wiki/Quantum_entanglement

[5] Quantum Consciousness

https://realitysandwich.com/155783/quantum_consciousness_0/

[6] Quantum Biology

<http://www.nature.com/news/2011/110615/pdf/474272a.pdf>

[7] Quantum entanglement in photoactive prebiotic systems

<https://globalmedicaldiscovery.com/key-drug-discovery-articles/quantum-entanglement-photoactive-prebiotic-systems/>

[8] http://www.businessinsider.com/groundbreaking-idea-of-lifes-origin-2014-12?pundits_only=0&comments_page=0#comment-548823e769bedd444b63cd87

[9] Information – Entropy Theory of Physics

https://www.academia.edu/3836084/Information_-_Entropy_Theory_of_Physics

[10] Scientists discover secret code hidden within human DNA

<http://rt.com/news/scientists-discover-dna-code-222/>

[11] Quantum entanglement between the electron clouds of nucleic acids in DNA

<http://arxiv.org/abs/1006.4053>

[12] Second layer of information in DNA confirmed

<http://phys.org/news/2016-06-layer-dna.html>

[13] Scientists work toward storing digital information in DNA

<http://phys.org/news/2016-07-scientists-digital-dna.html>

[14] IBM lab-on-a-chip breakthrough aims to help physicians detect cancer

<http://phys.org/news/2016-08-ibm-lab-on-a-chip-breakthrough-aims-physicians.html>

[15] Fighting cancer with space research

<http://phys.org/news/2016-09-cancer-space.html>

[16] When push comes to shove: Size matters for particles in our bloodstream

<http://phys.org/news/2016-10-shove-size-particles-bloodstream.html>

[17] New drug limits and then repairs brain damage in stroke

<http://medicalxpress.com/news/2016-11-drug-limits-brain.html>

[18] Visualizing gene expression with MRI

<http://phys.org/news/2016-12-visualizing-gene-mri.html>

[19] Super-resolution system reveals mechanics of tiny 'DNA walker'

<https://phys.org/news/2017-02-super-resolution-reveals-mechanics-tiny-dna.html>

[20] Computing with biochemical circuits made easy

<https://phys.org/news/2017-02-biochemical-circuits-easy.html>

[21] One day, more than your genes will be stored in DNA

<https://cosmosmagazine.com/technology/one-day-more-than-your-genes-will-be-stored-in-dna>

[22] Researchers launch plasmons with controlled amounts of angular momentum

<https://phys.org/news/2017-03-plasmons-amounts-angular-momentum.html>

[23] A novel 'soft' magnetic material could enable faster computer memory

<https://phys.org/news/2017-03-soft-magnetic-material-enable-faster.html>

[24] Spintronic technology advances with newly designed magnetic tunnel junctions

<https://phys.org/news/2017-03-spintronic-technology-advances-newly-magnetic.html>