

Cosmological Collider

In a recent research, scientists from HKUST and Harvard University revealed the connection between those two aspects, and argued that our universe could be used as a particle physics "collider" to study the high energy particle physics. [8]

According to Einstein's theory of relativity, the curvature of spacetime was infinite at the big bang. In fact, at this point all mathematical tools fail, and the theory breaks down. [7]

The universe may have existed forever, according to a new model that applies quantum correction terms to complement Einstein's theory of general relativity. The model may also account for dark matter and dark energy, resolving multiple problems at once. [6]

This paper explains the Accelerating Universe, the Special and General Relativity from the observed effects of the accelerating electrons, causing naturally the experienced changes of the electric field potential along the moving electric charges. The accelerating electrons explain not only the Maxwell Equations and the Special Relativity, but the Heisenberg Uncertainty Relation, the wave particle duality and the electron's spin also, building the bridge between the Classical and Relativistic Quantum Theories.

The Big Bang caused acceleration created the radial currents of the matter and since the matter composed of negative and positive charges, these currents are creating magnetic field and attracting forces between the parallel moving electric currents. This is the gravitational force experienced by the matter, and also the mass is result of the electromagnetic forces between the charged particles. The positive and negative charged currents attracts each other or by the magnetic forces or by the much stronger electrostatic forces. The gravitational force attracting the matter, causing concentration of the matter in a small space and leaving much space with low matter concentration: dark matter and energy.

Contents

Scientists reveal new connections between small particles and the vast universe	2
No Universe without Big Bang.....	3
No Big Bang? Quantum equation predicts universe has no beginning	4
Old ideas revisited	5
No singularities, nor dark stuff.....	5

New gravity particle	5
The Big Bang	5
Evidence for an accelerating universe	6
Equation	7
Explanatory models.....	8
Lorentz transformation of the Special Relativity	8
The Classical Relativistic effect	8
Electromagnetic inertia and Gravitational attraction	9
Electromagnetic inertia and mass.....	9
Electromagnetic Induction	9
Relativistic change of mass.....	9
The frequency dependence of mass	9
Electron – Proton mass rate	10
Gravity from the point of view of quantum physics	10
The Gravitational force	10
The Graviton	11
Conclusions	11
References	12

Scientists reveal new connections between small particles and the vast universe

Our observable universe is the largest object that physicists study: It spans a diameter of almost 100 billion light years. The density correlations in our universe, for example, correlations between numbers of galaxies at different parts of the universe, indicate that our vast universe has originated from a stage of cosmic inflation.

On the other hand, elementary particles are the smallest object that physicists study. A particle physics Standard Model (SM) was established 50 years ago, describing all known particles and their interactions.

Are density distributions of the vast universe and the nature of smallest particles related? In a recent research, scientists from HKUST and Harvard University revealed the connection between those two aspects, and argued that our universe could be used as a particle physics "collider" to study the high energy particle physics. Their findings mark the first step of cosmological collider phenomenology and pave the way for future discovery of new physics unknown yet to mankind.

The research was published in the journal *Physical Review Letters* on June 29, 2017 and the preprint is available online.

"Ongoing observations of cosmological microwave background and large scale structures have achieved impressive precision, from which valuable information about primordial density perturbations can be extracted," said Yi Wang, a co-author of the paper and an assistant professor at HKUST's department of physics. "A careful study of this SM background would be the prerequisite for using the cosmological collider to explore any new physics, and any observational signal that deviates from this background would then be a sign of physics beyond the SM."

The team carried out a two-step task to work out the background of the SM model. The first step was to work out the SM spectrum during inflation, which turned out to be dramatically different from that obtained from the particle physics calculation in flat space. The second one was to figure out how the SM fields entered the cosmological density correlation functions.

"Just like the line pattern of the light you see when observing a mercury lamp through a spectrometer, the mass distribution of the fundamental particles in SM also presents a special pattern, or a 'mass spectrum', which can be viewed as the fingerprint of SM," explained Zhong-Zhi Xianyu, a co-author and physicist at Center for Mathematical Sciences and Applications in Harvard University, "However, this fingerprint is subject to change if we change the ambient conditions. Just like the light spectrum changes when applying strong magnetic field to the lamp, the spectrum of the SM particles turns out to be very different at the time of inflation from it is now due to the inflationary background." The team carefully examined all effects from inflation and showed how the mass spectrum of SM would look like for different inflation models.

"Through inflation, the spectrum of elementary particles is encoded in the statistics of the distribution of the contents of the universe, such as the galaxies and cosmic microwave background, that we observe today", explains Xingang Chen, a co-author and scientist in the Harvard-Smithsonian Center for Astrophysics. "This is the connection between the smallest and largest."

Many problems along this direction remain to be explored. "In our minimal setup, the Standard Model particles interact with the inflaton (the driving force of inflation) rather weakly. But if some new particles can mediate stronger interactions between these two sectors, we would expect to observe a stronger signal of new physics," said Wang. "The cosmological collider is an ideal arena for new physics beyond SM." [8]

No Universe without Big Bang

According to Einstein's theory of relativity, the curvature of spacetime was infinite at the big bang. In fact, at this point all mathematical tools fail, and the theory breaks down. However, there remained the notion that perhaps the beginning of the universe could be treated in a simpler manner, and that the infinities of the big bang might be avoided. This has indeed been the hope expressed since the 1980s by the well-known cosmologists James Hartle and Stephen Hawking with their "no-boundary proposal", and by Alexander Vilenkin with his "tunnelling proposal". Now scientists at the Max Planck Institute for Gravitational Physics (Albert Einstein Institute/AEI) in Potsdam and at the Perimeter Institute in Canada have been able to use better mathematical methods to show that these ideas cannot work. The big bang, in its complicated glory, retains all its mystery.

One of the principal goals of cosmology is to understand the beginning of our universe. Data from the Planck satellite mission shows that 13.8 billion years ago the universe consisted of a hot and dense soup of particles. Since then the universe has been expanding. This is the main tenet of the hot big bang theory, but the theory fails to describe the very first stages themselves, as the conditions were too extreme. Indeed, as we approach the big bang, the energy density and the curvature grow until we reach the point where they become infinite.

As an alternative, the "no-boundary" and "tunneling" proposals assume that the tiny early universe arose by quantum tunnelling from nothing, and subsequently grew into the large universe that we see. The curvature of spacetime would have been large, but finite in this beginning stage, and the geometry would have been smooth - without boundary (see Fig. 1, left panel). This initial configuration would replace the standard big bang. However, for a long time the true consequences of this hypothesis remained unclear. Now, with the help of better mathematical methods, Jean-Luc Lehners, group leader at the AEI, and his colleagues Job Feldbrugge and Neil Turok at Perimeter Institute, managed to define the 35 year old theories in a precise manner for the first time, and to calculate their implications. The result of these investigations is that these alternatives to the big bang are no true alternatives. As a result of Heisenberg's uncertainty relation, these models do not only imply that smooth universes can tunnel out of nothing, but also irregular universes. In fact, the more irregular and crumpled they are, the more likely (see Fig. 1, right panel). "Hence the "no-boundary proposal" does not imply a large universe like the one we live in, but rather tiny curved universes that would collapse immediately", says Jean-Luc Lehners, who leads the "theoretical cosmology" group at the AEI.

Hence one cannot circumvent the big bang so easily. Lehners and his colleagues are now trying to figure out what mechanism could have kept those large quantum fluctuations in check under the most extreme circumstances, allowing our large universe to unfold. [7]

No Big Bang? Quantum equation predicts universe has no beginning

The widely accepted age of the universe, as estimated by general relativity, is 13.8 billion years. In the beginning, everything in existence is thought to have occupied a single infinitely dense point, or singularity. Only after this point began to expand in a "Big Bang" did the universe officially begin.

Although the Big Bang singularity arises directly and unavoidably from the mathematics of general relativity, some scientists see it as problematic because the math can explain only what happened immediately after—not at or before—the singularity.

"The Big Bang singularity is the most serious problem of general relativity because the laws of physics appear to break down there," Ahmed Farag Ali at Benha University and the Zewail City of Science and Technology, both in Egypt, told Phys.org.

Ali and coauthor Saurya Das at the University of Lethbridge in Alberta, Canada, have shown in a paper published in Physics Letters B that the Big Bang singularity can be resolved by their new model in which the universe has no beginning and no end.

Old ideas revisited

The physicists emphasize that their quantum correction terms are not applied ad hoc in an attempt to specifically eliminate the Big Bang singularity. Their work is based on ideas by the theoretical physicist David Bohm, who is also known for his contributions to the philosophy of physics. Starting in the 1950s, Bohm explored replacing classical geodesics (the shortest path between two points on a curved surface) with quantum trajectories.

Using the quantum-corrected Raychaudhuri equation, Ali and Das derived quantum-corrected Friedmann equations, which describe the expansion and evolution of universe (including the Big Bang) within the context of general relativity. Although it's not a true theory of quantum gravity, the model does contain elements from both quantum theory and general relativity. Ali and Das also expect their results to hold even if and when a full theory of quantum gravity is formulated.

No singularities, nor dark stuff

In addition to not predicting a Big Bang singularity, the new model does not predict a "big crunch" singularity, either. In general relativity, one possible fate of the universe is that it starts to shrink until it collapses in on itself in a big crunch and becomes an infinitely dense point once again.

Ali and Das explain in their paper that their model avoids singularities because of a key difference between classical geodesics and Bohmian trajectories. Classical geodesics eventually cross each other, and the points at which they converge are singularities. In contrast, Bohmian trajectories never cross each other, so singularities do not appear in the equations.

In cosmological terms, the scientists explain that the quantum corrections can be thought of as a cosmological constant term (without the need for dark energy) and a radiation term. These terms keep the universe at a finite size, and therefore give it an infinite age. The terms also make predictions that agree closely with current observations of the cosmological constant and density of the universe.

New gravity particle

In physical terms, the model describes the universe as being filled with a quantum fluid. The scientists propose that this fluid might be composed of gravitons—hypothetical massless particles that mediate the force of gravity. If they exist, gravitons are thought to play a key role in a theory of quantum gravity.

In a related paper, Das and another collaborator, Rajat Bhaduri of McMaster University, Canada, have lent further credence to this model. They show that gravitons can form a Bose-Einstein condensate (named after Einstein and another Indian physicist, Satyendranath Bose) at temperatures that were present in the universe at all epochs. [6]

The Big Bang

The Big Bang caused acceleration created radial currents of the matter, and since the matter is composed of negative and positive charges, these currents are creating magnetic field and attracting forces between the parallel moving electric currents. This is the gravitational force experienced by the matter, and also the mass is result of the electromagnetic forces between the charged particles.

The positive and negative charged currents attracts each other or by the magnetic forces or by the much stronger electrostatic forces!?


The gravitational force attracting the matter, causing concentration of the matter in a small space and leaving much space with low matter concentration: dark matter and energy.

There is an asymmetry between the mass of the electric charges, for example proton and electron, can be understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

Evidence for an accelerating universe

One of the observational foundations for the big bang model of cosmology was the observed expansion of the universe. [4] Measurement of the expansion rate is a critical part of the study, and it has been found that the expansion rate is very nearly "flat". That is, the universe is very close to the critical density, above which it would slow down and collapse inward toward a future "big crunch". One of the great challenges of astronomy and astrophysics is distance measurement over the vast distances of the universe. Since the 1990s it has become apparent that type Ia supernovae offer a unique opportunity for the consistent measurement of distance out to perhaps 1000 Mpc. Measurement at these great distances provided the first data to suggest that the expansion rate of the universe is actually accelerating. That acceleration implies an energy density that acts in opposition to gravity which would cause the expansion to accelerate. This is an energy density which we have not directly detected observationally and it has been given the name "dark energy".

The type Ia supernova evidence for an accelerated universe has been discussed by Perlmutter and the diagram below follows his illustration in Physics Today.



The data summarized in the illustration above involve the measurement of the redshifts of the distant supernovae. The observed magnitudes are plotted against the redshift parameter z . Note that there are a number of Type Ia supernovae around $z=0.6$, which with a Hubble constant of 71 km/s/mpc is a distance of about 5 billion light years.

Equation

The cosmological constant Λ appears in Einstein's field equation [5] in the form of

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu},$$

where R and g describe the structure of spacetime, T pertains to matter and energy affecting that structure, and G and c are conversion factors that arise from using traditional units of measurement. When Λ is zero, this reduces to the original field equation of general relativity. When T is zero, the field equation describes empty space (the vacuum).

The cosmological constant has the same effect as an intrinsic energy density of the vacuum, ρ_{vac} (and an associated pressure). In this context it is commonly moved onto the right-hand side of the equation, and defined with a proportionality factor of 8π : $\Lambda = 8\pi\rho_{\text{vac}}$, where unit conventions of general relativity are used (otherwise factors of G and c would also appear). It is common to quote values of energy density directly, though still using the name "cosmological constant".

A positive vacuum energy density resulting from a cosmological constant implies a negative pressure, and vice versa. If the energy density is positive, the associated negative pressure will drive an accelerated expansion of the universe, as observed. (See dark energy and cosmic inflation for details.)

Explanatory models

Models attempting to explain accelerating expansion include some form of dark energy, dark fluid or phantom energy. The most important property of dark energy is that it has negative pressure which is distributed relatively homogeneously in space. The simplest explanation for dark energy is that it is a cosmological constant or vacuum energy; this leads to the Lambda-CDM model, which is generally known as the Standard Model of Cosmology as of 2003-2013, since it is the simplest model in good agreement with a variety of recent observations.

Lorentz transformation of the Special Relativity

In the referential frame of the accelerating electrons the charge density lowering linearly because of the linearly growing way they takes every next time period. From the referential frame of the wire there is a parabolic charge density lowering.

The difference between these two referential frames, namely the referential frame of the wire and the referential frame of the moving electrons gives the relativistic effect. Important to say that the moving electrons presenting the time coordinate, since the electrons are taking linearly increasing way every next time period, and the wire presenting the geometric coordinate. The Lorentz transformations are based on moving light sources of the Michelson - Morley experiment giving a practical method to transform time and geometric coordinates without explaining the source of this mystery.

The real mystery is that the accelerating charges are maintaining the accelerating force with their charge distribution locally. The resolution of this mystery that the charges are simply the results of the diffraction patterns, that is the charges and the electric field are two sides of the same thing. Otherwise the charges could exceed the velocity of the electromagnetic field.

The increasing mass of the electric charges the result of the increasing inductive electric force acting against the accelerating force. The decreasing mass of the decreasing acceleration is the result of the inductive electric force acting against the decreasing force. This is the relativistic mass change explanation, especially importantly explaining the mass reduction in case of velocity decrease.

The Classical Relativistic effect

The moving charges are self maintain the electromagnetic field locally, causing their movement and this is the result of their acceleration under the force of this field.

In the classical physics the charges will distributed along the electric current so that the electric potential lowering along the current, by linearly increasing the way they take every next time period because this accelerated motion.

Electromagnetic inertia and Gravitational attraction

Since the magnetic induction creates a negative electric field as a result of the changing acceleration, it works as an electromagnetic inertia, causing an electromagnetic mass.

It looks clear that the growing acceleration results the relativistic growing mass - limited also with the velocity of the electromagnetic wave.

Since $E = h\nu$ and $E = mc^2$, $m = h\nu/c^2$ that is the m depends only on the ν frequency. It means that the mass of the proton and electron are electromagnetic and the result of the electromagnetic induction, caused by the changing acceleration of the spinning and moving charge! It could be that the m_0 inertial mass is the result of the spin, since this is the only accelerating motion of the electric charge. Since the accelerating motion has different frequency for the electron in the atom and the proton, they masses are different, also as the wavelengths on both sides of the diffraction pattern, giving equal intensity of radiation.

If the mass is electromagnetic, then the gravitation is also electromagnetic effect caused by the accelerating Universe! The same charges would attract each other if they are moving parallel by the magnetic effect.

The Planck distribution law explains the different frequencies of the proton and electron, giving equal intensity to different lambda wavelengths! Also since the particles are diffraction patterns they have some closeness to each other – can be seen as a gravitational force.

Electromagnetic inertia and mass

Electromagnetic Induction

Since the magnetic induction creates a negative electric field as a result of the changing acceleration, it works as an electromagnetic inertia, causing an electromagnetic mass. [1]

Relativistic change of mass

The increasing mass of the electric charges the result of the increasing inductive electric force acting against the accelerating force. The decreasing mass of the decreasing acceleration is the result of the inductive electric force acting against the decreasing force. This is the relativistic mass change explanation, especially importantly explaining the mass reduction in case of velocity decrease.

The frequency dependence of mass

Since $E = h\nu$ and $E = mc^2$, $m = h\nu/c^2$ that is the m depends only on the ν frequency. It means that the mass of the proton and electron are electromagnetic and the result of the electromagnetic induction, caused by the changing acceleration of the spinning and moving charge! It could be that the m_0 inertial mass is the result of the spin, since this is the only accelerating motion of the electric charge. Since the accelerating motion has different frequency for the electron in the atom and the proton, they masses are different, also as the wavelengths on both sides of the diffraction pattern, giving equal intensity of radiation.

Electron – Proton mass rate

The Planck distribution law explains the different frequencies of the proton and electron, giving equal intensity to different lambda wavelengths! Also since the particles are diffraction patterns they have some closeness to each other – can be seen as a gravitational force. [1]

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

Gravity from the point of view of quantum physics

The Gravitational force

The gravitational attractive force is basically a magnetic force.

The same electric charges can attract one another by the magnetic force if they are moving parallel in the same direction. Since the electrically neutral matter is composed of negative and positive charges they need 2 photons to mediate this attractive force, one per charges. The Big Bang caused parallel moving of the matter gives this magnetic force, experienced as gravitational force.

Since graviton is a tensor field, it has spin = 2, could be 2 photons with spin = 1 together.

You can think about photons as virtual electron – positron pairs, obtaining the necessary virtual mass for gravity.

The mass as seen before a result of the diffraction, for example the proton – electron mass rate $M_p=1840 m_e$. In order to move one of these diffraction maximum (electron or proton) we need to intervene into the diffraction pattern with a force appropriate to the intensity of this diffraction maximum, means its intensity or mass.

The Big Bang caused acceleration created radial currents of the matter, and since the matter is composed of negative and positive charges, these currents are creating magnetic field and attracting forces between the parallel moving electric currents. This is the gravitational force experienced by the matter, and also the mass is result of the electromagnetic forces between the charged particles. The positive and negative charged currents attracts each other or by the magnetic forces or by the much stronger electrostatic forces!?

The gravitational force attracting the matter, causing concentration of the matter in a small space and leaving much space with low matter concentration: dark matter and energy.

There is an asymmetry between the mass of the electric charges, for example proton and electron, can understood by the asymmetrical Planck Distribution Law. This temperature dependent energy distribution is asymmetric around the maximum intensity, where the annihilation of matter and antimatter is a high probability event. The asymmetric sides are creating different frequencies of electromagnetic radiations being in the same intensity level and compensating each other. One of these compensating ratios is the electron – proton mass ratio. The lower energy side has no compensating intensity level, it is the dark energy and the corresponding matter is the dark matter.

The Graviton

In physics, the graviton is a hypothetical elementary particle that mediates the force of gravitation in the framework of quantum field theory. If it exists, the graviton is expected to be massless (because the gravitational force appears to have unlimited range) and must be a spin-2 boson. The spin follows from the fact that the source of gravitation is the stress-energy tensor, a second-rank tensor (compared to electromagnetism's spin-1 photon, the source of which is the four-current, a first-rank tensor). Additionally, it can be shown that any massless spin-2 field would give rise to a force indistinguishable from gravitation, because a massless spin-2 field must couple to (interact with) the stress-energy tensor in the same way that the gravitational field does. This result suggests that, if a massless spin-2 particle is discovered, it must be the graviton, so that the only experimental verification needed for the graviton may simply be the discovery of a massless spin-2 particle. [2]

Conclusions

Motivated by the model's potential to resolve the Big Bang singularity and account for dark matter and dark energy, the physicists plan to analyze their model more rigorously in the future. Their future work includes redoing their study while taking into account small inhomogeneous and anisotropic perturbations, but they do not expect small perturbations to significantly affect the results. "It is satisfying to note that such straightforward corrections can potentially resolve so many issues at once," Das said. [6]

The accelerating Universe fits into the accelerating charges of the electric currents, because the Big Bang caused radial moving of the matter.

Needless to say that the accelerating electrons of the steady stationary current are a simple demystification of the magnetic field, by creating a decreasing charge distribution along the wire, maintaining the decreasing U potential and creating the \mathbf{A} vector potential experienced by the electrons moving by \mathbf{v} velocity relative to the wire. This way it is easier to understand also the time dependent changes of the electric current and the electromagnetic waves as the resulting fields moving by c velocity.

It could be possible something very important law of the nature behind the self maintaining \mathbf{E} accelerating force by the accelerated electrons. The accelerated electrons created electromagnetic fields are so natural that they occur as electromagnetic waves traveling with velocity c . It shows that the electric charges are the result of the electromagnetic waves diffraction.

One of the most important conclusions is that the electric charges are moving in an accelerated way and even if their velocity is constant, they have an intrinsic acceleration anyway, the so called spin, since they need at least an intrinsic acceleration to make possible their movement.

The bridge between the classical and quantum theory is based on this intrinsic acceleration of the spin, explaining also the Heisenberg Uncertainty Principle. The particle – wave duality of the electric charges and the photon makes certain that they are both sides of the same thing. Basing the gravitational force on the accelerating Universe caused magnetic force and the Planck Distribution Law of the electromagnetic waves caused diffraction gives us the basis to build a Unified Theory of the physical interactions.

The electric currents causing self maintaining electric potential is the source of the special and general relativistic effects. The Higgs Field is the result of the electromagnetic induction. The Graviton is two photons together. [3]

References

[1] 3 Dimensional String Theory

Author: George Rajna

Publisher: Academia.edu

http://www.academia.edu/3834454/3_Dimensional_String_Theory

[2] Graviton Production By Two Photon and Electron-Photon Processes In Kaluza-Klein Theories With Large Extra Dimensions

<http://arxiv.org/abs/hep-ph/9909392>

[3] http://www.academia.edu/4158863/Higgs_Field_and_Quantum_Gravity

[4] <http://hyperphysics.phy-astr.gsu.edu/hbase/astro/univacc.html>

[5] http://en.wikipedia.org/wiki/Einstein_field_equations

[6] No Big Bang? Quantum equation predicts universe has no beginning

<http://phys.org/news/2015-02-big-quantum-equation-universe.html>

[7] No Universe without Big Bang

<https://phys.org/news/2017-06-universe-big.html>

[8] Scientists reveal new connections between small particles and the vast universe

<https://phys.org/news/2017-07-scientists-reveal-small-particles-vast.html>