

The Average Computed In Primes Basis {File Closing Version 1}. ISSN 1751-3030

Author:

Ramesh Chandra Bagadi

Data Scientist

INSOFE (International School Of Engineering),

Hyderabad, India.

rameshcbagadi@uwalumni.com

+91 9440032711

Research Manuscript

Abstract

In this research investigation, the author has detailed a novel method of finding the average of a sequence in Primes Basis.

Theory

Given any Sequence of the kind,

$$S = \{y_1, y_2, y_3, \dots, y_{n-1}, y_n\}$$

We first write them as

$$S = \{y_1 = {}^N P_{j_1 + \delta_1}, y_2 = {}^N P_{j_2 + \delta_2}, y_3 = {}^N P_{j_3 + \delta_3}, \dots, y_{n-1} = {}^N P_{j_{n-1} + \delta_{n-1}}, y_n = {}^N P_{j_n + \delta_n}\}$$

where in ${}^N P_{j_1 + \delta_1}$, N is the Order Number of the Higher Order Sequence Of Primes in which the number y_1 is slated, $(j_1 + \delta_1)$ is the position number of the Prime Metric Basis Element.

For Example,

7 which is the 4th Prime Metric Basis Element of the Standard Primes, the Order [1] of which can be taken to 1. Therefore 7 can be written as ${}^1 P_4$. In a similar fashion, 8 can be written as ${}^1 P_{4 + \left(\frac{8-7}{11-7}\right)}$ where 7 is the nearest previous prime number of 8 and 11 is the next nearest prime number of 8. Here, in the notation ${}^1 P_{4 + \left(\frac{8-7}{11-7}\right)}$, we can consider $\left(\frac{8-7}{11-7}\right)$ as the δ , the 4 as the j and the 1 as

the N . We can also denote any number in a similar fashion using Higher Order Primes as well. [1] i.e., $N > 1$.

We then compute the sum $\left(\frac{\sum_{i=1}^n (j_i + \delta_i)}{n} \right)$. Let this be $(k + \beta)$. We then use [2] to

evolve $(k + \beta)$ by one step. Now, we find ${}^N p_{E^1(k+\beta)}$ where $E^1(k + \beta)$ is one step evolved $(k + \beta)$. This ${}^N p_{E^1(k+\beta)}$ can be used as the Average Computed in the Primes Basis.

Example.

Considering, $S = \{2, 4, 6\}$, we note that they are actually, ${}^{N=1} p_1$, ${}^{N=1} p_{1+\left(\frac{4-3}{5-3}\right)}$ and

${}^{N=1} p_{3+\left(\frac{6-5}{7-5}\right)}$. Therefore, $\left(\frac{\sum_{i=1}^n (j_i + \delta_i)}{n} \right) = 2$.

The number 2 when evolved using [2] gives 3. Hence the average is 5.

References

1. Bagadi, R. (2016). Field(s) Of Sequence(s) Of Primes Of Positive Integral Higher Order Space(s). *PHILICA.COM Article number 622*.
http://philica.com/display_article.php?article_id=622
2. Bagadi, R. (2017). One Step Evolution Of Any Positive Real Number. ISSN 1751-3030. *PHILICA.COM Article number 1106*.
http://philica.com/display_article.php?article_id=1106
3. <http://www.philica.com/advancedsearch.php?author=12897>
4. http://www.vixra.org/author/ramesh_chandra_bagadi