

Scheme For Finding The Next Term Of A Sequence Based On Evolution {File Closing Version 4}. ISSN 1751-3030

Author:

Ramesh Chandra Bagadi

Data Scientist

INSOFE (International School Of Engineering),

Hyderabad, India.

rameshcbagadi@uwalumni.com

+91 9440032711

Research Manuscript

Abstract

In this research investigation, the author has detailed a novel method of finding the next term of a sequence based on Evolution.

Theory

Given any Sequence of the kind,

$$S = \{y_1, y_2, y_3, \dots, y_{n-1}, y_n\}$$

We first write them as

$$S = \left\{ \begin{array}{l} y_1 = {}^N P_{j_1 + \delta_1}, y_2 = {}^N P_{j_2 + \delta_2}, y_3 = {}^N P_{j_3 + \delta_3}, \dots, y_{n-1} = {}^N P_{j_{n-1} + \delta_{n-1}}, \\ y_n = {}^N P_{j_n + \delta_n} \end{array} \right\}$$

where in ${}^N P_{j_1 + \delta_1}$, N is the Order Number of the Higher Order Sequence Of Primes in which the number y_1 is slated,

$(j_i + \delta_i)$ is the position number of the Prime Metric Basis Element. Here, j_i 's are Positive Integers and $0 < \delta_i < 1$.

For Example,

7 which is the 4th Prime Metric Basis Element of the Standard Primes, the Order [1] of which can be taken to 1. Therefore 7 can be written as 1p_4 . In a similar fashion, 8 can be written as

${}^1P_{4+\left(\frac{8-7}{11-7}\right)}$ where 7 is the nearest previous prime number of 8 and 11 is the next nearest prime number of 8. Here, in the

notation ${}^1P_{4+\left(\frac{8-7}{11-7}\right)}$, we can consider $\left(\frac{8-7}{11-7}\right)$ as the δ , the 4 as

the j and the 1 as the N . We can also denote any number in a similar fashion using Higher Order Primes as well. [1] i.e., $N > 1$.

Given $(j_i + \delta_i)$, a method of calculating the Decimal (Pseudo)

Prime corresponding to $(j_i + \delta_i)$ in ${}^N P_{(j_i + \delta_i)}$.

If δ_i is equal to $\left(\frac{a_1 a_2 a_3 \dots a_{k-1} a_k}{10^k}\right)$ where $0 < a_l < 10$ for $l = 1$ to k , we write

$${}^N P_{(j_i+\delta_i)} = {}^N P_{j_i} + \left\{ \frac{\left(a_1 a_2 a_3 \dots a_{k-1} a_k \right)^{th} \text{ PrimeNumber}}{\left(10^k \right)^{th} \text{ PrimeNumber}} \right\} \left\{ {}^N P_{j_i+1} - {}^N P_{j_i} \right\}$$

of N^{th} Order

of N^{th} Order

Given $\binom{N}{P}$, a method of calculating the Decimal (Pseudo) Position $(j_i + \delta_i)$, i.e., the Prime Metric Basis Element Position corresponding to $\binom{N}{P}$ in the Sequence of N^{th} Order Sequence Of Primes.

We write the given number (Real) say $\frac{a}{b}$ as

$\frac{a}{b} = {}^N P_{j_i} + \frac{c}{d}$ where c, d are some N^{th} Order Primes such that the equation holds roughly.

We then find the positions of c, d along the N^{th} Order Sequence of Primes, say they are l, m .

We then write the Position of $\binom{N}{P}$ as

$$j_i + \delta_i =$$

Prime Basis Position of (c) in the N^{th} Order

$$j_i + \frac{\text{Sequence of Primes}}{\text{Prime Basis Position of (d) in the } N^{th} \text{ Order}} \\ \text{Sequence of Primes}$$

For Simplicity, we can take $N = 1$.

For our representational simplicity, we label our $S = \{y_1, y_2, y_3, \dots, y_{n-1}, y_n\}$ as

$S = \{ {}_1y_1, {}_1y_2, {}_1y_3, \dots, {}_1y_{n-1}, {}_1y_n \}$ where the left subscript 1 indicates that these numbers are at the level 1 (Base) of the triangle we are going to build.

We now compute the Evolution Orders

$E^{(j+1)y_i} \left({}_jy_i \right) = {}_jy_{(i+1)}$ where $(j+1)y_i$ is the Evolution Order.

By Evolution Order, we mean the difference between the

Prime Basis Position Number of ${}_jy_{(i+1)}$ (when slated

thusly) and the Prime Basis Position Number of ${}_jy_i$ given

that we are considering this in $N = 1$. This means that ${}_jy_i$

needs to be evolved $(j+1)y_i$ times to get ${}_jy_{(i+1)}$. Please see

[2] for Evolution method. We repeat this process till we get

$E^{(n)y_i} \left({}_{(n-1)}y_i \right) = {}_{(n-1)}y_{(i+1)}$. We now evolve ${}_{(n-1)}y_{(i+1)}$ by one

step using [2] and get $E^1\left(\binom{(n-1)y_{(i+1)}}{(n-1)}\right)$. Note that $E^{\left\{E^1\left(\binom{(n-1)y_{(i+1)}}{(n-1)}\right)\right\}}\left(\binom{(n-2)y_i}{(n-2)}\right) = \binom{(n-2)y_{(i+1)}}{(n-2)}$.

We repeat this procedure, downwards, repeatedly to find $E^{\left\{(2)y_n\right\}}\left(\binom{(1)y_n}{(1)}\right) = \binom{(1)y_{(n+1)}}{(1)}$.

We illustrate this with an Example of three terms.

Considering

$S = \{y_1, y_2, y_3\}$ which we write as

$S = \left\{\binom{1y_1, 1y_2, 1y_3}{1}\right\}$ for future representational simplicity.

$E^{2y_1}\left(\binom{1y_1}{1}\right) = \binom{1y_2}{1}$, $E^{2y_2}\left(\binom{1y_2}{1}\right) = \binom{1y_3}{1}$. Now, we write

$E^{3y_1}\left(\binom{2y_1}{2}\right) = \binom{2y_2}{2}$. We now evolve $\binom{3y_1}{3}$ by one step using [2], i.e., perform $E^1\left(\binom{3y_1}{3}\right)$. And now, we write $E^{\left\{E^1\left(\binom{3y_1}{3}\right)\right\}}\left(\binom{2y_2}{2}\right) = \binom{2y_3}{2}$. Finally, we write, $E^{\left(2y_3\right)}\left(\binom{1y_3}{1}\right) = \binom{1y_4}{1} = y_4$ which is the next term of the sequence $S = \{y_1, y_2, y_3\}$. This method can be used advantageously for forecasting.

Note that $E^1(0) = 0$ and $E^1(1) = E^1\left(\frac{2}{2}\right) = \frac{3}{2}$, from [2].

References

1. Bagadi, R. (2016). Field(s) Of Sequence(s) Of Primes Of Positive Integral Higher Order Space(s). *PHILICA.COM Article number 622*.

Bagadi, R. (2017). Scheme For Finding The Next Term Of A Sequence Based On Evolution {File Closing Version 3}. ISSN 1751-3030. *PHILICA.COM Article number 1140*.
http://www.philica.com/display_article.php?article_id=1140

http://philica.com/display_article.php?article_id=622

2. Bagadi, R. (2017). One Step Evolution Of Any Positive Real Number. ISSN 1751-3030. *PHILICA.COM Article number 1106*.

http://philica.com/display_article.php?article_id=1106

3. <http://www.philica.com/advancedsearch.php?author=12897>

4. http://www.vixra.org/author/ramesh_chandra_bagadi