

Newton's Gravity from Heisenberg's Uncertainty Principle An In-Depth Study of the McCulloch Derivation

Espen Gaarder Haug*
Norwegian University of Life Sciences

September 6, 2018

Abstract

Mike McCulloch has derived Newton's gravity from Heisenberg's uncertainty principle in an innovative and interesting way. Upon deeper examination, we will claim that his work has additional important implications, when viewed from a different perspective. Based on recent developments in mathematical atomism, particularly those exploring the nature of Planck masses and their link to Heisenberg's uncertainty principle, we uncover an insight on the quantum world that leads to an even more profound interpretation of the McCulloch derivation than was put forward previously.

Key words: Heisenberg's uncertainty principle, Newtonian gravity, Planck momentum, Planck mass, Planck length, gravitational constant, Lorentz symmetry.

1 Heisenberg's Uncertainty Principle, Newton's Theory of Gravitation, and McCulloch's Derivation

McCulloch has derived Newton's gravitational force from the Heisenberg uncertainty principle; see [1, 2]. At first, this might sound absurd, as Newtonian gravity not is assumed to be consistent with the quantum world and the Heisenberg principle was developed expressly for the subatomic quantum world. However, we observe that McCulloch is using the Planck mass to obtain his derivation and, as recent work by Haug strongly indicates, all observable gravitational phenomena may be dependent on the Planck mass, which is directly linked to the Planck length.

In this paper, we will examine the McCulloch derivation, based on our understanding of the Planck masses and the link to quantum gravity. Heisenberg's uncertainty principle [3] is given by

$$\Delta p \Delta x \geq \hbar \quad (1)$$

Next McCulloch goes on to say "Now $E = pc$ so" :

$$\Delta E \Delta x \geq \hbar c \quad (2)$$

However, he does not mention that $E = pc$ only can hold for the Planck mass momentum. Further, we claim that it can only hold for a Planck mass particle. While the Planck mass, Planck length, Planck time, and Planck energy were introduced in 1899 by Max Planck himself [4, 5], the Planck mass particle was likely first introduced by Lloyd Motz, when he was working at the Rutherford Laboratory in 1962; see [6, 7, 8].

The momentum formula that holds for any particle is the relativistic momentum formula given by Einstein [10]

$$p = \frac{mv}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (3)$$

Multiplying this by c clearly does not give us the kinetic energy formula or the rest-mass energy formula. So, how does the Planck momentum fit in here? As shown by Haug [9], the Planck mass is truly unique and this means the McCulloch derivation likely holds. Our point is that the way McCulloch reaches his relation $\Delta E \Delta x \geq \hbar c$ likely can only be done with regard to Planck mass particles, or at least Planck masses. This is also essential for understanding gravity at a quantum level. The Planck mass particle is, in our view, the building

*e-mail espenhaug@mac.com. Thanks to Victoria Terces for helping me edit this manuscript, and also thanks to Doug Danforth for finding an important typo.

block of all other particles, an idea first suggested by Motz. The Planck mass seems to be much too large for this to be possible, but if it only lasts for one Planck second, then this can be shown to cause no problems. In previous work, we have claimed that indeed the Planck mass particle does only last for one Planck second, after which it goes from mass to pure energy. Looking at this scenario, we also get a hint from the Planck acceleration, $a_p = \frac{c^2}{l_p} \approx 5.56092 \times 10^{51} \text{ m/s}^2$. The Planck acceleration is assumed to be the maximum possible acceleration by some physicists; see [11, 12], for example. The velocity of a particle that undergoes Planck acceleration will actually reach the speed of light within one Planck second: $a_p t_p = \frac{c^2 l_p}{l_p c} = c$. However, we know that nothing with rest-mass can travel at the speed of light, thus no “normal” particle can undergo Planck acceleration if the shortest possible acceleration time interval is the Planck second. The solution is simple. The Planck acceleration is an internal acceleration inside the Planck particle that within one Planck second turns the Planck mass particle into pure energy. This also explains why the Planck momentum is so special, namely always $m_p c$, unlike for any other particles, which can take a wide range of velocities and therefore a wide range of momentums. The Planck mass can be described as

$$m_p = \frac{\hbar}{l_p} \frac{1}{c} \quad (4)$$

We naturally do not know the Planck mass exactly, just as we do not know the Planck length exactly; this is something we will discuss later in this paper. Yet, the more precisely we know the Planck length, the more precisely we will know the Planck momentum. From the union of this line of thought with Heisenberg’s uncertainty principle, we have claimed that the Planck mass momentum likely collapses into a certainty principle, as recently described by Haug [9, 13]:

$$m_p c l_p = \hbar \quad (5)$$

Further, from equation 2 McCulloch describes

$$\Delta E \Delta x = \sum_i^N \sum_j^n (\hbar c)_{i,j} \quad (6)$$

where \sum_i^N is the number of Planck masses in a smaller mass, so simply $\frac{m}{m_p}$ and \sum_j^n is the mass in a larger mass $\frac{M}{m_p}$.

This gives us the equation

$$\Delta E = \frac{\hbar c m M}{m_p^2 \Delta x} \quad (7)$$

However, unlike McCulloch, we will claim that there is no uncertainty in the energy here, because the particular energy in this case is the Planck mass energy. As explained previously, we claim that the Heisenberg uncertainty principle collapses at the Planck scale and this may be why McCulloch has been able to get gravity from the quantum scale. Again, McCulloch obtained ΔE from a momentum of the form pc , which can only apply to the Planck mass particle momentum.

Therefore, in our view we must have

$$\begin{aligned} \Delta E &= \frac{\hbar c m M}{m_p^2 \Delta x} \\ N n E_P &= \frac{\hbar c}{m_p^2} \frac{m M}{\Delta x} \\ N n m_p c^2 &= \frac{\hbar c}{m_p^2} \frac{m M}{\Delta x} \end{aligned} \quad (8)$$

This is only true if $\Delta x = l_p$. This can be linked to the Planck acceleration, $m_p a_p = m_p \frac{c^2}{l_p}$ and based on this we have

$$\begin{aligned} N n m_p a_p &= \frac{\hbar c}{m_p^2} \frac{m M}{(\Delta x)^2} \\ N n m_p \frac{c^2}{l_p} &= \frac{\hbar c}{m_p^2} \frac{m M}{(\Delta x)^2} \\ F &= \frac{\hbar c}{m_p^2} \frac{m M}{l_p^2} \end{aligned} \quad (9)$$

Be aware that $\Delta x = l_p$ is the minimum position uncertainty in the Heisenberg principle, according to strong evidence given by Haug [14]:

$$F = \frac{\hbar c}{m_p^2} \frac{mM}{r^2} \quad (10)$$

which is the same formula that McCulloch first derived and is very similar to Newton's gravitational formula, except that here we have $\frac{\hbar c}{m_p^2}$ (as first derived by McCulloch), instead of Newton's gravitational constant. Now pay attention to the fact that numerically

$$\frac{\hbar c}{m_p^2} \approx 6.67384 \times 10^{-11} \quad (11)$$

This is well inside the experimentally observed value of the gravitational constant. However, this is the gravitational constant derived from Heisenberg's uncertainty principle, which we have claimed becomes a certainty principle at the Planck scale and further, the Lorentz symmetry breaks down at the Planck scale.

The fact that gravity does not obey Lorentz symmetry is known, but seldom discussed, and is quite obvious even from simple gravitational time dilation experiments. If we run a clock experiment with someone on a mountain top and someone in a valley, they will agree that the clock on the mountain top is running faster than the one in the valley. This lies in contrast to special relativity, where Lorentz symmetry holds all the way up to our newly-introduced maximum velocity of matter, at which point we reach the Planck scale and Lorentz symmetry is broken.

Gravity has to do with the Planck mass particle that, in our view, makes up all matter and energy in the universe. Gravity has to do with the very collision points of indivisible particles inside the atoms themselves. In a series of recently published papers, [15, 16, 17, 18] Haug has suggested and shown strong evidence for the idea that Newton's gravitational constant must be a composite constant of the form

$$G = \frac{l_p^2 c^3}{\hbar} \quad (12)$$

This has been derived from dimensional analysis [15], but one can also derive it directly from the Planck length formula. McCulloch's Heisenberg-derived gravitational constant $G = \frac{\hbar c}{m_p^2}$ is naturally the same as the

Planck mass, and is directly linked to the Planck length, $m_p = \frac{\hbar}{l_p c}$.

Many physicists will likely protest here and claim that we cannot find the Planck mass before we know the Newton gravitational constant. However, Haug [17] has shown that one easily can measure the Planck length with a Cavendish apparatus without any knowledge of Newton's gravitational constant. And if we have the Planck length, then we can easily find the Planck mass. Further, Haug has shown that the standard uncertainty in the Planck length experiments must be exactly half of that of the standard uncertainty in the gravitational constant, which is likely a composite constant. See [19].

2 Other Comments on the McCulloch Paper in Relation to Newton's Gravity Derivation

Close to equation 6 in his paper [2], McCulloch asks, "What if we assume that the sum of the kinetic energy and the energy uncertainty is conserved?" and he provides the following equation

$$\frac{1}{2}m(\Delta v)^2 + \frac{\hbar c}{m_p^2} \frac{mM}{\Delta x} = constant \quad (13)$$

Let's examine this formulation in greater detail. In our framework, the $\frac{1}{2}$ should not be there in the first place, as Planck mass particles are unique and do not have standard kinetic energy. Instead, they have rest-mass energy that then bursts into pure energy. This is why the momentum of a Planck mass particle is $p = mc$, unlike for any other particle. Is there a way to unite this view though, with the revolutionary developments that are coming out from atomism? The same challenge has also arisen when deriving Newton's bending of light, where some authors have used the kinetic energy formula, in spite of the fact that they are working with photons that are naturally moving at the speed of light. A possible solution that gives a modified Newtonian quantum gravity theory and the right bending of light has recently been suggested by Haug; see [19].

We can claim the Planck mass particle has zero kinetic energy, or alternatively (and just as correct) that all of its rest-mass energy is kinetic energy, which is unlike any other particle. Also, we think there should be a minus in McCulloch's formula above, and that the constant in his formula is simply zero.

$$nNm_p c^2 - \frac{\hbar c}{m_p^2} \frac{mM}{l_p} = 0 \quad (14)$$

We think that McCulloch's hypothesis leads to something new and important, but that it requires further examination before it can be fully understood and adjusted if necessary. Then we can build a more solid framework around it.

Almost nothing revolutionary has happened in the study of gravity or theoretical physics since the time of Einstein. We think that recent breakthroughs in understanding the Planck units, combined with new use of Heisenberg's uncertainty principle, have just started to propel us to a new epoch of fundamental physics.

3 Conclusion

We have taken an in-depth look at McCulloch's interesting derivation of Newtonian gravity from Heisenberg's uncertainty principle. We have pointed out that that McCulloch has done this from Planck masses and the Planck momentum without reference to a transformation that we have hypothesized in our work, namely that Heisenberg's uncertainty principle seems to collapse and becomes the certainty principle at the Planck scale. This is important in understanding this new and interesting work and we are studying the viability of bringing the two ideas together.

Another important point is that all macroscopic observable gravity phenomena actually depend on the Planck mass or the Planck length – they are two sides of the same coin. McCulloch's theory supports our view that the gravitational constant must be a composite constant and that all gravitational phenomena depend on Planck-scale physics. An alternative view worthy of consideration is that the Heisenberg uncertainty principle become a certainty principle at the Planck scale and that gravity is dependent on the Planck scale. Therefore we can use the Heisenberg principle to derive Newton's gravity formula, with deeper implications for physics to come in the near future.

References

- [1] M. E. McCulloch. Gravity from the uncertainty principle. *Astrophysics and Space Science*, 349(2), 2014.
- [2] M. E. McCulloch. Quantised inertia from relativity and the uncertainty principle. *EPL (Europhysics Letters)*, 115(6), 2016.
- [3] W. Heisenberg. Über den anschaulichen inhalt der quantentheoretischen kinematik und mechanik. *Zeitschrift für Physik*, (43):172–198, 1927.
- [4] M. Planck. *Naturlische Masseinheiten*. Der Königlich Preussischen Akademie Der Wissenschaften, p . 479., 1899.
- [5] M. Planck. *Vorlesungen über die Theorie der Wärmestrahlung*. Leipzig: J.A. Barth, p. 163, see also the English translation “The Theory of Radiation” (1959) Dover, 1906.
- [6] L. Motz. Gauge invariance and the structure of charged particles. *Il Nuovo Cimento*, 26(4), 1962.
- [7] L. Motz. A gravitational theory of the mu meson and leptons in general. *Rutherford Observatory, Columbia University*, 1966.
- [8] L. Motz. The quantization of mass. *Rutherford Observatory, Columbia University*, 1971.
- [9] E. G. Haug. Revisiting the derivation of Heisenberg's uncertainty principle: The collapse of uncertainty at the Planck scale. *preprints.org*, 2018.
- [10] A. Einstein. On the electrodynamics of moving bodies. *Annalen der Physik, English translation by George Barker Jeffery 1923*, (17), 1905.
- [11] G. Scarpetta. *Letter Nuovo Cimento*,, 51, 1984.
- [12] D. F. Falla and P. T. Landsberg. Black holes and limits on some physical quantities. *European Journal of Physics*, 15, 1994.
- [13] E. G. Haug. Does Heisenberg's uncertainty collapse at the Planck scale? Heisenberg's uncertainty principle becomes the certainty principle. *www.viXra.org 1803.0038*, 2018.
- [14] E. G. Haug. Does Heisenberg's uncertainty principle predict a maximum velocity for anything with rest-mass below the speed of light ? *viXra:1801.0218*, 2018.
- [15] E. G. Haug. The gravitational constant and the Planck units. A simplification of the quantum realm. *Physics Essays Vol 29, No 4*, 2016.
- [16] E. G. Haug. Planck quantization of Newton and Einstein gravitation. *International Journal of Astronomy and Astrophysics*, 6(2), 2016.
- [17] E. G. Haug. Can the Planck length be found independent of big G ? *Applied Physics Research*, 9(6), 2017.
- [18] E. G. Haug. Newton and Einstein's gravity in a new perspective for Planck masses and smaller sized objects. *International Journal of Astronomy and Astrophysics*, 2018.
- [19] E. G. Haug. A simple Newtonian quantum gravity theory that predicts the same light bending as GR. And also a new gravitational prediction that can be tested! *www.viXra.org 1802.0169*, 2018.