

Enough of the trap of non-existent dark matter.

Angel Garcés Doz

March 30, 2018

angel1056510@gmail.com

Abstract

Since the experimental event of abnormal speeds within galaxies was discovered; the so-called dark matter is incessantly sought. But all the experiments still do not find such a ghost. Its justification is supported, in the apparent, simplest explanation: the existence of a matter additional to the baryonic matter within the galaxies. But there is another possible explanation more natural and just as simple: A modification of the gravity due to the occurrence of effects of quantum gravity produced by the repulsive acceleration of the vacuum. A repulsion of the vacuum would produce, as we demonstrated with the experiments with magnets of equal polarity, a pressure towards the center of mass and, therefore, a fictitious increase in mass. In this short article; we calculate the average speed of rotation within the newly discovered ultradiffuse galaxy NGC1052-DF2. This calculation shows that the equation, type MOND, agrees perfectly with the estimated velocities. Therefore, some voices that too quickly have announced that this galaxy (almost without dark matter or anything of it) does not agree with the theories of modified gravitation (we refer to the quantum gravitation that would quantize the equations of Einstein's gravitation, or GR); they are simply avoiding the real explanation: effects of quantum gravity produced by the vacuum (very large masses of galaxies).

Introduction.

We will refer directly to the summary of the article published in nature and in arxiv, as the best way to start exposing the experimental finding; to then apply our theoretical equation that can be found theoretically deduced in the bibliography of this article (bibliography of some of our previous articles).

“Studies of galaxy surveys in the context of the cold dark matter paradigm have shown that the mass of the dark matter halo and the total stellar mass are coupled through a function that varies smoothly

with mass. Their average ratio $M_{\text{halo}}/M_{\text{stars}}$ has a minimum of about 30 for galaxies with stellar masses near that of the Milky Way (approximately 5×10^{10} solar masses) and increases both towards lower masses and towards higher masses 1,2. The scatter in this relation is not well known; it is generally thought to be less than a factor of two for massive galaxies but much larger for dwarf galaxies 3,4. Here we report the radial velocities of ten luminous globular-cluster-like objects in the ultra-diffuse galaxy 5 NGC1052–DF2, which has a stellar mass of approximately 2×10^8 solar masses. **We infer that its velocity dispersion is less than 10.5 kilometres per second with 90 per cent confidence**, and we determine from this that its total mass within a radius of 7.6 kiloparsecs is less than 3.4×10^8 solar masses. This implies that the ratio $M_{\text{halo}}/M_{\text{stars}}$ is of order unity (and consistent with zero), a factor of at least 400 lower than expected 2. NGC1052–DF2 demonstrates that dark matter is not always coupled with baryonic matter on galactic scales.”

A galaxy lacking dark matter

Calculation of the average speed of rotation in the galaxy NGC1052-DF2.

$H = \text{Hubble - Constant}$, $a_0 = \text{constant repulsive vacuum acceleration} = 6.911519372842 \cdot 10^{-10} \text{ m/s}^2$

$$a_0 = H \cdot c$$

$M(\text{NGC1052-DF2}) = \text{Mass of the galaxy NGC1052 - DF2} = 2 \cdot 10^8 \times 1.9885 \times 10^{30} \text{ Kg}$

$\bar{v}_r(\text{NGC1052-DF2}) = \text{average speed of rotation}$

$$2 \cdot \ln(M(\text{NGC1052-DF2})/m_{PK}) = \text{Entropy}$$

$m_{PK} = \text{Planck mass}$

$$\bar{v}_r(\text{NGC1052-DF2}) = \sqrt[4]{\frac{M(\text{NGC1052-DF2}) \cdot a_0 \cdot G_N}{2 \cdot \ln(M(\text{NGC1052-DF2})/m_{PK})}} = 17.1302252 \text{ Km/s}$$

The previous equation could be modified by a reduction correction factor ($\approx \varphi$); that possibly would depend, among other factors, on the shape of the galaxy.

$$\varphi = \frac{\sqrt{5} + 1}{2}$$

References

- [1] Angel Garcés Doz, Experiments with Powerful Neodymium Magnets: Magnetic Repulsion.phenomenological Equivalences: the Reverse Casimir Effect, or Repulsive (Spherical Shell) and the Spherical Shell of the Macroscopic Universe (Obsevable Sphere of the Universe), <http://vixra.org/abs/1603.0416>
- [2] Angel Garcés Doz, Axiomatization of Unification Theories: the Fundamental Role of the Partition Function of Non-Trivial Zeros (Imaginary Parts) of Riemann’s Zeta Function. Two Fundamental Equations that Unify Gravitation with Quantum Mechanics, <http://vixra.org/abs/1701.0042>
- [3] Angel Garcés Doz, The Experiments Of The Bottle And The Beam For The Lifetime Of The Neutron: A Theoretical Approximation Derived From The Casimir Effect, <http://vixra.org/abs/1803.0303>
- [4] Pieter van Dokkum, Shany Danieli, Yotam Cohen, Allison Merritt, Aaron J. Romanowsky, Roberto Abraham, Jean Brodie, Charlie Conroy, Deborah Lokhorst, Lamiya Mowla, Ewan O’Sullivan & Jielai Zhang, A galaxy lacking dark matter, Nature volume 555, pages 629–632 (29 March 2018), <https://www.nature.com/articles/nature25767>
- [5] Pieter van Dokkum, Shany Danieli, Yotam Cohen, Allison Merritt, Aaron J. Romanowsky, Roberto Abraham, Jean Brodie, Charlie Conroy, Deborah Lokhorst, Lamiya Mowla, Ewan O’Sullivan, Jielai Zhang, A galaxy lacking dark matter, 27 Mar 2018, <https://arxiv.org/abs/1803.10237>
- [6] Galaxy without any dark matter baffles astronomers, <https://www.theguardian.com/science/2018/mar/28/galaxy-without-any-dark-matter-baffles-astronomers>