

The X-cohomology

A.Balan

October 5, 2018

Abstract

With help of a closed form over the manifold, a X-cohomology of the manifold is defined

1 Definition

We consider a closed 1-form X over the manifold M ; $dX = 0$. Then we can define:

$$d_X(\alpha) = d(\alpha) + X \wedge \alpha$$

It is easy to verify that:

Theorem 1

$$d_X \circ d_X = 0$$

Demonstration 1 *Indeed:*

$$d(d\alpha + X \wedge \alpha) + X \wedge (d\alpha + X \wedge \alpha) = 0$$

So, we can have a X-cohomology of the manifold M :

Definition 1

$$XH^*(M, \mathbf{R}) = \text{Ker}(d_X) / \text{Im}(d_X)$$

2 The X-cohomology as a module

Theorem 2 *The X-cohomology $XH^*(M, \mathbf{R})$ is a module over the cohomology of de Rahm viewed as an algebra, $H^*(M, \mathbf{R})$.*

Demonstration 2 *If:*

$$d_X(\alpha) = 0$$

then:

$$d_X(\alpha \wedge \beta) = (-1)^{\text{deg}(\alpha)}(\alpha \wedge d\beta)$$

References

- [GHL] S.Gallot, D.Hulin, J.Lafontaine, "Riemannian Geometry", Springer-Verlag, Berlin, 2004.
- [J] N.Jacobson, "Basic Algebra" I and II, Dover, New-York, 2017.
- [S] R.Switzer, "Algebraic Topology-homotopy and homology", Springer-Verlag, Berlin, 2002.