

Weak Force and Weak Charge in Granular Spacetime

I Abstract

The standard model predicts a weak charge of SM prediction, $Q_p w(SM) = 0.0710 \pm 0.0007$. This value, SM prediction, $Q_p w(SM) = 0.0710 \pm 0.0007$, is the ratio of the weak charge to elementary charge. This paper shows that granular spacetime equations show a ratio of weak charge to elementary charge of $.071776 \pm .000001$. We also show that the ratio of weak charge and elementary charge are simply the square root of the mass of the neutron to the mass of two times the z boson or a combination.

II. Review of Equations of Force

The following equations have been developed to help unite the equations of force. So far the forces of gravity[1], electromagnetic[1], and the weak interaction[2].

Charge appears to come from electron sized charge.

$$q^2 = \frac{T\pi^3 hc\epsilon Me}{2Mn} \quad [2]$$

Where

$$T^2 = \frac{1}{\sqrt{1 - \left(2^{0.5} \frac{\pi Me}{3 * 3Mn}\right)^2}} \left[\left(\frac{Mp - Me}{Mn}\right)^2 + \left(\frac{Mn}{Mn}\right)^2 + \left(\frac{Mn}{Mn}\right)^2 \right] \quad [2.1]$$

Gravity appears to come from Proton sized mass

$$Mn^2 = \frac{2\pi^3 hcMp}{NGMn} \quad [3]$$

The Weak force appears to come from the Z bosons

$$q^2 = \frac{T\pi^3 hc\epsilon Me}{4Mz} \quad [4]$$

Note that q for equation 4 is a weak force charge and Mz is the z boson mass.

Weak Force and Weak Charge in Granular Spacetime

III Discrete calculation of weak charge

In this section we develop a potential discrete method for calculating weak charge from Planck's constant and the masses of the proton, neutron, and electron. The equation developed is the following.

$$q^2 = \frac{T\pi^3hc\epsilon Me}{4Mz} \quad [4]$$

where

Sum Angular Momentum Scalar

$$T^2 = \frac{1}{\sqrt{1 - (2^{0.5} \frac{\pi Me}{3 * 3Mn})^2}} \left[\left(\frac{Mp - Me}{Mn} \right)^2 + \left(\frac{Mn}{Mn} \right)^2 + \left(\frac{Mn}{Mn} \right)^2 \right] \quad [1] \quad [2.1]$$

Compton Wavelength of Z boson pairs $r = \frac{h}{cMz}$ [2.2]

Compton Radius of Neutron $r = \frac{h}{cMn}$ [2.2.1]

Compton Frequency of Neutron $f = \frac{Mnc^2}{h}$ [2.3]

where q is w boson charge

where pi or π is pi or π

where h is plancks constant

where c is speed of light

where ε is dielectric permittivity

To start we use the traditional equation of force between two charges.

$$Fe = \frac{q^2}{4\pi\epsilon r^2} \cdot \quad [2.4]$$

where Me is the mass of the electron

where Mn is the mass of the neutron

where Mp is the mass of the proton

T is defined as above

Mz is the mass of the z boson.

Weak Force and Weak Charge in Granular Spacetime

In this case we are calculating the weak charge and weak force max and estimating the mass of the Z boson. The value q will be the weak charge.

$$F_w = \frac{q^2}{4\pi\epsilon r^2} \quad [2.5]$$

Let us propose that charge can be calculated by summing the combined vector of the proton and electron force signified by the value "T".

$$T^2 = \frac{1}{\sqrt{1 - \left(2^{0.5} \frac{\pi M_e}{3 * 3 M_n}\right)^2}} \left[\left(\frac{M_p - M_e}{M_n}\right)^2 + \left(\frac{M_n}{M_n}\right)^2 + \left(\frac{M_n}{M_n}\right)^2 \right] \quad [1] \quad [2.1]$$

$$TF = \frac{q^2}{4\pi\epsilon r^2} \quad [2.6]$$

Since the electron, proton, and neutron all contain elementary charge within the quarks or electron, and these charges are all 1/3 or 2/3 or 3/3 of elementary charge within the quarks or electron, it is not unlikely that some relationship of this sort is possible. It should be noted that, in this situation, force can be modeled as a rate of angular momentum since there are two frequencies in the equation for force. One could be the rate of spinning, the other the rate of emission of angular momentum carriers.

It is known that $F=ma$, substituting yields;

$$Tma = \frac{q^2}{4\pi\epsilon r^2} \quad [2.7]$$

What is the acceleration of, A square, a circle, a sphere, a spherical shell? A spherical shell works for both force of charge and force of gravity. When attempts to pack spheres concentrically around other spheres a certain amount of defect space is made in relation to perfect packing. It can be shown that this amount of defect space is equal to the outer layer of spheres. Therefore, this is justification for using a hollow sphere when the actual geometry is not an actual hollow sphere. Therefore, the equation for acceleration of a spherical shell is as follows in equation [2.8]

The distribution of these discontinuities can be summed to be a spherical shell. This is shown in the paper "The Holographic Principle and How can the Particles and Universe be Modeled as a Hollow Sphere"(4)

Weak Force and Weak Charge in Granular Spacetime

$$a = \frac{2}{3} R(2\pi)^2 f^2 \quad [2.8]$$

Substituting for “a” yields;

$$Tm \frac{2}{3} R(2\pi)^2 f^2 = \frac{q^2}{4\pi\epsilon r^2} \quad [2.9]$$

Propose that the mass on the left hand side of the equation “m” is the mass of the electron “Me”

$$TMe \frac{2}{3} R(2\pi)^2 f^2 = \frac{q^2}{4\pi\epsilon r^2} \quad [2.10]$$

Propose that all masses and charges are divided by 3. Therefore, the equation becomes

$$TMe2R(2\pi)^2 f^2 = \frac{q^2}{4\pi\epsilon r^2} \quad [2.11]$$

Propose that radii are different, depending which force they are experiencing. The rationale for this is explained later in the discussion. It has to do with how the discontinuities are more concentrated at the center and the concentration of defects decreases inversely proportional to the radius. A radius of 10 would have approximately 20 percent defects, but a radius of 20 has only about 10 percent defects. To compensate for a large sphere the radii “r” are each divided by 4. Thus the equation becomes;

$$\frac{T}{2} MeR\pi^2 f^2 = \frac{q^2}{4\pi\epsilon r^2} \quad [2.12]$$

Substituting in Equation 2.2 and 2.3

Compton Wavelength of z boson pairs $R = \frac{h}{c2Mz} \quad [2.2]$

Weak Force and Weak Charge in Granular Spacetime

$$\text{Compton Radius of Neutron} \qquad r = \frac{h}{cMn} \qquad [2.2.1]$$

$$\text{Compton Frequency of Neutron} \qquad f = \frac{Mnc^2}{h} \qquad [2.3]$$

where q is weak charge
 where pi or π is pi or π
 where h is Planck's constant
 where c is speed of light
 where Me is the mass of the electron
 where Mn is the mass of the neutron
 where Mp is the mass of the proton
 where Mz is the mass of the Z bosons
 and T is defined as above

Which simplifies to

$$q^2 = \frac{T\pi^3hc\epsilon Me}{4Mz} \qquad [2.13]$$

Substituting the values from the appendix

$$q = .071776 \pm .000001 * 1.602176622 * 10^{-19} \text{ for the Z boson}$$

Both the ratio of weak charge to elementary charge for the Z boson of 0.071776 ± 0.000001 compare well to the 0.0719 ± 0.0045 ratio of the weak charge of the proton discussed in nature[4] and the 0.0710 ± 0.0007 value calculated from the Standard Model.[5]

Please note that the ratio of weak charge and elementary charge are simply the square root of the mass of the neutron to the mass of two times the

Weak Force and Weak Charge in Granular Spacetime

$$q_{\text{ratioweakcharge}} \text{ to elementary charge} = \sqrt{\frac{Mn}{2Mz}} = 0.71776 \pm 0.000001 \text{ for the z boson}$$

IV Discussion

The value of "T", that is proposed in section II, shown below;

Sum Angular Momentum

$$T^2 = \frac{1}{\sqrt{1 - \left(2^{0.5} \frac{\pi Me}{3 * 3Mn}\right)^2}} \left[\left(\frac{Mp - Me}{Mn}\right)^2 + \left(\frac{Mn}{Mn}\right)^2 + \left(\frac{Mn}{Mn}\right)^2 \right] \quad [2.1]$$

may have other factors affecting the forces in the x, y, and z dimension. The neutrinos mass appears to be so small, that if there is a neutrino were incorporated into the mass of the proton or neutron it might affect the mass in the 10th, or 11th digit. Note that equation 2.1 was used in Proton Electron Universe and Their Directions of Force[1]. It is presumed here that this same value may be used for the weak charge. The actual weak charge is not known to much accuracy so it did not seem of value to speculate if it is different than the value for elementary charge.

V Conclusions

We see that the same form of equations can be used for calculating the elementary charge, elementary graviton, and the weak charge. These equations are shown below.

$$q^2 = \frac{T\pi^3 hc \epsilon Me}{2Mn} \quad [2]$$

Where

Gravity appears to come from Proton sized mass

Weak Force and Weak Charge in Granular Spacetime

$$Mn^2 = \frac{2\pi^3 hcMp}{NGMn} \quad [3]$$

The Weak force appears to come from the Z bosons

$$q^2 = \frac{T\pi^3 hc\epsilon Me}{4Mz} \quad [4]$$

Note that q for equation 4 is a weak force charge and Mz is the z boson mass.

$$q_{\text{ratioweakcharge to elementary charge}} = \sqrt{\frac{Mn}{2Mz}} = 071776 \text{ for the z boson}$$

Weak Force and Weak Charge in Granular Spacetime

Appendix A

Fundamental Physical Constants (18)

1. $c=2.99792458 \times 10^8$ m/s
2. $h=6.626\ 070\ 040(81) \times 10^{-34}$ J s $6.626\ 06957(33) \times 10^{-34}$ J s
3. Mass of Neutron = $M_n=1.674\ 927\ 471(21) \times 10^{-27}$ kg $1.674\ 927\ 351(74) \times 10^{-27}$ kg
4. Mass of Proton = $M_p=1.672\ 621\ 898(21) \times 10^{-27}$ kg $1.672\ 621\ 777(74) \times 10^{-27}$ kg
5. Mass of Electron = $M_e=9.109\ 383\ 56(11) \times 10^{-31}$ kg $9.109\ 382\ 91(40) \times 10^{-31}$ kg.
6. q = unit charge = $1.602\ 176\ 6208(98) \times 10^{-19}$ C $1.602\ 176\ 565(35) \times 10^{-19}$ C
7. ϵ = Dielectric Permittivity = $8.854187817 \times 10^{-12}$
8. $G=6.674\ 08(31) \times 10^{-11}$ m³ kg⁻¹ s⁻² $6.67384(80) \times 10^{-11}$ m³ kg⁻¹ s⁻²
9. Z boson mass= $1.62557e-25$ kg or 91.1876(21) GeV

References

1. <http://vixra.org/pdf/1807.0519v1.pdf>
2. https://www.science20.com/tommaso_dorigo/things_that_can_decay_to_boson_pairs-228083
3. <http://vixra.org/pdf/1712.0672v3.pdf>
4. <https://www.nature.com/articles/d41586-018-05037-9>
5. <http://iopscience.iop.org/article/10.1088/1742-6596/678/1/012026/pdf>