

# Pythagoras theorem is an alternate form of Ptolemy's theorem

Radhakrishnamurty Padyala  
A-102, Cedar, Brigade Orchards,  
Devanahalli, Bengaluru – 562110, India.  
Email: padyala1941@yahoo.com

## Abstract

Generally, the proofs given to demonstrate Ptolemy's theorem prove Pythagoras theorem as a special case of Ptolemy's theorem when certain special conditions are imposed. We prove in this article, that Pythagoras theorem follows from Ptolemy's theorem in all cases. Therefore, we may say that Pythagoras rediscovered Ptolemy's theorem.

## Key words

Ptolemy's theorem, geometry, Pythagoras theorem, cyclic quadrilateral, mean proportional

## Introduction

Ptolemy's theorem is a well-known theorem in geometry. Many different methods of proof are available in literature<sup>1-4</sup>. In a recent article<sup>1</sup> we gave a geometrical proof of Ptolemy's theorem. Therein we showed that the products of pairs of opposite sides of a cyclic quadrilateral can be geometrically represented by rectangles. Similarly the product of the diagonals of the same quadrilateral can be represented by another rectangle. Then we showed that the sum of the areas of the rectangles obtained from the products of sides is equal to the area of the rectangle obtained from the product of the diagonals.

In this article we show the products of pairs of opposite sides as well as the product of the diagonals of a cyclic quadrilateral can be geometrically represented by circles. Then we show that the sum of the areas of the circles obtained from the products of sides is equal to the area of the circle obtained from the product of the diagonals. To obtain the radii of these circles we take the mean proportional of the pairs of opposite sides of the cyclic quadrilateral and the mean proportional of the pair of its diagonals.

## Statement of Ptolemy's theorem<sup>2</sup>

Ptolemy's theorem states that, the sum of the products of pairs of opposite sides of a cyclic quadrilateral is equal to the product of the diagonals.

## Proof

Draw any circle. Choose any four points A, B, C and D on it. Join A, B; B, C; C, D; A, D; A, C; and B, D. Now ABCD is a cyclic quadrilateral. AB, CD is a pair of opposite sides, AD, BC is the other pair of opposite sides and AC, BD are the diagonals (see Fig. 1).




Fig. 1. ABCD is a cyclic quadrilateral. Ptolemy's theorem says,  $AB \cdot DC + AD \cdot BC = AC \cdot BD$ .

We can write Ptolemy's theorem in the form of equation (1) below.

$$AB \times CD + AD \times BC = AC \times BD \quad (1)$$

Instead of taking the products of the line segments, we can take their mean proportionals. Let  $r_1$  be the mean proportional of the pair of opposite sides AB, CD;  $r_2$  be the mean proportional of the pair of opposite sides AD, BC and  $r_3$  be the mean proportional of the diagonals AC, BD. Then we get,

$$AB \times CD = r_1^2 \quad (2)$$

$$AD \times BC = r_2^2 \quad (3)$$

$$AC \times BD = r_3^2 \quad (4)$$

We can now state Ptolemy's theorem in an alternate form as follows:

#### **Alternate statement of Ptolemy's theorem**

The sum of the squares of the mean proportionals of the pairs opposite sides of a cyclic quadrilateral is equal to the square of the mean proportional of the diagonals. We can write this alternate form of statement of Ptolemy's theorem in the form of Eq (5) below.

$$(r_1^2 + r_2^2) = r_3^2 \quad (5)$$

Equation (5) is a statement of Pythagorus' theorem.






Fig. 3. Circles are drawn with A as center and the mean proportionals as radii.

We draw the three concentric circles with A as center and radii  $r_1$  (green),  $r_2$  (red),  $r_3$  (blue). The areas of these circles are  $\pi r_1^2$ ,  $\pi r_2^2$ ,  $\pi r_3^2$ . From Eq. (5) we see that the sum of the areas of the two inner circles is equal to the area of the outer most circle. To demonstrate that it is so, we proceed as follows.

Draw a circle with AG as diameter (see Fig. 4). From the opposite ends A, G of AG draw circles with radii,  $r_1$ ,  $r_2$ . These circles intersect at four points marked  $P_1$ ,  $P_2$ ,  $P_1'$ ,  $P_2'$ . We join A,  $P_1$ ;  $P_1$ , G; A,  $P_2'$  and G  $P_2'$ . We get the cyclic quadrilateral A  $P_1 G P_2'$ . We apply Ptolemy's theorem to this quadrilateral and get Eq. (5).

Since the radii  $r_1$ ,  $r_2$  are mean proportionals of the pairs of opposite sides, and  $r_3$ , the mean proportional of the diagonals of the cyclic quadrilateral ABCD it follows that the sum of the squares of the mean proportionals of the pairs of opposite sides, is equal to the square of the mean proportional of the diagonals of a cyclic quadrilateral. Thus we find Pythagoras' theorem is an alternate form Ptolemy's theorem. Therefore, we may say Pythagoras rediscovered Ptolemy's theorem.




Fig. 4. Circles are drawn with AG as diameter. From the opposite ends A and G as centers, circles are drawn with radii  $r_1$  and  $r_2$ . The points of intersection are marked as  $P_1$ ,  $P_2$ ,  $P_1'$ ,  $P_2'$ . Cyclic quadrilateral  $AP_1GP_2'$  is drawn.

## Acknowledgement

My sincere thanks are due to Mr. Arun Mozhi Selvan Rajaram, EPI group, Chennai, India, without whose assistance and support, this article could not have been completed.

## References

1. Radhakrishnamurty Padyala, A geometrical proof of Ptolemy's theorem <http://viXra.org/abs/2005.0026>, RG DOI: 10.13140/RG.2.2.20787.17446.
2. Shailesh Shirali, At Right angles, November (2016)
3. Zvezdelina Stankova, Ptolemy's Theorem, <https://www.youtube.com/watch?v=bJOuzqu3MUQ>
4. Marano Perez de la Cruz, Ptolemy via Cross Ratio, cut-the-knot.org/proofs/ptolemyCrossRatio.shtml