

Impact-area of an Ejecta-Ray from the Port Hedland Crater, located near Southern-Cross (Western Australia)

- Raman Spectra of selected Rock Samples - by Harry K. Hahn, 30.12.2021 -

Summary :

The visited area is located near the small town Southern-Cross in Western Australia. The Gravity Anomaly Map indicates that **Ejecta-material from the assumed \varnothing 400 x 350 km Port Hedland Crater (Impact) in all probability impacted here and formed these linear structures.** The Port Hedland Crater is located north of the town Port Hedland on the sea-floor of the Indian Ocean. The Port Hedland Crater, which is unknown yet, in all probability is a large secondary crater caused by the Permian-Triassic Impact Event.

Another possible source of the linear ejecta rays could be the Victoria Lake Impact (see map on next page)

For a detailed description of the Permian-Triassic Impact (PTI) Hypothesis please read **Part 1 (P1)** of my hypothesis. And for more information to the \varnothing 400 x 350 km Port Hedland Crater (PHC) please read pages 14-16, 20-21 and 24-27 of **Part 3 (P3)** and page 33 of **Part 2 (P2)** of my hypothesis.

The geological map identifies the nearly linear structures as structures (low mountain ranges) which consist of different rock types than the surrounding plains of the Yilgarn Craton. These linear structures seem to penetrate the Yilgarn Craton down to a depth of around 6 km (see geologic cross-section A-B)

I have collected some rock-samples from these nearly linear structures in the Southern-Cross area and have analysed these samples, mostly quartz, with Micro-Raman-Spectroscopy, to find out if they were exposed to a shock pressure which may indicate an Impact Event. And indeed that seems to be the case !

The Raman-spectra of quartz from the Sample Sites 1, 9, 16 and 18 provide first evidence for an impact event as the probable cause of the linear ray-structure in the Southern-Cross area.

The clear shifts of the main Raman peaks, of the analysed quartz samples, to the lower frequencies 463, 261, 204 and 126 cm^{-1} (Site 1), to 463, 261 and 205 cm^{-1} (Site 9 & 16) and to 463, 261, 204 and 125 cm^{-1} (Site 18) provide first proof for an Impact Event as the cause of the linear ray-structure !

(\rightarrow see explanation in **Appendix 1** at page **25** : Overview : The Raman bands (peaks) of shocked Quartz)

Microscopic images of some analysed quartz grains seem to provide further proof for a shock event.

PDFs (planar deformation features) seem to be present in some samples (\rightarrow images on **pages 4 to 11**)

All spectra were made with a **BRUKER Senterra-II Raman Microscope** (wavenumber precision $<0.1\text{cm}^{-1}$)


\rightarrow Images of the analysed rock samples and photos of the sample sites are in the Appendix at **page 18**.

\rightarrow More images of all sample sites are available on www.permiantriassic.de or www.permiantriassic.at


\rightarrow **References** : see **page 26** / and pages 14-16, 20-21 and 24-27 of **Part 3 (P3)** of my hypothesis.

Note : A shock pressure of 20 GPa exceeds every pressure caused by normal terrestrial metamorphism. The indicated shock pressures of $\approx 20\text{-}22$ GPa therefore in general point to an impact shock event.


Gravity Anomaly Map of Southern-Cross area




\varnothing 400 x 350 km Port Hedland Crater




Port Hedland Crater (= Bengal Bay Crater) + surrounding area (Gravity Anomaly Map)



Gravity Anomaly Map of SW-Australia




Geological Map of South-West-Australia
→ with sample site area marked on the map




The Ejecta-Ray area near the town Southern-Cross is marked in green on the Geological Map above

The thin ejecta-ray-structures visible on the gravity anomaly map of Western Australia as linear red (positive) anomalies, were caused either by the $\varnothing 400 \times 350 \text{ km}$ Port Hedland Crater

$\varnothing 400 \times 350 \text{ km}$ Port Hedland Crater




(=Bengal Bay Crater) or by the Victoria Lake Impact Crater. According to my Permian-Trassic Impact (PTI) hypothesis the Port Hedland Crater (PHC) is a big secondary-crater caused by the **PT-impact Event**, which is also responsible for the formation of the Bengal Bay (= Bengal Bay Crater) The topographic map below shows the original situation at the time of the PT-impact Event. The gravity anomaly map indicates a number of linear ejecta-ray-structures (red) on the Yilgarn Craton which are (nearly) parallel




This map shows the probable arrangement of the "tectonic plates" before the PTI

Topographic map


Sample Site 1: Stone 1_spectra 1 indicates: Quartz





Sample:



Sample Site 1: Stone 2_spectra 1 indicates: Quartz



Sample :




Sample Site 1: Stone 2 : Quartz


Detail of Microscopic Image

Image size : ~ 250 x 200 µm

Sample Site 9: Stone 1_spectra 1 indicates: Quartz




Sample :




Microscopic Images : Sample from Site 1 and 9 → original state (no preparation for analysis)


Sample Site 1 : Stone 1 : Quartz (Image ~ 300 x 250 μm)




Sample Site 1 : Stone 2 : Quartz (Image ~ 500 x 400 μm)




Sample Site 9 : Stone 1 : Quartz (Image: ~ 500 x 400 μm)





Detail : Image size : ~ 250 x 200 μm



Sample Site **16** : Stone 1_spectra 1 indicates : **Quartz**



Sample :




Sample Site **16**: Stone 1 : **Quartz**


Detail of Microscopic Image

Image size : ~ 250 x 200 μm

Sample Site 18: Stone 1_spectra 1 indicates: Quartz




Sample :




Sample Site 18: Stone 1 : Quartz

Detail of Microscopic Image


Image size : ~ 250 x 200 μm



Sample Site 2: Stone 1_spectra 1 indicates: Quartz




Sample:




Microscopic Images : Sample from Site 2, 16 and 18 → original state (no preparation for analysis)


Sample Site **16** : Stone 1 : Quartz (Image ~ 500 x 400 μm)




Sample Site **18** : Stone 2 : Quartz (Image ~ 500 x 400 μm)




Sample Site **2** : Stone 1 : Quartz (Image: ~ 500 x 400 μm)



Detail : Image size : ~ 250 x 200 μm



Sample Site **16** : Stone 2_spectra 1 indicates : **Quartz**



Sample :







Image size : ~ 400 x 300 μm

Detail : Image size : ~ 200 x 150 μm



Sample Site **11** : Stone 1_spectra 1 indicates: **Amicite_Quartz_Dachiardite-Na** (→ see RRUFF_CS results)




Sample :



CrystalSleuth: EXTRACT_11_SC-WA_stein_1_black-white_0_000000_0_NK_Y_G1

File Edit Mode Help




% Match	Spectrum Name	RRUFF ID
96	-) Amicite (532nm)	R080066
95	-) Monacite-(Ce) (532nm)	R060925
94	-) Dachiardite-Na (532nm)	R061116
93	Bassetite (532nm)	R080027
92	Arhbarite (532nm)	R060730
91	Polyhalite (532nm)	R070209
91	Keystoneite (532nm)	R070661
91	Wilamaninite (532nm)	R060514
90	Sugilite (532nm)	R070684
89	-) Quartz (532nm)	X080015
89	Quartz (532nm)	X080016
89	Zemmarite (532nm)	R050273
89	Fraserite (532nm)	R070416

X080015
Quartz
SiO₂
Synthetic


R080066
Amicite
K₂Na₂(Si₄Al₄O₁₆·18H₂O
Kirovski Drive, Khibiny, Kola Peninsula, Russia

R061116
Dachiardite-Na
Na₄(Si₂₀Al₄O₄₈·18H₂O
Alpe di Siusi, Bolzano, Trentino-Alto Adige, Italy



Sample Site **11** : Stone 1_spectra 2 indicates: **Quartz**

(→ see RRUFF_CS results)




Note the black- and white laminated structure of the Quartz

Sample :




Detail of microscopic image



CrystalSleuth: EXTRACT_11_SC-WA_stein_1_black-white_0_000007_0_NK_G1


File Edit Mode Help



File Manager | SpecEdit | Raman Library | X-Ray

% Match	Spectrum Name	RRUFF ID
90	< > Quartz (532nm)	X080016
90	Quartz (532nm)	X080015
90	Quartz (532nm)	R060604
89	Quartz (532nm)	R040031
88	Quartz (532nm)	R050125
84	Amicite (532nm)	R080066
83	Dachardite-Na (532nm)	R061116
82	Edgarbeleyite (532nm)	R060500
79	Sugilite (532nm)	R070684
79	Monazite-(Ce) (532nm)	R060925
76	Sodalite (532nm)	R060405
75	Sodalite (532nm)	R060435
75	Sodalite (532nm)	R040141

X080016
Quartz
SiO₂
Synthetic






Sample Site **11** : Stone 2_spectra 1 indicates: **Quartz_**

(→ see RRUFF_CS results)


Note the red- and black laminated structure of the Quartz

The spectrum was measured in the red-colored quartz

Sample :





Sample Site **12**: Stone 1_spectra 1 indicates: **Erdite_Cechite_Florencite-(Ce)_Geothite** (→ see RRUFF_CS)




Note the laminated structure of the mineral

→ Iron-bearing mineral !

Sample :



Sample Site 12 : Stone 3_spectra 1 indicates : **Heulandite-Sr_Mordenite_Dachiardite-Na_Amicite** (→ RRUFF)




Sample :



CrystalSleuth: EXTRACT_12_SC-WA_stein 3_-black-glas.0_000006.0_NK_Y_G1

File Edit Mode Help




% Match:	Spectrum Name:	RRUFF ID:
86	< Heulandite-Sr (532nm)	R070272
84	< Mordenite (532nm)	R061118
82	< Dachiardite-Na (532nm)	R061116
78	Monazite-(Ce) (532nm)	R060925
77	< Amicite (532nm)	R080066
77	Epistilbite (532nm)	R061105
76	Ericssonite (532nm)	R070406
75	Nepheline (532nm)	R060581
74	Gismondine (532nm)	R060809
74	Quartz (532nm)	X080016
74	Stilwellite-(Ce) (532nm)	R060899
73	Mordenite (532nm)	R070524
73	Quartz (532nm)	X080015

R070272
Heulandite-Sr
Na5Sr_4(Sr_27_Al_9)_O_72_#183;24H_2O
Campel mine, Campel, Castiglione Chiavarese, Genova Province


R061118
Mordenite
(Na_2_Ca_K_2)_4(Al_8_Si_40)_96_#183;28H_2O
Chapman's Brook, Nova Scotia, Canada

R061116
Dachiardite-Na
Na_4(Sr_20_Al_4)_O_48_#183;13H_2O
Alpe di Susi, Bolzano, Trentino-Alto Adige, Italy


R080066
Amicite
K_2Na_2(Si_4Al_4)_O_16_#183;5H_2O
Krovaki Drive, Khibiny, Kola Peninsula, Russia





Detail of microscopic image :



Sample Site 17: Stone 1_spectra 1 indicates: **Oligoclase, Labradorite** (→ see RRUFF_CS results)



Sample :



Appendix 1 : Photos of the rock samples from the analysed sample sites :


→ See next page !

Please note : Photos of all Sample Sites & Rock Samples are available on my website :


→ [Samples from Southern Cross Area](#) or here : [Southern Cross Area](#)

Geological Map of SW-Australia




Location where samples were collected :




Southern Cross area



Geological Map : Jackson_sh5012




Sample sites No. : 1 to 24 marked in yellow on the maps



Cross-Section A - B

Geological Map : Jackson_sh5012





The sections of an ejecta-ray of the Port Hedland Crater (or VLC) → the nearly linear green-colored structures) seem have penetrated the Yilgarn Craton down to a **depth of around 6 km**.

see Detail 2


→ Geological Maps can be downloaded here :

<http://www.geoscience.gov.au/>

Then go to "Geology" – 1:250K Geological Maps and search for the required map











Sample site 12





Sample site 16





Appendix 2 : A short overview : The Raman bands (peaks) of Quartz shocked with 22-26 GPa


In order to verify a sample site as an impact site or impact structure, **shock-metamorphic effects** must be discovered in the rocks of the sample site. This can be done by different methods.

For example with the help of PDFs (planar deformation features) which are visible in the quartz with the help of a microscope. However this requires careful preparation of the samples and expertise.

Another, easier method, is the use of a RAMAN microscope. Micro-RAMAN Spectroscopy on quartz grains in the samples can provide the first evidence for a shock event, that was caused by an impact.


Mc Millan et al. (1992) and others have shown that the main RAMAN-peaks of Quartz shift towards lower frequencies if the Quartz was exposed the a shock-pressure > 15 GPa. → see diagram below

The shift of the main quartz RAMAN-peaks can be used to identify quartz that was shocked by an impact



Quartz shocked with 22 GPa and 26 GPa shows shifts of the main RAMAN-peaks of 1 - 4 cm⁻¹ to lower frequencies

Appendix 3 : Raman spectra of (W) weakly-shocked & (M) moderately-shocked Alkali-Feldspar



Weakly shocked alkali feldspar mainly developed irregular fractures and undulatory extinction. Note that the Raman-lines 210 and 765 are missing in the w-shocked feldspar, and an additional line at ≈ 150 appears.

The shock pressure for the w-shocked feldspar was estimated to be between 5 and 14 GPa

References :

Photos of Sample Sites & Rock Samples are available on : [Samples from Southern Cross Area](#) (or : [Southern Cross Area](#))

Find more information to the linear Ejecta-Ray structures in W-Australia in Parts 2 & 3 of my hypothesis - by Harry K. Hahn
Please read pages 14-16, 20-21 & 24-27 of [Part 3 \(P3\)](#) & page 33 of [Part 2 \(P2\)](#) of my hypothesis (→ [weblinks below!](#))

Please also read my Raman-analyses to rock samples from the [Kalgoorlie area](#) ; [Geraldton-area](#) & [Margaret-River area](#) !!
→ You can find these analyses either on [www.vixra.org](#) or on [www.archive.org](#) → under my author name : Harry K. Hahn

The Permian-Triassic (PT) Impact hypothesis - by Harry K. Hahn - 8. July 2017 :

Part 1 : [The 1270 X 950 km Permian-Triassic Impact Crater caused Earth's Plate Tectonics of the Last 250 Ma](#)

Part 2 : [The Permian-Triassic Impact Event caused Secondary-Craters and Impact Structures in Europe, Africa & Australia](#)

Part 3 : [The PT-Impact Event caused Secondary-Craters and Impact Structures in India, South-America & Australia](#)

Part 4 : [The PT-Impact Event and its Importance for the World Economy and for the Exploration- and Mining-Industry](#)

Part 5 : [Global Impact Events are the cause for Plate Tectonics and the formation of Continents and Oceans \(Part 5\)](#)

Part 6 : [Mineralogical- and Geological Evidence for the Permian-Triassic Impact Event](#)

Alternative weblinks for my Study **Parts 1 - 6 with slightly higher resolution** : [Part 1](#), [Part 2](#), [Part 3](#), [Part 4](#), [Part 5](#), [Part 6](#)

Parts 1 – 6 of my PTI-hypothesis are also available on my website : [www.permiantriassic.de](#) or [www.permiantriassic.at](#)

Shock-metamorphic effects in rocks and minerals - <https://www.lpi.usra.edu/publications/books/CB-954/chapter4.pdf>

Shock metamorphism of planetary silicate rocks and sediments: Proposal for an updated classification system

Stöffler - 2018 - Meteoritics & Planetary Science – Wiley: <https://onlinelibrary.wiley.com/doi/epdf/10.1111/maps.12912>

A Raman spectroscopic study of shocked single crystalline quartz - by P. McMillan, G. Wolf, Phillipe Lambert, 1992

<https://asu.pure.elsevier.com/en/publications/a-raman-spectroscopic-study-of-shocked-single-crystalline-quartz>

alternative : <https://www.semanticscholar.org/paper/A-Raman-spectroscopic-study-of-shocked-single-McMillan-Wolf/cfaaf6eb3e46fbd2912fb91c7acf40e88e721132>

Raman spectroscopy of natural silica in Chicxulub impactite, Mexico - by M. Ostroumov, E. Faulques, E. Lounejeva

https://www.academia.edu/8003100/Raman_spectroscopy_of_natural_silica_in_Chicxulub_impactite_Mexico

alternative : <https://www.sciencedirect.com/science/article/pii/S1631071302017005>

Shock-induced irreversible transition from α -quartz to CaCl₂-like silica - Journal of Applied Physics: Vol 96, No 8

<https://aip.scitation.org/doi/10.1063/1.1783609>

Shock experiments on quartz targets pre-cooled to 77 K - J. Fritz, K. Wünnemann, W. U. Reimold, C. Meyer

https://www.researchgate.net/publication/234026075_Shock_experiments_on_quartz_targets_pre-cooled_to_77_K

A Raman spectroscopic study of a fulgurite – by E. A. Carter, M.D. Hargreaves, ...

https://www.researchgate.net/publication/44655699_Raman_Spectroscopic_Study_of_a_Fulgurite

alternative : <https://royalsocietypublishing.org/doi/abs/10.1098/rsta.2010.0022>

Shock-Related Deformation of Feldspars from the Tenoumer Impact Crater, Mauritania - by Steven J. Jaret

<https://trace.tennessee.edu/cgi/viewcontent.cgi?article=1002&context=pursuit>

A Study of Shock-Metamorphic Features of Feldspars from the Xiuyan Impact Crater - by Feng Yin, Dequi Dai

https://www.researchgate.net/publication/339672303_A_Study_of_Shock-Metamorphic_Features_of_Feldspars_from_the_Xiuyan_Impact_Crater

https://www.researchgate.net/publication/339672303_A_Study_of_Shock-Metamorphic_Features_of_Feldspars_from_the_Xiuyan_Impact_Crater

Shock effects in plagioclase feldspar from the Mistastin Lake impact structure, Canada – A. E. Pickersgill – 2015

<https://onlinelibrary.wiley.com/doi/pdf/10.1111/maps.12495>

Shock Effects in feldspar: an overview - by A. E. Pickersgill

<https://www.hou.usra.edu/meetings/lmi2019/pdf/5086.pdf>

ExoMars Raman Laser Spectrometer RLS, a tool for the potential recognition of wet target craters on Mars

https://www.researchgate.net/publication/348675414_ExoMars_Raman_Laser_Spectrometer_RLS_a_tool_for_the_potential_recognition_of_wet_target_craters_on_Mars