

The black hole fallacy

A.J.Owen

October 31st 2022

Author's contact email address:

ajowen@physics.org

Abstract

It is shown here that the currently accepted solution in general relativity predicting black holes and an event horizon is unphysical. The spacetime outside a point mass is entirely regular, the effective gravitational mass decreases to zero as a test object comes into close proximity with it, and free-fall velocities do not exceed the speed of light. The ringlike radio images of galactic centres can be attributed to gravitational lensing of regular point sources.

1 Gravitational field due to a point mass

A solution using Albert Einstein's theory of general relativity (*GR*) [1] for the gravitational field due to a point mass was first obtained by Karl Schwarzschild [2] in 1916. A year later Droste [3] and Weyl [4] independently obtained a further variant of the solution. Subsequently, Hilbert [5] extended Droste and Weyl's solution in such a way that it showed a discontinuity in spacetime which is now interpreted as an event horizon obscuring the central mass, which is called a black hole.

The procedure for obtaining the solution is summarised, as follows. A metric line element in a curved spacetime (t, r, θ, ϕ) with spherical spatial symmetry about the origin may be written:

$$d\tilde{s}^2 = c^2 dt'^2 = A(r) c^2 dt^2 - B(r) dr^2 - C(r) d\Omega^2 \quad (1)$$

where $d\tilde{s}$ is a spacetime increment, c is the speed of light, dt' an increment of proper time, dt an increment of coordinate time, dr an increment of radial coordinate distance r , and $d\Omega^2 = d\theta^2 + \sin^2 \theta d\phi^2$; A, B and C are radially dependent functions describing the curvature of the time, radial and angular metric coefficients, respectively.

Lagrangian formalism is then used to obtain the geodesic equations in t, r, θ, ϕ , and from these the Christoffel curvature coefficients and Ricci tensor components are found. The Ricci tensor components are then set equal to zero, in order to satisfy Einstein's *GR* field equations for the vacuum outside the point mass. This leads to the following pair of simultaneous equations relating A, B and C :

$$\frac{A'}{A} + \frac{B'}{B} = \frac{2C''}{C'} - \frac{C'}{C} ; \quad \frac{A'}{A} - \frac{B'}{B} = -\frac{2C''}{C'} + \frac{4B}{C'} \quad (2)$$

where $A' = dA/dr$, $B' = dB/dr$, $C' = dC/dr$ and $C'' = d^2C/dr^2$. There are two independent equations for the three functions, A, B , and C , which means they cannot be solved explicitly, without some additional condition or assumption.

To proceed, it is customary to define a new radial coordinate which I shall denote as \tilde{r} in terms of C , where

$$\tilde{r} = \sqrt{C(r)} \quad (3)$$

and then the metric becomes:

$$d\tilde{s}^2 = c^2 A(\tilde{r}) dt^2 - B(\tilde{r}) d\tilde{r}^2 - \tilde{r}^2 d\Omega^2 \quad (4)$$

With this coordinate transformation, Equations 2 can be solved in terms of $(t, \tilde{r}, \theta, \phi)$, often called Schwarzschild coordinates, resulting in the solution:

$$d\tilde{s}^2 = c^2 \left(1 - \frac{\alpha}{\tilde{r}}\right) dt^2 - \left(1 - \frac{\alpha}{\tilde{r}}\right)^{-1} d\tilde{r}^2 - \tilde{r}^2 d\Omega^2 \quad (5)$$

i.e.

$$A(\tilde{r}) = \frac{1}{B(\tilde{r})} = 1 - \frac{\alpha}{\tilde{r}} \quad ; \quad C = \tilde{r}^2 \quad (6)$$

α is a positive, real constant of integration that is found by comparing the solution with known physics (to be discussed later). In addition, since $A(\tilde{r})B(\tilde{r}) = 1$, we have

$$A(r)B(r) = \left(\frac{d\tilde{r}}{dr}\right)^2 \quad (7)$$

The exact relationship between \tilde{r} and r is not specified or deducible from the spherical geometry alone, so the above procedure merely expresses A and B in terms of C (or the coordinate \tilde{r}). Many teachers of relativity ignore any difference between \tilde{r} and r , and regard \tilde{r} as the

"true" radial coordinate distance r from the mass M .¹ This opens the door to the idea of a black hole, since $(1 - \alpha/r)$ changes sign if r could go from above to below α .

In his original solution, Schwarzschild [2] had already realised that $(1 - \alpha/\tilde{r})$ becomes negative if $\tilde{r} < \alpha$. However, he regarded this as unphysical, and to prevent it happening defined an auxiliary radial coordinate \tilde{r} by the function:

$$\tilde{r} = (r^3 + \alpha^3)^{1/3}$$

so that as $r \rightarrow 0$, $\tilde{r} \rightarrow \alpha$, and the discontinuity is forced to occur at the origin. Thus, Schwarzschild himself rejected the idea of A or B changing sign or becoming infinite in free space, and he therefore did not predict black holes, even though current popular science suggests he did.

Soon afterwards, both Droste and Weyl published a different variant, essentially in which $\tilde{r} = r$, but they avoided the discontinuity by limiting the range of \tilde{r} to $\alpha < \tilde{r} < \infty$. Subsequently, Hilbert extended Droste and Weyl's solution to the range $\tilde{r} < \alpha$, i.e. into the region where the functions A and B become negative, and then spacetime is discontinuous. In modern language this is called an event horizon, behind which a black hole is obscured. Hilbert has a good argument for extending the solution to $\tilde{r} \rightarrow 0$, even if \tilde{r} is not equal to r , on the grounds that GR is intended to be a generally covariant theory, implying that a change of coordinates should not alter the physics of the situation.

Although this extension by Hilbert is currently the generally accepted solution, several mathematicians have questioned it. According to Abrams [6] and Crothers [7], the complete set of possible solutions for \tilde{r} as a function of r is given by

$$\tilde{r} = (|r - r_0|^n + \alpha^n)^{1/n} \quad (8)$$

where n is a positive integer, r_0 an arbitrary constant, and α a real positive constant. Schwarzschild's original solution is represented here by $n = 3$ and $r_0 = 0$. Droste and Weyl's solution has $n = 1, r_0 = \alpha, r > r_0$, and in a paper by Brillouin [8], there is a solution with $n = 1, r_0 = 0$. Crothers [7] points out that Hilbert's extension of Droste and Weyl's solution does not belong to this set, since the modulus of $(r - r_0) \geq 0$, and therefore $\tilde{r} \geq \alpha$. The function $(1 - \alpha/\tilde{r})$ then cannot become negative, and so black holes are not actually predicted

¹By "true" is meant here the distance according to an observer in a flat frame of reference a very long way from the mass causing gravitation, or in a flat frame that would have existed if the effect of the gravitational mass M were somehow removed.

mathematically. However, this objection has not been accepted by the scientific community to date, and so Hilbert's extension to $\tilde{r} < \alpha$ is still regarded as correct.

2 Relating GR to the physical world

The quantity α relates the geometrical scale of GR to gravitational forces in the real world. To obtain this connection, one can consider the equation of motion of a test object free-falling from rest at infinity along a radial path towards the central mass at the coordinate origin. The required equation is obtained from the geodesic in r and given by

$$\ddot{r} + \frac{A'}{2B}c^2\dot{t}^2 + \frac{B'}{2B}\dot{r}^2 = 0 \quad (9)$$

where \ddot{r} ($= d^2r/dt'^2$) is the proper acceleration and \dot{r} ($= dr/dt'$) is the proper velocity. Using the metric to eliminate \dot{t} ($= dt/dt'$), this becomes

$$\ddot{r} + \frac{A'}{2AB}c^2 + \frac{1}{2}\left(\frac{A'}{A} + \frac{B'}{B}\right)\dot{r}^2 = 0 \quad (10)$$

This expression in terms of the curved geometry of spacetime is then compared with Newton's law of gravitation in the regime where Newton's law is valid, which is considered to be in the weak-field region at large r . Newton's law may be written:

$$a + \frac{GM}{r^2} = 0 \quad (11)$$

where a is the classical acceleration, G Newton's gravitational constant and M the mass causing gravity. Superficially, there is no obvious agreement, but if one temporarily ignores any difference between \tilde{r} and r , it follows from Equation 7 that $B = 1/A$, and then by eliminating B Equation 10 would become

$$\ddot{r} + \frac{1}{2}c^2A' = 0 \quad (12)$$

Equating the proper acceleration \ddot{r} to the Newtonian acceleration a , would then give

$$\frac{1}{2}c^2A' = \frac{GM}{r^2} \quad (13)$$

and by integrating this differential equation one obtains

$$A = 1 - \frac{2GM/c^2}{r} \quad (14)$$

(where the constant of integration is obtained by setting $A = 1$ for $r \rightarrow \infty$). This then gives

$$\alpha = \frac{2GM}{c^2} \quad (15)$$

However, the above step that leads to A changing sign at $r = \alpha$ in Equation 14, which leads to the the black-hole prediction, is fundamentally incorrect, for the following reason. Whereas the radial equation of motion in *GR* (Equation 10) describes gravity as a combination of both time and radial space curvature, the spatial geometry of Newton's inverse square law of gravitation is strictly Euclidean, i.e. in Newton's law there is by definition no spatial curvature.

Thus, in order to compare Equation 10 rigorously with Newton's law, we must set $B = 1$ (not $B = 1/A$). The following differential equation is then obtained:

$$\ddot{r} + \frac{1}{2}(c^2 + \dot{r}^2)\frac{A'}{A} = 0 \quad [Newton; B = 1] \quad (16)$$

This differs crucially from Equation 12. Inserting Newtonian expressions for the acceleration and free-fall velocity then gives

$$-\frac{GM}{r^2} + \frac{1}{2}\left(c^2 + \frac{2GM}{r}\right)\frac{A'}{A} = 0 \quad (17)$$

and after some rearrangement one obtains

$$\frac{A'}{A} = \frac{\alpha/r}{(\alpha + r)} \quad (18)$$

where $\alpha = 2GM/c^2$, as before. Integrating this finally delivers the correct expression for $A(r)$, viz.

$$A = \left(1 + \frac{\alpha}{r}\right)^{-1} = \frac{r}{r + \alpha} \quad (19)$$

The curvature function A is now completely regular for $0 < r < \infty$ (see Figure 1).

Newton's inverse-square law of gravity can thus be thought of as describing that aspect of gravity caused exclusively by the curvature of the time coordinate, which is dominant for most cases we consider, such as planetary motion. However, space curvature becomes significant when speeds approach the speed of light, and distances to the central mass become small, and this will modify gravity from being purely Newtonian.

There is no reason to suppose that the time curvature (determined by Newton's law) would deviate from Equation 19 even in strong fields,




Figure 1: Time curvature A

but in order to obtain the space curvature (not described by Newton's law at all) we must now satisfy Einstein's *GR* field equations for the vacuum, viz.

$$A = \frac{1}{B} = 1 - \frac{\alpha}{\tilde{r}} = \left(1 + \frac{\alpha}{r}\right)^{-1} \quad (20)$$

which gives the following relationship between \tilde{r} and r :

$$\tilde{r} = r + \alpha \quad (21)$$

and $C = \tilde{r}^2 = (r + \alpha)^2$.

Next, we see that $A(r) = 1/B(r)$, since \tilde{r} and r are linearly related by Equation 7, and by integrating Equation 12 we then obtain for the proper velocity of free-fall:

$$\frac{\dot{r}^2}{c^2} = 1 - A$$

which gives here

$$\frac{\dot{r}^2}{c^2} = 1 - \left(1 + \frac{\alpha}{r}\right)^{-1} = \frac{\alpha}{r + \alpha}$$

or

$$\dot{r} = c \sqrt{\frac{\alpha}{r + \alpha}} \quad (22)$$

This means for $r \rightarrow 0$ we have $\dot{r} \rightarrow c$. On the other hand, Hilbert's solution predicts

$$\frac{\dot{r}^2}{c^2} = 1 - A = \frac{\alpha}{r}$$

or

$$\dot{r} = c\sqrt{\frac{\alpha}{r}} \quad (23)$$

which implies $\dot{r} \rightarrow c$ for $r \rightarrow \alpha$, and $\dot{r} \rightarrow \infty$ for $r \rightarrow 0$.

The small difference between \tilde{r} and r clearly becomes crucially significant the smaller r becomes; \tilde{r} never becomes smaller than α as $r \rightarrow 0$, meaning that A and B do not change sign at $r = \alpha$. Put another way, the spacetime manifold does not exist for $\tilde{r} < \alpha$, so Hilbert's extension into that region is invalid, and therefore an event horizon and black hole do not occur.




Figure 2: Effective gravitational mass

Furthermore, the radial free-fall acceleration may be written

$$\ddot{r} = -\frac{1}{2}c^2 \frac{\alpha}{(r + \alpha)^2} = -\frac{GM}{(r + 2GM/c^2)^2} \quad (24)$$

which shows classical Newtonian behaviour for $r \gg \alpha$ but deviates from inverse-square law behaviour for r of the order of α . The acceleration increases as $1/(r + \alpha)^2$ as $r \rightarrow 0$, from which an effective gravitational

mass M_{eff} may be defined as

$$\frac{M_{eff}}{M} = \frac{r^2}{(r + \alpha)^2} = \left(1 + \frac{\alpha}{r}\right)^{-2} \quad (25)$$

For large r , Newton's law is obeyed, but as $r \rightarrow 0$, the effective gravitational mass goes to zero (see Figure 2).




Figure 3: Due to angular curvature the point source appears as a ring

The proposed spacetime solution also suggests an interpretation for the ring-like images obtained by the so-called Event Horizon Telescope of sources at galactic centres. The function $C = \tilde{r}^2$ in the solution for a point mass is the square of the radial coordinate r if angular curvature is ignored, but if we note correctly that $\tilde{r} = r + \alpha$, we can construct a diagram such as in Figure 3 to show that the point source at $r = 0$ will be imaged as a ring at $\tilde{r} = 0$. Thus, the alleged "shadow of a black hole" is in reality the gravitationally lensed image of a regular point source.

3 Conclusion

Newton's law of gravitation relates solely to the contribution to gravity made by time curvature, space being Euclidean, and this fact must be used to link GR with classical physics. This means that the currently accepted black-hole solution is flawed fundamentally. Replacing it with a solution first mentioned in a paper by Brillouin, which I have expounded here, satisfies Einstein's vacuum field equations and agrees with all the usual predictions of GR , such as the bending of starlight

and the perihelion rotation of Mercury, but not the false prediction relating to black holes. The solution shows there is no horizon in space-time, because the Schwarzschild radial coordinate \tilde{r} is offset from the radial coordinate r by a distance $\alpha = 2GM/c^2$. This provides an explanation for the ringlike images of galactic centres published from EHT data [9]. The solution also shows that the velocity of a free-falling test object cannot exceed the speed of light, which is a result that would also be intuitively expected from the kinematics of special relativity.

The deductions outlined in this paper deviate markedly from the current paradigm. It would be satisfying if existing observations and calculations of trajectories that have previously led to the alleged presence of black holes could be re-examined on the basis of the model presented here, where there is no event horizon or black hole.

References

- [1] A.Einstein, Sitzungsberichte der Preussischen Akademie der Wissenschaften zu Berlin: 778, (11 November 1915)
- [2] K.Schwarzschild, *On the gravitational field of a point mass according to Einstein's theory*, Sitzung der Koeniglich Preussischen Akademie der Wissenschaft zu Berlin, Phys.-Math.189-196, (Jan.1916)
- [3] J.Droste, *The field of a single centre in Einstein's theory of gravitation*, Proc.K.Ned.Akad.Wet.19,197(1917)
- [4] H.Weyl, *Zur Gravitationstheorie*, Ann.Phys(Leipzig) 54,117(1917)
- [5] D.Hilbert, Math.Ann.92,1(1924), (also arXiv: physics/0310104)
- [6] L.S.Abrams, *Black holes: the legacy of Hilbert's error*, Can.J.Phys, 67,919(1989) (also arXiv:gr-qc/0201044v1 14 Jan 2002)
- [7] S.J.Crothers, *On the general solution to Einstein's vacuum field equations and its implications for relativistic degeneracy*, Progress in Physics, 1,68-73(2005)
- [8] M.Brillouin, *The singular points of Einstein's universe*, Journ Phys Radium, 23,43,(1923)
(see www.crothers.plasmareources.com/brillouin.pdf)
- [9] J.W.Moffat, *The Shadow of the Black Hole*, Oxford University Press (2020)