

**For calculating Nontrivial Zeros of Riemann Zeta function- $\zeta$ , the definition**

**$\xi(s) = \frac{s}{2}(s-1)\pi^{-s/2}\Gamma\left(\frac{s}{2}\right)\zeta(s)$  of Riemann Xi function- $\xi$  is not appropriate.**

**Author**

Ashok Kumar (Kashyap)

ORCID Id: <https://orcid.org/0000-0002-6345-7249>

Formerly, Assistant Professor of Physics

Department of Applied Sciences and Humanities

DEC- Faridabad, INDIA-121004

E-mail- [ak.research.ph@gmail.com](mailto:ak.research.ph@gmail.com)

Contact: +91-7557498649

**MSC-2020 Codes: 11M06, 11M26**

## ABSTRACT

We show that for calculating nontrivial zeros of the Riemann Zeta function  $\zeta$ , the form of the definition  $\xi(s) = (s/2)(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$ ,  $s \in \mathbb{C}$  of the function  $\xi$  and the followed

deduction that nontrivial zeros of functions  $\zeta(s)$  and  $\xi(s)$  are identical is not appropriate.

The definition of function  $\xi$  in which both functions  $\xi$  and  $\zeta$  are functions of same complex variable  $s$  and the assumption of identicalness of nontrivial zeros of  $\xi$  and  $\zeta$  is ambiguous, so

may be the deep reason, the Riemann hypothesis could not be resolved yet. However, the

definition  $\xi(t) = (s/2)(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$ ,  $t = \alpha + i\beta$  and  $s = \underline{1/2} + it$ , introduced by B.

Riemann (1859) leads the results: (i) when  $\xi(\alpha + i\beta) = 0$ ,  $\beta = 0$ ,  $\alpha \in \mathbb{R}$ , corresponding

nontrivial zero of the function  $\zeta(s)$  are of the form  $s = \underline{1/2} + i\alpha$  and (ii) when  $t = \alpha + i\beta$  and

$\xi(\alpha + i\beta) = 0$ , nontrivial zeros of the function  $\zeta(s)$  are of the form  $s = (\underline{1/2} - \beta) + i\alpha$  which lie

on both sides of the line  $\alpha = 1/2$ . Here, we sketch the zeros of the function  $\zeta(s)$  those

correspond to real zeros of the function  $\xi(s)$  that shows the Riemann hypothesis is true only

when nontrivial zeros of functions  $\xi(s)$  and  $\zeta(s)$  lie on two perpendicular lines.

**Keywords:** Zeta function, Riemann's Xi Function, nontrivial zeros, critical strip, critical line.

## 1 INTRODUCTION

In 1859, B. Riemann [1] in his research report introduced a function  $\zeta(s)$   $s = \sigma + it, \sigma, t \in \mathbb{R}$  known as the Riemann's zeta function  $\zeta(s)$  with the definition,

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \quad \dots (1)$$

Riemann further created another function known as the Riemann Xi-function  $\xi(t), t = \alpha + i\beta$  defined as:

$$\xi(t) = (s/2)(s-1)\pi^{-s/2}\zeta(s), s = \underline{1/2} + it \quad \dots (2)$$

The definition (2) is the **original definition** of the function  $\xi$ . But in Mathematics literature present day authors e.g. [2], [3], [4] and other use an alternative definition of function- $\xi$  as

$$\xi(s) = (s/2)(s-1)\pi^{-s/2}\zeta(s) \quad \dots (3)$$

With the definition (3) authors claim that nontrivial zeros of functions  $\xi(s)$  and  $\zeta(s)$  are identical.

In this research article, we show that the use of the definition (3) of the function- $\xi$  cannot be justified as it creates mathematical ambiguities. However, the original definition (2) of the function  $\xi$  corroborated with Riemann's statement: "it is clear that  $\xi(t)$  can vanish only if the imaginary part of  $t$  lies between  $i/2$  and  $-i/2$ ." which indicates that  $t$  is a complex number produces the results : (i) Corresponding to each complex zero  $t = \alpha + i\beta$  of the function  $\xi(t)$ , there exists a complex zero  $s = (1/2 - \beta) + i\alpha$  of the function  $\zeta(s)$ , i.e., zeros of functions  $\xi(t)$  and  $\zeta(s)$  are a distance apart (not identical). (ii) Corresponding to each real zero  $t = \alpha, \alpha \in \mathbb{R}$  of the function  $\xi(t)$ , there exists a complex zero  $s = \underline{1/2} + i\alpha$  of the function  $\zeta(s)$ . Perceived zeros of results (i) and (ii), are shown in Fig. 1(a) and 1(b) on the last page.

## 2 RESULTS

Recall the definition (3) connecting functions  $\xi$  and  $\zeta$  both of same complex variable  $s$ ,

$$\xi(s) = (s/2)(s-1)(\pi^{-s/2})\zeta(s), \quad s = \mu + i\lambda \quad \dots (4)$$

Clearly,  $\xi(0) = 0$  and  $\xi(1) = 0$ , so  $s = 0, s = 1$  are real zeros of  $\xi(s)$ . Suppose zeros of functions  $\xi(s)$  and  $\zeta(s)$  are identical, then  $s = 0, s = 1$  must also be zeros of  $\zeta(s)$  but according to definition (1) of  $\zeta(s)$ ,  $\zeta(0) = \infty$  and  $\zeta(1) = \infty$ , therefore,  $s = 0, s = 1$  are not zeros of  $\zeta(s)$ , so not of the function  $\xi(s)$ . That is ambiguity in definition (3). Actually, when  $s$  is a real number, all zeros of  $\zeta(s)$  necessarily are zeros of the function  $\xi(s)$  but when  $s$  is a complex number zeros of functions  $\xi(s)$  and  $\zeta(s)$  may be different can be shown as:

Suppose  $\xi = G + iH, (s/2)(s-1)(\pi^{-s/2}) = C + iD$  and  $\zeta(s) = A + iB$ , then from result (4),

$$G + iH = (CA - DB) + i(AD + BC) \quad \dots (5)$$

Zeros of  $\xi(s)$  can be obtained choosing  $G=0$  and  $H=0$  which means  $CA - DB = 0$  and

$AD + BC = 0$ . This system of equations produces  $A = 0, B = 0, A = iB, C = 0, D = 0$  and

$C = iD$ . Moreover, the function  $\zeta(s)$  can be written as  $\zeta(s) = \sqrt{A^2 + B^2} (\cos\phi + i\sin\phi)$  with

$\phi = \tan^{-1}\left(\frac{B}{A}\right)$ . Now, if  $\xi(s) = 0, (s/2)(s-1)(\pi^{-s/2}) \neq 0$ , then  $\sqrt{A^2 + B^2} (\cos\phi + i\sin\phi) = 0$

which implies the equation  $\zeta(s) = 0$  is unsolvable as when  $\sin\phi = 0 \Rightarrow \cos\phi \neq 0$ .

Further, suppose that  $s = a_i, a_i \in \mathbb{R}$  or  $\mathbb{C}, i = 1, 2, 3, \dots, n$  are zeros of the function  $\xi(s)$  and

$s = b_j, b_j \in \mathbb{R}$  or  $\mathbb{C}, j = 1, 2, 3, \dots, m$  zeros of the function  $\zeta(s)$ , i.e.,  $\xi(s) = \prod_{i=1}^n (s - a_i)$  and

$\zeta(s) = \prod_{j=1}^m (s - b_j)$ . Therefore, the result (4) can be expressed as,

$$\prod_{i=1}^n (s - a_i) = (s/2)(s-1)\pi^{-s/2}\Gamma(s/2) \prod_{j=1}^m (s - b_j) \quad \dots (6)$$

There are two cases:

**Case I:** At least one zero  $s = a$  (say) is common to both functions  $\xi$  and  $\zeta$  then,

$$\xi(s) = (s-a) \prod_{i=1}^{n-1} (s - a_i), \text{ and } \zeta(s) = (s-a) \prod_{i=1}^{m-1} (s - b_i), \text{ therefore,}$$

$$(s-a) \prod_{i=1}^{n-1} (s - a_i) = (s/2)(s-1)\pi^{-s/2}\Gamma(s/2)(s-a) \prod_{i=1}^{m-1} (s - b_i).$$

Further, write  $\prod_{i=1}^{n-1} (s - a_i) = \xi_1(s)$  and  $\prod_{i=1}^{m-1} (s - b_i) = \zeta_1(s)$ , then

$$\left. \begin{aligned} (s-a) \left[ \xi_1(s) - (s/2)(s-1)\pi^{-s/2}\Gamma(s/2)\zeta_1(s) \right] &= 0 \\ \xi_1(s) - (s/2)(s-1)\pi^{-s/2}\Gamma(s/2)\zeta_1(s) &\neq 0 \\ \left[ \xi_1(s) - (s/2)(s-1)\pi^{-s/2}\Gamma(s/2)\zeta_1(s) \right]_{s=a} &= \{0/(s-a)\}_{s=a} \\ \left[ \xi_1(a) - (a/2)(a-1)\pi^{-a/2}\Gamma(a/2)\zeta_1(a) \right] &= 0/0 \end{aligned} \right\} \quad \dots (7)$$

Thus there exists at least one case that for  $s = a$ ;  $\left[ \xi_1(a) - (a/2)(a-1)\pi^{-a/2}\Gamma(a/2)\zeta_1(a) \right]$  is not non-zero but indeterminate. But, in general  $\left[ \xi_1(a) - (a/2)(a-1)\pi^{-a/2}\Gamma(a/2)\zeta_1(a) \right]$  is considered a non-zero.

**Case II:** Functions  $\xi(s)$  and  $\zeta(s)$  have same number say  $p$  of identical zeros. Let  $\gamma$  be one of such zeros, then

$$\prod_{\gamma} (s - \gamma)^p \left[ 1 - (1/2)s(s-1)\pi^{-s/2}\Gamma(s/2) \right] = 0$$

$$\Rightarrow 1 - (1/2)\gamma(\gamma-1)\pi^{-\gamma/2}\Gamma(\gamma) = 0/0 \quad \dots (8)$$

If  $1 - (1/2)\gamma(\gamma-1)\pi^{-\gamma/2}\Gamma(\gamma)$  is nonzero then from result (8), either  $0 = 0$  or

$1 - (1/2)\gamma(\gamma-1)\pi^{-\gamma/2}\Gamma(\gamma)$  is indeterminate. Also, if  $\gamma$  equals 1, then  $1 = 0/0$  and if

$1 - (1/2)\gamma(\gamma-1)\pi^{-\gamma/2}\Gamma(\gamma)$  equals zero then  $0 = 0/0$ , i.e. 0 is itself indeterminate.

Whatever be the case I or II discussed above but even one common zero  $s = \alpha$  (say) results

$[\xi_1(\alpha) - (1/2)\alpha(\alpha-1)\pi^{-\alpha/2}\Gamma(\alpha/2)\zeta_1(\alpha)] = 0/0$  which shows 0 is not a free number, its use is

conditional. Thus, from the above discussion it can be concluded that (i) the definition (3) of

the function  $\xi$  is not a proper definition for calculating nontrivial zeros of the function  $\zeta(s)$

and (ii) to solve an equation like  $f(x) \times g(x) = 0$ ,  $f(x)$  or  $g(x) \in \mathbb{C}$ , the definition of zero

requires investigation because the conclusion from the equation  $X + iY = 0$ ,  $X, Y \in \mathbb{R}$

implies  $X = 0$  and  $Y = 0$  is not always true. The consideration  $\zeta(s) = X + iY = 0$  implies

$X = 0$  and  $Y = 0$  is the foremost reason; the Riemann hypothesis could not have been

resolved yet, also the claimed nontrivial zeros 14.134725142, 21.022039639, 25.010857580

and so on may not be nontrivial zeros of the function  $\zeta(s)$  but are of some other function.

That we will show elsewhere.

Now, using the definition  $\xi(t) = (s/2)(s-1)(\pi^{-s/2})\zeta(s)$ , we establish a relation between

nontrivial zeros of function  $\xi(t)$  and  $\zeta(s)$ ,  $s = \underline{1/2} + it$ .

Riemann states: "It is clear that  $\xi(t)$  can vanish only if the imaginary part of  $t$  lies between

$i/2$  and  $-i/2$ ." That suggests  $t$  is a complex variable. Suppose  $t = \mu + i\lambda$  (say) and

$\zeta(s)$ ,  $s = \underline{1/2} + it$ . Therefore, from the definition (3),

$$\xi(\mu+i\lambda) = (1/2)(1/2-\lambda+\mu i)(1/2-\lambda+\mu i-1)\pi^{-(1/2-\lambda+\mu i)/2}\Gamma[(1/2)(1/2-\lambda+\mu i)]\zeta(1/2-\lambda+\mu i)$$

Substitute, 0 for  $\mu$  and  $1/2$  for  $\lambda$  (or  $t = i/2$ )

$$\xi(i/2) = (1/2)(0)(-1+0i)\pi^0\Gamma[0]\zeta(0) = 0 \quad \dots (9)$$

Substitute, 0 for  $\mu$  and  $-1/2$  for  $\lambda$  (or  $t = -i/2$ )

$$\xi(-i/2) = (1/2)(1)(0)\pi^{-1/2}\Gamma[(1/2)(1)]\zeta(1) = 0 \quad \dots (10)$$

That shows  $t = -i/2$  and  $t = i/2$  are nontrivial zeros of the function  $\xi(t)$  but corresponding

to  $\xi(-i/2)$  and  $\xi(i/2)$  the values  $\zeta(0)$  and  $\zeta(1)$  are undefined. To avoid this ambiguity

Riemann states: “ $\xi(t)$  can vanish only if the imaginary part of  $t$  lies between  $i/2$  and  $-i/2$ ”.

The results (9) and (10) show if nontrivial zeros of function  $\xi(t)$  lie between  $t = -i/2$  to

$t = i/2$ , then corresponding zeros of the function  $\zeta(s)$  lie between  $s = 1$  to  $s = 0$ . Thus, the

range of nontrivial zeros of the function  $\zeta(s)$  is  $s \in [0, 1]$  which is the critical strip for

nontrivial zeros of function  $\zeta(s)$ .

The critical strip for nontrivial zeros of  $\zeta(s)$  can also be determined as:

Suppose  $t = \alpha \pm i\beta$  are zeros of the function  $\xi(t)$ , then according to Riemann's statement,

$$\begin{aligned} -i/2 \leq t \leq i/2 \\ \Rightarrow i^2 1/2 \leq -it \leq -i^2 1/2 \Rightarrow -1/2 \leq -i \alpha \pm i\beta \leq 1/2 \\ \Rightarrow -1/2 \leq -i\alpha \mp \beta \leq 1/2 \Rightarrow 1/2 \geq i\alpha \pm \beta \geq -1/2 \\ \Rightarrow 1 \geq 1/2 \pm \beta + i\alpha \geq 0 \Rightarrow 0 \leq 1/2 \pm \beta + i\alpha \leq 1 \end{aligned}$$

But  $1/2 \pm \beta + i\alpha$  is variable of the function  $\zeta(s)$  corresponding to  $t = \alpha \pm i\beta$ . Thus, if zeros

of function  $\xi(t)$  lie between  $t = -i/2$  to  $t = i/2$ , then zeros of the function  $\zeta(s)$  lie between

$s = 0$  to  $s = 1$ .

Thus, nontrivial zeros of the function  $\zeta(s)$  are of the form  $\frac{1}{2} \mp i\beta$  that lie in the region  $0 \leq \frac{1}{2} \mp \beta \leq 1$  that verbalize the Riemann hypothesis. Further, if  $\beta$  equals zero, i.e. all zeros of function  $\xi(t = \alpha \pm i\beta)$  are real then zeros of  $\zeta(s)$  are of the form  $\frac{1}{2} + i\alpha$  that lie in the region  $0 \leq \frac{1}{2} \leq 1$  on the line  $a = 1/2$ . Clearly, the functions  $\xi(t = \alpha \pm i\beta)$  and  $\zeta(s = \frac{1}{2} + it)$  have same number of zeros and there is one-to-one correspondence between real zeros of the function  $\xi(t)$  and nontrivial complex zeros of the function  $\zeta(s)$ .

The perceived (not calculated) nontrivial zeros of functions  $\xi(t)$  and  $\zeta(s)$  when (i)  $t \in \mathbb{C}$  a complex number, and (ii), when  $t$  is real number are shown in Fig. 1(a) and Fig. 1(b) respectively. **Note:** Here, for to show the relative locations of zeros of the function  $\zeta(s)$ , zeros of the function  $\xi(t)$  are perceived (not calculated).




Fig. 1(a): Zeros of functions  $\xi(t)$  and  $\zeta(s)$  when  $t$  is a complex variable




Fig. 1(b): Zeros of functions  $\xi(t)$  and  $\zeta(s)$  when  $t$  is a real variable



Thus, if  $t = \alpha \pm i\beta$  is zero of the function  $\xi(t)$ , then corresponding zero of the function  $\zeta(s)$

is  $s = \left(\frac{1}{2} \mp \beta\right) \pm i\alpha$ . That show zeros of functions  $\xi$  and  $\zeta$  cannot have same form and same

variable and in the context of the Riemann hypothesis the form of definition of function  $\xi$

$\xi(s) = (s/2)(s-1)(\pi^{-s/2})\zeta(s)$ ,  $s = \mu + i\lambda$  is ambiguous.

### Acknowledgement

The author is grateful to M. Srinivasan Ramanujan (Mathematician 1887-1920), Albert Einstein (Physicist 1879-1955) whose early life struggle and odds skilled him how to remain academically steady in odd livelihood conditions.

### References

- [1] Riemann, B. (1859), Über die Anzahl der Primzahlen unter einer gegebenen Grösse Monatsberichte der Berliner Akademie, November 1859, pp. 671-680. English translation “on the number of Primes Less Than a Given Magnitude,” in the Appendix of [3], pp. 299-305.
- [2] Bombieri E (2000). The Riemann Hypothesis- official problem description, Clay Mathematics Institute.
- [3] Titchmarsh, E.C. (1974). The Theory of the Riemann Zeta-Function. OUP.
- [4] Edwards, Harold M. (2001), Riemann’s Zeta Function, Dover ed. (2001).
- [5] Apostol T.M. Introduction to Analytical Number Theory, Springer, 5<sup>th</sup> Printing 1998, Edition (1976).
- [6] H. Davenport. Multiplicative Number Theory, Volume 74, Springer-Verlag, Second Edition (1980).

### Declaration

The Author does not have any compelling interest writing this research article. The Author communicates this research article through this pre-print repository to share the knowledge to the interested audience.

**Additional Information:** Corresponding to non-trivial zero  $\alpha + i\beta$  of the function  $\xi$ , non-trivial zero of the function  $\zeta$  is  $\left(\frac{1}{2} - \beta\right) + i\alpha$ .




Fig. 1(a): Zeros of functions  $\xi(t)$  and  $\zeta(s)$  when  $t$  is a complex variable




Fig. 1(b): Zeros of functions  $\xi(t)$  and  $\zeta(s)$  when  $t$  is a real variable