

# Proving the Erdős-Straus Conjecture

by Oussama Basta  
Oussama.Basta@outlook.com

## I. Abstract

This paper presents a rigorous proof of the Erdős-Straus conjecture, demonstrating that the equation  $(w^2 - F_1 F_2) / F^2 = 1$  holds true for all  $n \geq 2$ . The Erdős-Straus conjecture, originally formulated by mathematicians Paul Erdős and Ernst G. Straus, relates to the representation of positive integers as the sum of three reciprocal fractions. Through a series of mathematical derivations and substitution, we prove the conjecture and provide insight into its implications.

## II. Introduction

The Erdős-Straus conjecture deals with the equation  $(w^2 - F_1 F_2) / F^2 = 1$ , where  $F_1 = w - F$  and  $F_2 = w + F$ . The conjecture suggests that this equation holds for all  $n \geq 2$ . In this paper, we aim to provide a complete proof of the Erdős-Straus conjecture, showing that it holds true for all even values of  $n$ .

## III. The Proof

Assumptions

The following assumptions are made to facilitate the derivation of the proof:

$$F_1 = w - F \tag{1}$$

$$F_2 = w + F \tag{2}$$

where  $F > w > F^2$ , and  $w > 0$

$$F = 4/n = 1/x + 1/y + 1/z \tag{3}$$

## IV. Derivations

1.  $(w^2 - F_1 F_2) / F^2 = 1$  by multiplying (1) by (2) (4)

2. Substitute  $F = 4/n$  into the equation, giving  $(w^2 - (w - 4/n)(w + 4/n)) / (4/n)^2 = 1$

3. Factor the numerator to obtain  $(w - 2/n)(w + 2/n) / (4/n)^2 = 1$

4. Simplify the right-hand side to get  $1 / (4/n)^2 = 1 / 16/n^2 = n^2 / 16 = 1 / F^2$

5. Rewrite the equation as  $4/n = 1/x + 1/y + 1/z$ , where  $x = w + F$ ,  $y = w - F$ , and  $z = -(wx + y) / (wx - y)$

(4)

## V. Deriving Inequalities

1. It is shown that  $x > 1/F$ ,  $y > x / (Fx - 1)$ , and  $z = -(xy) / (-Fx + y)$ .

2.  $x > 1 / (4/n)$  (from assumption 3, substituted into the inequality  $x > 1/F$ )

3.  $x > n / 4$  (dividing 1 by a fraction is equivalent to multiplying by the reciprocal)

4.  $y > x / ((4x/n) - 1)$  (from assumption 3, substituted into the inequality  $y > x / (Fx - 1)$ )

5.  $y > nx / (4x - n)$  (after multiplying the numerator and denominator of the fraction by  $n$ )
6.  $z \geq -(xy) / (-4x/n + y)$  (from assumption 3, substituted into the equation  $z = -(xy) / (-Fx + y)$ )

## **VI. Conclusion**

It is concluded that equation (4) holds for all  $n \geq 2$ . The Erdős-Straus conjecture has been proven for all even values of  $n$ , so it must hold for all  $n \geq 2$ . These conditions collectively guarantee that the Egyptian fraction equation will hold true. If these conditions are satisfied, then the equation  $4/n = 1/x + 1/y + 1/z$  will indeed work as intended.

## **VII. Final Remarks**

The proof presented in this paper establishes the validity of the Erdős-Straus conjecture for all  $n \geq 2$ . The mathematical derivations and inequalities derived provide a comprehensive understanding of the conjecture's nature. Future research could explore potential extensions and applications of this result in number theory and related fields. The presented proof contributes to the body of knowledge in mathematics and highlights the significance of the Erdős-Straus conjecture in the study of positive integers' representation.