

On real flat quantum-spacetime linelement from first principles

Holger Döring
DPG-departement matter and cosmos
section: GRT and gravity
IQ-Wissen Berlin
Germany
e-mail: holgerdoering@alumni.tu-berlin.de

Abstract:

In an uncomplicated way there can be derived a quantum-lineelement of local tangent-spacetime of Minkowski-type from a fundamental form of physical announcements. This term includes a variable, which can be interpreted as combination of Ricci-scalar and cosmological constant. In this method, the Planck-constant is included in the lineelement-form in a natural way. If this ansatz is seen as real from first principles, then classical Minkowski-lineelement is wrong. So since it is the limiting case for GRT without gravity, quantum gravity-equations also must be false and be modified. Local invariant Lorentz-Planck-transformations can be derived for some special conditions of velocity values. This explorations for quantum in tangential-spacetime may lead to a deeper understanding of quantizing gravity.

Key-words: Planck-Lorentz-transformation; Planck-Minkowski-lineelement; quantized flat spacetime; Ricci-scalar; scalar-curvature; cosmological constant; minimal Planck-length; first principles; quantizing uncurved spacetime.

1.Introduction:

There are two methods constructing a quantum-spacetime-theory from classical limits. Going from $0 \Rightarrow \hbar$ for quantum states against continuity and going from $\gamma \Rightarrow G_{\mu,\nu}$ for gravity and curvature from Newton-model to Einstein Tensor-description and then going from zero to \hbar . The first means going from flat Minkowski spacetime to a sort of quantized tangential spacetime and the second to go from spacetime without curvature to GRT with gravity-force and then to quantum \hbar . Mostly the second trying is used. [2],[3].

This paper here only constructs a quantum flat Planck- Minkowski- spacetime from fundamental Planck-length-squares. If this attempt will be physically, logically and mathematically consistent and succesful, must be seen. From this level then could be constructed a consistent quantum gravity, which limiting case this construction here would be.

Citation Wheeler [1.]: *"Space-time geometry is no longer high above the battle of matter and energy. It takes part in the struggle. Geometry tells matter how it should move, but mass in turn dictates the curvature to geometry."*

This is also a fact for quantum tangential spacetime, where is shown, that a gravityfree local spacetime TM of a manifold M isn't really flat and empty but depends on some physical variables.

2.Calculation:

From first principles is defined:

$$\text{If } \frac{1}{r^2_{PL}} := R_{Fund} \pm \Lambda := \frac{m^2_{PL} \cdot c^2}{\hbar^2} , \quad (1)$$

where R is the Ricci-scalar curvature and Λ is cosmological constant, then this definition can be developed to its general case of:

$$R \pm \Lambda = \frac{m^2 \cdot c^2}{\hbar^2} \quad (2.)$$

and of course R is defined over the two curvature-mainaxes (in two dimensions) as scalar invariance of curvature:

$$R := \left| \frac{2}{\rho_1 \cdot \rho_4} \right| . \quad (3.)$$

This leads to a modification of Einsteins local energy-equation for flat spacetime of:

$$E = m \cdot c^2 - \frac{\hbar^2 \cdot (R \pm \Lambda)}{m} \quad (4.)$$

This result leads directly to corrected line-element of quantized Planck-Minkowski-space in its new form of:

$$ds^2 = x^2 \cdot \frac{\hbar^2 \cdot (R \pm \Lambda)}{c^2 \cdot m_0^2} - c^2 \cdot t^2 \cdot \left(\frac{\hbar^2 \cdot (R \pm \Lambda)}{c^2 \cdot m_0^2} - 1 \right) \quad (5a.)$$

which also can be written as:

$$ds^2 = c^2 \cdot t^2 + \frac{\hbar^2 \cdot (R \pm \Lambda)}{c^2 \cdot m_0^2} \cdot (x^2 - c^2 \cdot t^2) \quad (5b.)$$

As is seen, the timelike coordinate is quantized and exists in a classical form, which gives this system of description a similiar partial form of a bimetric whereby the spacelike dimension-coordinate appears only in a quantum description.

This equation (5a./5b.) is from now on called „Planck-Minkowski-lineelement“.

As is seen, this new lineelement is conform to classical Minkowski- description of local spacetime and differs from it only for the conformal term of:

$$K := \frac{\hbar^2 \cdot (R \pm \Lambda)}{c^2 \cdot m_0^2} . \quad (6.)$$

In this form the new local lineelement can be written as:

$$ds^2 = c^2 \cdot t^2 + K \cdot (x^2 - c^2 \cdot t^2) \quad (7a.)$$

or

$$ds^2 = K \cdot x^2 - c^2 \cdot t^2 \cdot (K - 1) \quad , \quad (7b.)$$

which comes from [4.]:

$$ds^2 = d x^i \cdot dx^k \cdot \eta_{i,k}$$

with to the classic Minkoski tensor varied compared metric fundamental tensor for uncurved tangential spacetime :

$$\eta_{i,k} = \begin{pmatrix} K & 0 \\ 0 & 1 - K \end{pmatrix} \quad (7c.)$$

This term leads directly to the Planck-Lorentz-transformations, described with its transformation-matrix (formulated for two dimensions) extended from [10.]:

$$A = \begin{pmatrix} \sqrt{K} & \sqrt{K} \cdot \beta \\ \sqrt{K} \cdot \beta & \sqrt{K} \end{pmatrix} \quad (8.)$$

$$\text{with } \begin{pmatrix} x' \\ c \cdot t' \end{pmatrix} = \begin{pmatrix} x \\ c \cdot t \end{pmatrix} \cdot A \quad (9.)$$

This shows directly the Planck-Lorentz-transformations for two dimensions of uncurved but quantized local spacetime in its conformal form:

$$\begin{aligned} x' &= \sqrt{K} \cdot (x + \beta \cdot c \cdot t) \\ c \cdot t' &= \sqrt{K} \cdot (x \cdot \beta + c \cdot t) \end{aligned} \quad , \text{with } \beta = \frac{v}{c} \quad (10a./10b)$$

where

$$\sqrt{K} := \frac{\hbar}{c \cdot m_0} \cdot \sqrt{R \pm \Lambda} \quad (11.)$$

Since there is the Planck- condition $\sqrt{K} = \gamma = \left(1 - \frac{v^2}{c^2}\right)^{-\frac{1}{2}}$ to Lorentz-factor, this leads to the corrected maximal limiting condition in physical velocity for material fermion bodies of:

$$v \leq c \cdot \sqrt{1 - \frac{c^2 \cdot m_0^2}{\hbar^2 \cdot (R \pm \Lambda)}} \quad (12.a)$$

instead of:

$$v \leq c \quad . \quad (12b.)$$

3.Conclusion:

It is possible to construct a flat quantum-spacetime in Minkowski-like coordinates with its local quantum invariance conditions for coordinate transformations. If this description would be true, so every real quantum-gravity must lead in its limit without gravity- force to this Planck-Minkowski-metric, like classical gravity of GRT must lead without force in its limit to classical flat spacetime of Minkowski-lineelement. In all formulas above either Ricci-scalar term R or cosmological constant Λ can be set to zero. If both are set to zero, classical, local Minkowski spacetime remains at tangential spacetime to manifold M . Also the local geometric invariance form is reviewed: the light cone will change its form a little bit, which size-changing depends from now on, controlled by R, Λ and m_0 . Not only in gravity-determined spaces [6.],[7.] but even in tangential spacetime, it is no longer a static form [8.],[9.].

4.Summary:

From first principles of a fundamental length-square with identifying square of inverse Planck-length with Ricci-scalar and cosmological constant there can be constructed a local uncurved quantum-spacetime without gravity as a corrected form of tangential spacetime TM_Q for manifold M . But this construction will include Ricci curvature scalar R and may lead to a consistent form of a description of quantum-gravity which in limit for zero gravity is this description of quantum tangential spacetime mentioned above.

5.Acknowledgements:

Many thanks to Prof. Philip E.A. Mortimer (PhD), former member of university of Glasgow and Edinburgh (Edinburgh) for help- and colourful ideas, discussions and suggestions.

6.References:

- . [1.] Wheeler, J. A., Einsteins Vision, Springer Verlag Berlin-Heidelberg-New York, 1968.
- . [2.] Döring, H.A.W., some brief qualitative letter-comments on quantizing gravity: thirteen hypotheses.2023.hal-04278409.<https://hal.science/hal-04278409>
- . [3.] Giulini, D., Kiefer, C., Lämmerzahl, C., (eds.): Lecture Notes in Physics, quantum-gravity – from theory to experimental research. Springer-Verlag, Berlin-Heidelberg-New York, 2003
- . [4.] Minkowski, H., Raum und Zeit, Vortrag; Leipzig und Berlin, Druck und Verlag von B.G. Teubner 1909
- . [5.] Misner, C.W., Thorne, K.S., und Wheeler, J.A.: *Gravitation*. Freeman, San Francisco 1973, ISBN 0-7167-0334-3.
- . [6.] Penrose, R.: Techniques of Differential Topology in Relativity. Society for Industrial and Applied Mathematics (SIAM): CBMS-NSF Regional Conference Series in Applied Mathematics, 1972; ISBN 0-89871-005-7, [doi:10.1137/1.9781611970609](https://doi.org/10.1137/1.9781611970609).

- . [7.] Minguzzi,E.: Lorentzian causality theory. In: Living Reviews in Relativity, Band 22, Nr.3, 3. Juni 2019; [doi:10.1007/s41114-019-0019-x](https://doi.org/10.1007/s41114-019-0019-x).
- . [8.] Papadopoulos, K., Acharjee,S., Papadopoulos,B.,K.: *The order on the light cone and its induced topology*. In: *International Journal of Geometric Methods in Modern Physics*. 15. Jahrgang, Nr.5, 1.Mai 2018, S.1850069–18](arxiv.org [PDF]).
- . [9.] Giulini, D.: [Globale versus lokale Strukturen von Raum-Zeiten](#). Tutorium der AGjDPG, DPG-Frühjahrstagung 2017, Bremen, 13. März 2017.
- . [10.] Einstein, A.: Zur Elektrodynamik bewegter Körper, *Annalen der Physik*,**322**,10,1905. doi:<https://doi.org/10.1002/andp.19053221004>

7. Verification:

This paper is written without help from a chatbot like Chat-GPT4 or other AIs. It's fully human work.

January 2024

