


# A Geometric Algebra solution to a Contest Problem

April 22, 2024

James Smith  
LinkedIn group ("Pre-University Geometric Algebra")

## Abstract

We show how to use rotations of vectors in GA to solve the following problem: "The following are known about a triangle: The ratio of the lengths of two sides; the angle formed by those sides; and the length of that angle's bisector. Find the length of the side opposite that angle."



Given  $\alpha$ ,  $d$ , and  $k$  (but not  $l$ ), find the length  $BC$ .




Figure 1: Given  $\alpha$ ,  $d$ , and  $k$  (but not  $l$ ), find the length  $BC$ .


## 1 Statement of the Problem

The problem that we solve here is a generalization of <https://www.youtube.com/watch?v=BeuLmUjPFsk>. We state that generalization as: Given  $\alpha$ ,  $d$ , and  $k$  (but not  $l$ ), find the length  $BC$  (Fig. 1).

## 2 Ideas that We Will Use

1. For any two parallel vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\mathbf{u} \wedge \mathbf{v} = 0$ .
2. For any two vectors  $\mathbf{u}$  and  $\mathbf{v}$ ,  $\mathbf{u} \wedge \mathbf{v} = \langle \mathbf{u}\mathbf{v} \rangle_2$ .
3. Ideas concern the rotation of a vector  $\mathbf{v}$  that is parallel to a given bivector  $\mathbf{i}$ :

- (a) The vectors  $\mathbf{v}e^{i\theta}$  and  $\mathbf{v}e^{-i\theta}$  are rotations of  $\mathbf{v}$  by the same angle  $\theta$ , but in opposite directions.



- (b) For any angle  $\theta$ , and any vector  $\mathbf{v}$  that is parallel to bivector  $\mathbf{i}$ ,  $\mathbf{v}e^{i\theta} = e^{-i\theta}\mathbf{v}$ . Here is a proof:




Figure 2: Formulation of the problem in terms of vectors. Note that  $\mathbf{b} = kl\hat{\mathbf{d}}e^{-i\alpha}$ ;  $\mathbf{d} = d\hat{\mathbf{d}}$ ; and  $\mathbf{c} = l\hat{\mathbf{d}}e^{i\alpha}$ .

$$\begin{aligned}
 \mathbf{v}e^{i\theta} &= \mathbf{v} [\cos \theta + \mathbf{i} \sin \theta] \\
 &= \mathbf{v} \cos \theta + \mathbf{v}\mathbf{i} \sin \theta \\
 &= \mathbf{v} \cos \theta - \mathbf{i}\mathbf{v} \sin \theta \\
 &= [\cos \theta - \mathbf{i} \sin \theta] \mathbf{v} \\
 &= e^{-i\theta} \mathbf{v}.
 \end{aligned}$$

(c) From  $\mathbf{v}e^{i\theta} = e^{-i\theta} \mathbf{v}$ , we can see that  $\mathbf{v}e^{i\theta} \mathbf{v} = v^2 e^{-i\theta}$ , and that  $\hat{\mathbf{v}}e^{i\theta} \hat{\mathbf{v}} = e^{-i\theta}$ .

### 3 Solution Strategy

We will use rotation of vectors to determine  $l$ , then (knowing  $l$ ) we will use the Law of Cosines to calculate  $BC$ .

### 4 Formulation in Terms of Vectors

We formulate the problem as shown in Fig. 2.

## 5 Solution

We begin by recognizing that  $[\mathbf{c} - \mathbf{d}] \parallel [\mathbf{d} - \mathbf{b}]$ . Therefore,  $[\mathbf{c} - \mathbf{d}] \wedge [\mathbf{d} - \mathbf{b}] = 0$ , and

$$\hat{\mathbf{d}}e^{i\alpha}\hat{\mathbf{d}} = e^{-i\alpha}.$$

$$\langle d^2 \rangle_2 = 0.$$

$$\begin{aligned} \langle [\mathbf{c} - \mathbf{d}] [\mathbf{d} - \mathbf{b}] \rangle_2 &= 0 \\ \langle [l\hat{\mathbf{d}}e^{i\alpha} - d\hat{\mathbf{d}}] [d\hat{\mathbf{d}} - kl\hat{\mathbf{d}}e^{-i\alpha}] \rangle_2 &= 0 \\ \langle ld\hat{\mathbf{d}}e^{i\alpha}\hat{\mathbf{d}} - kl^2\hat{\mathbf{d}}e^{i\alpha}\hat{\mathbf{d}}e^{-i\alpha} - d^2 + kdle^{-i\alpha} \rangle_2 &= 0 \\ \langle lde^{-i\alpha} - kl^2e^{-i\alpha}e^{-i\alpha} + kdle^{-i\alpha} \rangle_2 &= 0 \\ \langle de^{-i\alpha} - kle^{-2i\alpha} + kde^{-i\alpha} \rangle_2 &= 0 \\ -d \sin \alpha + kl \sin 2\alpha - kd \sin \alpha &= 0 \\ \therefore l &= \frac{(k+1)d \sin \alpha}{k \sin 2\alpha}. \end{aligned}$$

Because  $\sin 2\alpha = 2 \sin \alpha \cos \alpha$ ,

$$l = \frac{(k+1)d \sin \alpha}{2k \cos \alpha}.$$

Finally, from the Law of Cosines,

$$\begin{aligned} BC &= \sqrt{AB^2 + AC^2 - 2(AB)(AC) \cos 2\alpha} \\ &= \sqrt{(kl)^2 + l^2 - 2kl^2 \cos 2\alpha} \\ &= l\sqrt{k^2 + 1 - 2k \cos 2\alpha} \\ &= \left[ \frac{(k+1)d \sin \alpha}{2k \cos \alpha} \right] \sqrt{(k+1)^2 - 2k(1 + \cos 2\alpha)} \\ &= \left[ \frac{(k+1)d \sin \alpha}{2k \cos \alpha} \right] \sqrt{(k+1)^2 - 4k \cos^2 \alpha}. \end{aligned}$$

For any angle  $\theta$ ,

$$\cos \frac{\theta}{2} = \sqrt{\frac{1 + \cos \theta}{2}}.$$