

Vector kinematic analysis of Keplerian velocity from Hamilton's hodograph

Hervé Le Cornec*

Dated 26/07/2025

Abstract: Many authors in the literature focused on Hamilton's hodograph of the Keplerian motion but none of them developed a vector kinematic analysis of the peculiar velocity shown by the hodograph, so we perform it here. As expected, the analysis predicts Kepler's laws, the mathematical structure of Newton's acceleration, and it also leads to new expressions for the classical mechanical energy and the eccentricity. We then discuss the relationship between the geometric acceleration given by the kinematics and the physical one stated by Newton.

Introduction

In 1845 W.R. Hamilton proposed a new way of representing the motion of a body submitted to Newton's gravitation law, the hodograph^[1]. Many authors have since discussed this approach^[2-8], some of them giving figures of this geometric problem that lacked in Hamilton's communication. Among them is E.I. Butikov who pictured a most interesting property of the Keplerian motion, in the figure 4 of one of his article^[8]: the velocity of a Keplerian orbiter is always the simple addition of a constant rotation velocity, that he called \mathbf{u} while we will call it \mathbf{v}_R , and a constant translation velocity, that he called \mathbf{w} while we will call it \mathbf{v}_T . The overall velocity is then given by :

$$\mathbf{v} = \mathbf{v}_R + \mathbf{v}_T \quad (1)$$

Because the authors were focused on the structure of the hodograph rather than the kinematics of this peculiar velocity, we could not find in the literature any vector kinematic analysis of this very simple vector expression, therefore we decided to develop one.

We will show that, as expected, we are able to confirm that such a velocity is in full agreement with the three laws of Kepler and with the mathematical expression of Newton's gravitational acceleration. Nonetheless, Newton's constant physical factor GM (G being the constant of gravitation and M the mass at the focus of the orbit) is replaced by another constant factor purely kinematic, which is trivial as far as the above equation (1) embeds no physical parameter but only geometric ones.

We are then driven to discuss the equality between the kinematic acceleration and Newton's physical one.

Kinematic analysis

As explained in the introduction, the velocity \mathbf{v} of any Keplerian orbiter is simply the vector addition of two uniform velocities, one of rotation, \mathbf{v}_R , plus one of translation, \mathbf{v}_T (see equation 1). Its simplest mathematical expression is then as follows:

$$\mathbf{v} = \mathbf{v}_R + \mathbf{v}_T$$

with

$$\begin{aligned} \mathbf{v}_R &= \boldsymbol{\omega} \times \mathbf{r} \\ \mathbf{v}_R &= \|\mathbf{v}_R\| = \omega r = \text{constant} \\ \mathbf{v}_T &= \text{constant} \end{aligned} \quad (2)$$

In this expression $\boldsymbol{\omega}$ is the vector frequency of rotation, perpendicular to the plane of the orbit, and \mathbf{r} is the vector radius, from the focus of the orbit to the orbiter. Note that \mathbf{v}_T and \mathbf{v}_R are coplanar all along the orbit. Take care, in this expression the index R means "rotation" but not "radial", while the index T stands for "translation" but not "tangential". The figure 1 shows these two velocities on a typical Keplerian orbit.




Figure 1 : the Keplerian velocity \mathbf{v} decomposed into its rotation velocity \mathbf{v}_R and its translation velocity \mathbf{v}_T .

Now let us demonstrate that coming from this definition of the orbital velocity we can predict the existence of Kepler's laws as well as Newton's acceleration, or at least its mathematical structure.

The first consequence of the above expression (2) is the validity of the following one by derivation of \mathbf{v}_R with respect to time ($\boldsymbol{\omega}$ and $\dot{\boldsymbol{\omega}}$ being colinear) :

$$\dot{\boldsymbol{\omega}} \mathbf{r} = -\dot{r} \boldsymbol{\omega} \quad (3)$$

From the relations (2) and (3) we can then calculate the acceleration which is the derivative of the velocity with respect to time :

$$\mathbf{a} = \dot{\boldsymbol{\omega}} \times \mathbf{r} + \boldsymbol{\omega} \times \mathbf{v} = -\frac{\dot{\boldsymbol{\omega}}}{r^2} \times (\mathbf{r} \times (\mathbf{r} \times \mathbf{v})) \quad (4)$$

Defining the massless angular momentum as

$$\mathbf{L} = \mathbf{r} \times \mathbf{v} \quad (5)$$

the final expression of the acceleration is given by :

* herve.le.cornec@free.fr

$$\mathbf{a} = -\frac{k}{r^3} \mathbf{r} \quad \text{with} \quad k = L v_R = \text{constant} \quad (6)$$

$$L = r^2 \dot{\theta} \quad (14)$$

Therefore, the acceleration and the vector radius are colinear and this forces the angular momentum to be a constant, because $\dot{\mathbf{L}} = \mathbf{r} \times \mathbf{a} = \mathbf{0}$, as awaited for a central field motion :

$$\mathbf{L} = \text{constant} \quad (7)$$

Note that the expression (6) of the acceleration is mathematically identical to the one of Newton, at the condition that Kepler's constant k verifies:

$$GM = L v_R = k \quad (8)$$

Now from this we observe that the vector product of the rotation velocity with the angular momentum leads trivially to :

$$\mathbf{v}_R \times \mathbf{L} = v_R^2 \left(1 + \frac{\mathbf{v}_R \cdot \mathbf{v}_T}{v_R^2} \right) \mathbf{r} \quad (9)$$

The scalar version of this equation is therefore :

$$p = (1 + e \cos \theta) r$$

$$\text{with} \quad p = \frac{L}{v_R} = \frac{k}{v_R^2} \quad \text{and} \quad e = \frac{v_T}{v_R} \quad (10)$$

This is the equation of a conic where p is the semi-latus rectum, e is the eccentricity and θ is the true anomaly, i.e. the angle between \mathbf{v}_T and \mathbf{v}_R which is also the angle between the direction of the perigee and the vector radius. This is the expression of Kepler's first law^[9].

Now the vector multiplication of \mathbf{v} and \mathbf{L} leads to $\mathbf{v} \times \mathbf{L} = \mathbf{v}_T \times \mathbf{L} + \mathbf{v}_R \times \mathbf{L} = \mathbf{v}_T \times \mathbf{L} + k \mathbf{r}/r$, and thus to

$$\frac{\mathbf{v}_T \times \mathbf{L}}{k} = \frac{\mathbf{v} \times \mathbf{L}}{k} - \frac{\mathbf{r}}{r} \quad (11)$$

The right hand side of this equation is known as the eccentricity vector^[10], that can be thus given by a new expression :

$$\mathbf{e} = \frac{\mathbf{v}_T \times \mathbf{L}}{L v_R} \quad (12)$$

Therefore the translation velocity is always perpendicular to the main axis of the conic, which direction is the one of the vector eccentricity. The figure 1 exhibits both the rotation and the translation velocities at different positions on a conic.

Let us now notice that the scalar multiplication of the total velocity and the vector radius leads to :

$$\mathbf{r} \cdot \mathbf{v} = \mathbf{r} \cdot \mathbf{v}_T = r \dot{r} \quad \text{thus} \quad \dot{r} = v_T \sin \theta \quad (13)$$

Using this last expression associated to the time derivation of the conic equation (10), it is trivial to show that the angular momentum can be presented as the multiplication of the square of the vector radius and the derivative of the true anomaly with respect to time :

This last expression is very well known, being described for instance by L. Landau and E. Lifchitz in their course "Mechanics"^[11]. It shows that the areal velocity, defined as $\dot{f} = r^2 \dot{\theta}/2 = L/2$, must be a constant as far as the angular momentum also is. Therefore the expression (14) is nothing else but the second law of Kepler^[9].

Note that the time derivative of the true anomaly $\dot{\theta}$ and the frequency of rotation ω are then related by the following formula :

$$\dot{\theta} = \omega (1 + e \cos \theta) \quad \text{or} \quad r \dot{\theta} = p \omega \quad (15)$$

Now integrating the expression (14) over a complete period T of revolution for an ellipse (see L. Landau and E. Lifchitz^[9]), knowing that the surface of an ellipse is $f = \pi a b$, a being the semi major axis and $b = \sqrt{a p}$ the minor one, we can integrate as so :

$$\int_0^T L dt = L T = \int_0^T 2 \dot{f} dt = 2 f = 2 \pi a \sqrt{a p} \quad (16)$$

Then using the equations (10), we are trivially led to the following formula :

$$L v_R = 4 \pi^2 \frac{a^3}{T^2} = k \quad (17)$$

This is the expression of Kepler's third law^[9].

In addition, we can see a deep connection between this Keplerian kinematics and the classical mechanics. Indeed by calculating the square of the velocity (2) it is trivial to get the following relationship :

$$\frac{1}{2} v^2 - \frac{k}{r} = \frac{1}{2} v_R^2 (e^2 - 1) = \text{constant} \quad (18)$$

If we multiply this expression by the mass m of the orbiter, we get its classical mechanical energy^[9], which is therefore :

$$E_M = \frac{1}{2} m v^2 - \frac{mk}{r} = \frac{1}{2} m v_R^2 (e^2 - 1) \quad (19)$$

This expression of the classical energy by the means of the rotation velocity and the eccentricity has never been reported at our knowledge.

Discussion

As expected, the kinematic analysis of the velocity derived from Hamilton's hodograph of the Keplerian motion predicts Kepler's laws as well as the mathematical structure of Newton's acceleration.

In addition it gives two results that have not been described yet in the literature. The first one is the expression (12) of the eccentricity vector, so also of the Runge-Lenz vector^[10]. The second one is the value (19) of the mechanical energy depending on the rotation velocity and the eccentricity.

However, the kinematic description of the Keplerian motion is purely geometrical and mathematical, while Newton's description takes account of the physical parameters G and M . This causes a difference when considering the gravitational circular motion around a central body of mass M . Indeed, for Newton the velocity must then be:

$$v = \sqrt{\frac{GM}{r}} \quad (20)$$

Consequently, GM being postulated as a constant independent of r , the velocity can only decrease if the radius increases. The situation is different for the kinematics that states:

$$v = \sqrt{\frac{L v_R}{r}} = \sqrt{L \omega} \quad (21)$$

Therefore the velocity can be constant, whatever r , if $L \omega$ also is.

There is an experimental observation that could be concerned by this difference between the above velocities: the rotation of galaxies, when considering at a first approximation that the stars of the disk are in circular motion. Vera Rubin^[12] measured that the stars in the disk of spiral galaxies have approximately the same velocity, whatever their distance to the galactic bulb. This cannot be explained by Newton's velocity (20), but is consistent with the velocity (21) of the kinematics.

Noting that $L \omega$ has the dimension of an energy divided by a mass, the interpretation of the Keplerian kinematics of Vera Rubin's observation is that all the stars of the disk are in the same massless energy level $L \omega$. On the contrary, for the Newtonian interpretation there must exist a mass that we did not detect but speeds up the stars of the disk, the dark matter.

The whole connection between the kinematics's interpretation of the acceleration and Newton's one stands in the relation (8): $GM = L v_R$. On the left side of this equality is a constant physical factor, where G is postulated as the same universal constant, applicable to all masses, at all scales. On the contrary, on the right side is the kinematic factor $L v_R$, where L and v_R are defined for a given orbit but are both free to vary at any scale and for any phenomenon.

We may wonder which one nature is choosing to be the physical reality, the postulate of a genius of the 17th century, or the factual reality of the geometry. Whatever the choice, we must admit that the relation (8) is at least true at the solar system scale, because we experienced it so many times, even sending humans to the moon.

Actually, what suggests the kinematics is that G would not be a universal constant at all scales, but can vary regarding the scale and the phenomenon. Nonetheless it confirms that the mathematical structure of Newton's acceleration is indeed universal for all Keplerian motions.

Acknowledgment

Many thanks to Professor Emmanuel Trélat from Laboratoire Jacques-Louis Lions, CNRS, for his help on the kinematic part of this article.

References

- [1] W. R. Hamilton, "The hodograph, or a new method of expressing in symbolic language the Newtonian law of attraction" Proc. R. Ir. Acad. III, 344–353 ~1845–7
- [2] David Derbes, Reinventing the wheel: Hodographic solutions to the Kepler problems. American Journal of Physics 69, 481 (2001).
- [3] Orbit information derived from its hodograph, J. B. Eades, Tech. Rep. TM X-63301, NASA (1968)
- [4] H. Abelson, A. diSessa and L. Rudolph, Velocity space and the geometry of planetary orbits, Am. J. Phys. 43 , 579-589 (1975).
- [5] A. Gonzalez-Villanueva, H. N. Nunez-Yepe, and A. L. Salas-Brito, In velocity space the Kepler orbits are circular, Eur. J. Phys. 17 , 168-171 (1996).
- [6] T. A. Apostolatos, Hodograph: A useful geometrical tool for solving some difficult problems in dynamics, Am. J. Phys. 71 , 261-266 (2003).
- [7] José F Cariñena et al 2016, A new look at the Feynman 'hodograph' approach to the Kepler first law, Eur. J. Phys. 37 025004
- [8] E. I. Butikov, The velocity hodograph for an arbitrary Keplerian motion, Eur. J. Phys. 21 (2000) 1–10.
- [9] L. Landau & E. Lifchitz, Mechanics, Ed. Mir, Moscow, 1966, §15.
- [10] Cordani, Bruno (2003). The Kepler Problem. Birkhaeuser. p. 22. ISBN 3-7643-6902-7.
- [11] L. Landau & E. Lifchitz, Mechanics, Ed. Mir, Moscow, 1966, §14.
- [12] Rubin, V. C.; Ford Jr. W.K. & Thonnard, N., « Extended rotation curves of high-luminosity spiral galaxies. IV. Systematic dynamical properties, Sa -> Sc », The Astrophysical Journal, vol. 225, 1978, L107-L111