

Born Reciprocal (Non-inertial) Relativity, Phase Space Trajectories and Strings with variable Tension

Carlos Castro Perelman
Bahamas Advanced Science Institute and Conferences
Ocean Heights, Stella Maris, Long Island, the Bahamas
perelmanc@hotmail.com

August 2025

Abstract

Pursuing further our work on Born Reciprocal (Non-inertial) Relativity theory we find that the family of hyperbolic trajectories of particles living in a four-dim phase space of signature $(2, 2)$ automatically satisfy the maximal speed of light and the maximal proper force conditions. A direct link between the maximal proper force b and the minimum value of the initial positions $(x_o)_{min}$ of these hyperbolic paths is found. Furthermore, when the maximal proper force is set to be equal to the Planck-mass squared $b = m_P^2$ then $(x_o)_{min} = 2GM$ happens to coincide with the numerical value of the horizon radius of the Schwarzschild black hole in four spacetime dimensions. We proceed with a rigorous analysis of how open strings sweeping a two-dim world-sheet region inside a Rindler wedge in a $D = 1 + 1$ spacetime can be seen as a continuum of point masses that are being collectively accelerated along the Rindler hyperbolic trajectories. The upshot of this analysis reveals that these strings can naturally acquire a *variable* tension which is consistent with the dynamical tension model of strings and branes proposed by Guendelman in the past years [11]. We finalize by analyzing strings moving in phase space backgrounds within the context of non-inertial relativity theory.

Keywords : Born Reciprocal Relativity; Non-inertial Relativity; Phase Space; Strings; variable Tension string/brane models.

1 Introduction : Spacetime Trajectories

We shall begin by studying the behavior of the $D = 1 + 1$ spacetime trajectories of a massive particle of proper (rest) mass m subjected to a variable proper acceleration $g(\tau)$. The units are chosen to be $\hbar = c = 1$. After defining

$$u_i \equiv \frac{dx_i}{d\tau}, \quad \dot{u}_i \equiv \frac{du_i}{d\tau} = \frac{d^2x_i}{d\tau^2}, \quad i = 0, 1 \quad (1)$$

where the derivatives are taken with respect to the proper time τ , let us find solutions to the equations

$$(u_0)^2 - (u_1)^2 = 1, \quad (\dot{u}_0)^2 - (\dot{u}_1)^2 = -g^2(\tau), \quad (2)$$

corresponding, respectively, to the normalization condition on the timelike velocity of a massive particle, and subjected to a variable spacelike proper acceleration $g(\tau)$, in a $D = 1 + 1$ spacetime of signature $(+, -)$. Solving u_0 in terms of u_1 yields $u_0 = \sqrt{1 + (u_1)^2}$ in the first equation. Substituting this value into the second equation leads to the differential equation

$$\frac{(du_1/d\tau)^2}{1 + (u_1)^2} = g^2(\tau) \quad (3)$$

and whose solution is given by

$$u_1 = \frac{dx}{d\tau} = \sinh\left(\int g(\tau)d\tau\right) \Rightarrow x(\tau) = \int_0^\tau \sinh\left(\int_0^{\tau'} g(\tau'')d\tau''\right) d\tau' \quad (4)$$

The solution for $t(\tau)$ turns out to be

$$t(\tau) = \int_0^\tau \cosh\left(\int_0^{\tau'} g(\tau'')d\tau''\right) d\tau' \quad (5)$$

One can verify by a mere *substitution* that the solutions in eqs-(4,5) for $t(\tau), x(\tau)$ obey

$$\left(\frac{d^2t}{d\tau^2}\right)^2 - \left(\frac{d^2x}{d\tau^2}\right)^2 = -(g(\tau))^2 \quad (6)$$

for *all* functions $g(\tau)$.

In the special case that $g(\tau) = g_o$ constant, the integrals (4,5) lead to

$$t(\tau) = \frac{1}{g_o} \sinh(g_o\tau), \quad x(\tau) = \frac{1}{g_o} \cosh(g_o\tau) \quad (7)$$

and one recovers the hyperbolic trajectories $x^2 - t^2 = (g_o)^{-2}$ associated with the motion of a massive particle subjected to a uniform proper acceleration g_o .

In the next section we shall analyze the phase space trajectories after implementing the maximal proper force condition $|mg(\tau)| \leq b$. The proper force is bounded by the physical numerical constant b of units of $(mass)^2$, and is associated with the construction of the Born Reciprocal Relativity Theory (Non-inertial Relativity) in phase space involving x, t, p, E [3], [4], [5], [6], [10].

One can enforce the maximal proper force condition by writing

$$\frac{F(\tau)}{b} = \frac{mg(\tau)}{b} = \tanh[\xi(\tau)] \Rightarrow g(\tau) = \frac{b}{m} \tanh[\xi(\tau)] \quad (8)$$

where $\xi(\tau)$ is any arbitrary function of the proper time τ and is associated with a variable force-boost rapidity. Inserting $g(\tau)$ given by eq-(8) into eqs-(4,5) yields

$$t(\tau) = \int_0^\tau \cosh \left(\int_0^{\tau'} \frac{b}{m} \tanh[\xi(\tau'')] d\tau'' \right) d\tau' \quad (9)$$

$$x(\tau) = \int_0^\tau \sinh \left(\int_0^{\tau'} \frac{b}{m} \tanh[\xi(\tau'')] d\tau'' \right) d\tau' \quad (10)$$

One can verify that the velocity never exceeds the speed of light $c = 1$

$$v = \frac{dx}{dt} = \frac{(dx/d\tau)}{(dt/d\tau)} = \tanh \left(\int_0^\tau \frac{b}{m} \tanh[\xi(\tau'')] d\tau'' \right) \quad (11)$$

Because the hyperbolic tangent function is bounded between 1 and -1 , from eq-(11) one learns that $|v| = |(dx/dt)| \leq 1$ for all the expressions provided by $\xi(\tau')$.

We shall examine 3 cases :

Case **1** . When $\xi(\tau) = \xi_o$ is constant one arrives at the solutions described by eqs-(7) with $g_o = \frac{b}{m} \tanh(\xi_o)$.

Case **2** . When $\xi(\tau) = g_o\tau$ is linear in τ , and when $mg_o = b$ coincides with the maximal proper force, after some algebra and neglecting integration constants, one arrives at

$$t(\tau) = \frac{1}{2g_o} (\sinh(g_o\tau) + \arctan[\sinh(g_o\tau)]) \quad (12a)$$

$$x(\tau) = \frac{1}{2g_o} (\sinh(g_o\tau) - \arctan[\sinh(g_o\tau)]) \quad (12b)$$

Note the *different* functional behavior of the solutions (12,13) compared to the hyperbolic trajectories. The solutions provided in (12,13) implement the maximal force condition $|mg(\tau)| \leq b$ by construction.

Case **3** . When $\xi(\tau)$ is any arbitrary function one ends up with

$$t(\tau) = \int_0^\tau \cosh[\Omega(\tau')] d\tau', \quad x(\tau) = \int_0^\tau \sinh[\Omega(\tau')] d\tau',$$

$$\Omega(\tau') \equiv \int_0^{\tau'} \frac{b}{m} \tanh[\xi(\tau'')] d\tau'' \quad (13)$$

Having explored the trajectories in a $D = 1 + 1$ spacetime in the next section we shall extend our analysis to the trajectories of point masses in a four-dim phase space background of signature $(2, 2)$ associated with a $D = 1 + 1$ spacetime, and proceed with the study of the propagation of strings in phase space backgrounds.

2 Non-inertial Relativity, Phase Space Trajectories of Point Masses and Strings

The principle behind the concept of “Born reciprocal relativity theory”, or non-inertial relativity to be more precise¹, was advocated by [3], [4], [5] and it was based on the idea proposed long ago by [1] that coordinates and momenta should be unified on the same footing. Consequently, if there is a limiting speed (temporal derivative of the position coordinates) in Nature there should be a maximal force as well, since force is the temporal derivative of the momentum. Hence, a *maximal* speed limit (speed of light) must be accompanied with a *maximal* proper force (which is also compatible with a *maximal* and *minimal* length duality) [5].

The principle of maximal acceleration was advocated earlier on by [7]. A chapter on Reciprocity Theory and Born’s Quantum Metric Operator appeared early on in the book by [2]. The concept of Born reciprocity in order to provide a new point of view on string theory in which spacetime is a derived dynamical concept was advanced by [8].

The generalized velocity and force (acceleration) boosts (rotations) transformations of the *flat* 8D Phase space coordinates, where $X^i, t, E, P^i; i = 1, 2, 3$ are \mathbf{c} -valued (classical) variables which are *all* boosted (rotated) into each-other, were given by [3] based on the group $U(1, 3)$ and which is the Born version of the Lorentz group $SO(1, 3)$. The $U(1, 3) = SU(1, 3) \times U(1)$ group transformations leave invariant the symplectic 2-form $\Omega = -dt \wedge dE + \delta_{ij} dX^i \wedge dP^j; i, j = 1, 2, 3$ and also the following Born-Green line interval in the *flat* 8D phase-space

$$(d\omega)^2 = (dt)^2 - (dX)^2 - (dY)^2 - (dZ)^2 + \frac{1}{b^2} ((dE)^2 - (dP_x)^2 - (dP_y)^2 - (dP_z)^2), \quad (c = 1) \quad (14a)$$

The maximal proper force is set to be given by b . The symplectic group is relevant because $U(1, 3) = Sp(8, R) \cap O(2, 6)$; $U(3, 1) = Sp(8, R) \cap O(6, 2)$, and $U(2, 2) = Sp(8, R) \cap O(4, 4)$.

These transformations can be *simplified* drastically when the velocity and force (acceleration) boosts are both parallel to the x -direction and leave the transverse directions Y, Z, P_y, P_z intact. There is now a subgroup $U(1, 1) = SU(1, 1) \times U(1) \subset U(1, 3)$ which leaves invariant the following phase space line interval

$$(d\omega)^2 = (dt)^2 - (dX)^2 + \frac{(dE)^2 - (dP)^2}{b^2} = (d\tau)^2 \left(1 + \frac{(dE/d\tau)^2 - (dP/d\tau)^2}{b^2} \right) = (d\tau)^2 \left(1 - \frac{F^2}{F_{max}^2} \right), \quad F_{max} = b \quad (14b)$$

¹We thank one of the referees of a previous article for highlighting this fact in order to clarify the point that Born did not propose a reciprocal relativity theory

where one has factored out the non-vanishing proper time infinitesimal $(d\tau)^2 = dt^2 - dX^2 \neq 0$ in (14b). The numerical quantity F^2 is positive by definition. The proper force on a massive particle is given by $F = ma$, where a is the proper acceleration and m is the rest mass. We refrained from factoring out $(dt)^2$ in (14b) because it is not Lorentz invariant, whereas $(d\tau)^2$ is Lorentz invariant.

The $U(1, 1)$ group transformations involving the velocity and force boosts (along the X direction) acting on the phase-space coordinates X, t, P, E and which leave invariant the interval (14b) are given by [3], [4]

$$t' = t \cosh \xi + (\xi_v x + \frac{\xi_a P}{b}) \frac{\sinh \xi}{\xi} \quad (15a)$$

$$E' = E \cosh \xi + (b \xi_a X + \xi_v P) \frac{\sinh \xi}{\xi} \quad (15b)$$

$$X' = X \cosh \xi + (\xi_v t + \frac{\xi_a E}{b}) \frac{\sinh \xi}{\xi} \quad (15c)$$

$$P' = P \cosh \xi + (\xi_v E + b \xi_a t) \frac{\sinh \xi}{\xi} \quad (15d)$$

ξ_v is the velocity-boost rapidity parameter; ξ_a is the force (acceleration) boost rapidity parameter, and ξ is the net effective rapidity parameter of the primed-reference frame. The rapidity parameters ξ_a, ξ_v, ξ are defined, respectively, in terms of the spatial velocity $v = dx/dt$, and proper force $F = ma$, as follows

$$\tanh(\xi_v) = v; \quad \tanh(\xi_a) = \frac{F}{F_{max}}, \quad F_{max} = b, \quad \xi = \sqrt{(\xi_v)^2 + (\xi_a)^2} \quad (15e)$$

When $\xi_v \rightarrow \infty \Rightarrow v \rightarrow c = 1$. And $\xi_a \rightarrow \infty \Rightarrow F \rightarrow F_{max} = b$.

It is straight-forward to verify that the transformations of eqs-(15) leave invariant the phase space interval $(dt)^2 - (dX)^2 + ((dE)^2 - (dP)^2)/b^2$ but *do not* leave separately invariant the proper time interval $(d\tau)^2 = dt^2 - dX^2$, nor the interval in energy-momentum space $\frac{1}{b^2} [(dE)^2 - (dP)^2]$. Only the *combination*

$$(d\omega)^2 = (d\tau)^2 \left(1 - \frac{F^2}{F_{max}^2} \right) \quad (15f)$$

is truly left invariant under force (acceleration) boosts. They also leave invariant the symplectic 2-form (phase space areas) $\Omega = - dt \wedge dE + dX \wedge dP$.

It is very important to emphasize that there are *no* factors of $(1 + F^2/b^2)$ appearing in the above factorization process because in the superluminal case $(d\tau)^2 < 0$ (spacelike interval) one still has $m^2 a^2 < 0$ despite that $a^2 > 0$ (timelike proper acceleration), because $m^2 < 0$ due to the *imaginary* mass of tachyons. When $(d\tau)^2 > 0$ (timelike interval) $\Rightarrow m^2 a^2 < 0$ since $a^2 < 0$ (spacelike proper acceleration) and $m^2 > 0$. Hence we shall always have the factor $(1 - F^2/b^2)$ as expected with $m^2 a^2 = -F^2 < 0$. This is also compatible with the fact that if $(d\tau)^2 > 0$, then $(dE)^2 - (dP)^2 < 0$, and vice versa, if $(d\tau)^2 < 0$, then

$(dE)^2 - (dP)^2 > 0$. An exception occurs for a free particle (or at rest) giving $(dE)^2 - (dP)^2 = 0$ because $dE = dP = 0$.

The same results occur in higher dimensions. Given $E^2 - \vec{p} \cdot \vec{p} = E^2 - |\vec{p}|^2 = m^2 > 0$ when $(d\tau)^2 > 0$ gives upon differentiation

$\frac{dE}{d|\vec{p}|} = \frac{|\vec{p}|}{E} < 1$, and resulting in $(dE)^2 - d\vec{p} \cdot d\vec{p} = E^2 - |d\vec{p}|^2 < 0$. And vice versa, in the case of $E^2 - \vec{p} \cdot \vec{p} = E^2 - |\vec{p}|^2 = m^2 < 0$ (tachyon), when $(d\tau)^2 < 0$, gives upon differentiation $\frac{dE}{d|\vec{p}|} = \frac{|\vec{p}|}{E} > 1$, and resulting in $(dE)^2 - d\vec{p} \cdot d\vec{p} = E^2 - |d\vec{p}|^2 > 0$.

Consequently, the *negative* sign appearing inside the parenthesis in eqs (14b,15f) furnishes the analog of the Lorentz relativistic factor in special relativity and it involves the ratio of the square of two *proper* forces. The result (15) in the 4-dim phase space can be generalized to the $8D$ -dim phase space (and to higher dimensions)

In the case of hyperbolic trajectories associated with the motion of a massive particle subjected to a uniform proper acceleration g_o , the phase space infinitesimal interval (15f) is given by $d\omega = \sqrt{1 - \frac{m^2 g_o^2}{b^2}} d\tau$. Due to the $U(1,1)$ invariance of $d\omega$, under force boosts transformations one has the relation

$$d\omega = \sqrt{1 - \frac{(mg_o)^2}{b^2}} d\tau = \sqrt{1 - \frac{(m'g'_o)^2}{b^2}} d\tau' \quad (16)$$

which reveals that $F = mg_o \neq F' = m'g'_o$ and $\tau \neq \tau'$. Namely, the proper force mg_o and the proper time τ are *not* fully $U(1,1)$ invariant, they are ($SO(1,1)$) Lorentz invariant. The explicit transformations relating $m'g'_o$ with mg_o , and relating τ' with τ , in terms of the force-boost rapidity parameter ξ_a can be found in [6].

Therefore, ω is the truly $U(1,1)$ -invariant evolution parameter that must be used in order to describe the phase space trajectories of a particle instead of the proper time τ which is *not* $U(1,1)$ invariant. Consequently, instead of having the expressions associated to hyperbolic trajectories

$$t(\tau) = \frac{1}{g_o} \sinh(g_o\tau), \quad x(\tau) = \frac{1}{g_o} \cosh(g_o\tau) \quad (17)$$

$$p(\tau) = m \frac{dx}{d\tau} = m \sinh(g_o\tau), \quad E(\tau) = m \frac{dt}{d\tau} = m \cosh(g_o\tau) \quad (18)$$

one must have expressions defined in terms of ω as follows

$$Z_I(\omega) \equiv \left\{ t(\omega), x(\omega), \frac{1}{b} E(\omega), \frac{1}{b} p(\omega) \right\}, \quad I = 1, 2, 3, 4 \quad (19)$$

The analog of the timelike velocity condition $V_\mu V^\mu = 1$ in special relativity is now given in phase space by

$$\dot{Z}_I \dot{Z}^I = \left(\frac{dt}{d\omega}\right)^2 - \left(\frac{dx}{d\omega}\right)^2 + \frac{1}{b^2} \left(\frac{dE}{d\omega}\right)^2 - \frac{1}{b^2} \left(\frac{dp}{d\omega}\right)^2 = 1 \quad (20)$$

The extension to phase space of the spacelike condition on the proper acceleration in spacetime is

$$\ddot{Z}_I \ddot{Z}^I = \left(\frac{d^2 t}{d\omega^2}\right)^2 - \left(\frac{d^2 x}{d\omega^2}\right)^2 + \frac{1}{b^2} \left(\frac{d^2 E}{d\omega^2}\right)^2 - \frac{1}{b^2} \left(\frac{d^2 p}{d\omega^2}\right)^2 = -\mathcal{A}^2(\omega) \quad (21)$$

where $\mathcal{A}(\omega)$ is the phase space analog of the proper spacetime acceleration and must not be confused with $g(\tau)$. Given $\dot{Z}_I \dot{Z}^I = 1$, a differentiation yields $\ddot{Z}_I \dot{Z}^I = 0$, consequently the phase space analog of the proper acceleration is orthogonal to the phase space analog of velocity, hence if the velocity is timelike, the acceleration is spacelike, and vice versa.

A thorough study of the spacelike $(d\omega)^2 < 0$, null $(d\omega)^2 = 0$, and timelike $(d\omega)^2 > 0$ intervals in phase space, and their relation to the intervals $(d\tau)^2$ in spacetime, can be found in [6] where it was shown in many examples that there are *no* crossovers in the spacetime intervals when one performs a force boost transformation for any value of the force boost rapidity ξ_a rapidity parameter. Consequently, one has that if $(d\tau)^2 > 0 \Rightarrow (d\tau')^2 > 0$. And viceversa, if $(d\tau)^2 < 0 \Rightarrow (d\tau')^2 < 0$.

Before we discuss the energy and momentum it is very important to invoke the construction of the quadratic Casimir invariants of the Quaplectic group studied by Low [3], [4]. The Quaplectic group in four spacetime dimensions (eight phase space dimensions) is the semi-direct product of $U(1,3)$ with the translations in phase space and including the unit central element associated with the Weyl-Heisenberg algebra. The relevance of the quadratic Casimir is that it correctly defines the analog of proper mass in phase space \mathcal{M} . Therefore, upon defining

$$\mathcal{M} \frac{dt}{d\omega} = E, \quad \mathcal{M} \frac{dx}{d\omega} = p \quad (22)$$

where \mathcal{M} is the $U(1,1)$ -invariant proper mass in phase space, and which is not the same as m , the two eqs-(20,21) become a system of two simultaneous differential equations for the two functions $E(\omega), p(\omega)$ given by

$$\left(\frac{E}{\mathcal{M}}\right)^2 - \left(\frac{p}{\mathcal{M}}\right)^2 + \frac{1}{b^2} \left(\frac{dE}{d\omega}\right)^2 - \frac{1}{b^2} \left(\frac{dp}{d\omega}\right)^2 = 1 \quad (23)$$

$$\left(\frac{1}{\mathcal{M}}\right)^2 \left(\frac{dE}{d\omega}\right)^2 - \left(\frac{1}{\mathcal{M}}\right)^2 \left(\frac{dp}{d\omega}\right)^2 + \frac{1}{b^2} \left(\frac{d^2 E}{d\omega^2}\right)^2 - \frac{1}{b^2} \left(\frac{d^2 p}{d\omega^2}\right)^2 = -\mathcal{A}^2(\omega) \quad (24)$$

In the $b \rightarrow \infty$ limit, eq-(23) becomes $E^2 - p^2 = \mathcal{M}^2$, and one recovers the on-shell condition since $\mathcal{M} \rightarrow m$ in the $b \rightarrow \infty$ limit. Whereas eq-(24) becomes the spacelike proper force squared as it occurs in special relativity since $\omega \rightarrow \tau$ and $\mathcal{A}(\omega) \rightarrow g(\tau)$

$$\left(\frac{dE}{d\tau}\right)^2 - \left(\frac{dp}{d\tau}\right)^2 = -(mg(\tau))^2 \quad (25)$$

Let us find a one-parameter family of solutions to eqs-(23,24) parametrized by the parameter κ after setting $\mathcal{A} = \mathcal{A}(\kappa)$ to be a one-parameter family of

accelerations *independent* of the phase space evolution parameter ω . A careful inspection reveals that a one-parameter family of solutions to eqs-(23,24) is given by

$$t(\omega; \kappa) = \frac{\kappa}{\mathcal{A}(\kappa)} \sinh[\mathcal{A}(\kappa) \omega], \quad x(\omega; \kappa) = \frac{\kappa}{\mathcal{A}(\kappa)} \cosh[\mathcal{A}(\kappa) \omega] \quad (26)$$

$$E(\omega; \kappa) = \kappa \mathcal{M} \cosh[\mathcal{A}(\kappa) \omega], \quad p(\omega; \kappa) = \kappa \mathcal{M} \sinh[\mathcal{A}(\kappa) \omega] \quad (27)$$

where κ is a numerical parameter.

From eq-(26) one infers that as $\omega \rightarrow \infty$ the particle reaches the speed of light $\frac{dx}{dt} = \frac{(dx/d\omega)}{(dt/d\omega)} = \tanh[\mathcal{A}(\kappa)\omega] \rightarrow 1$ ($c = 1$).

Inserting the solutions given by eq-(27) into eqs-(23,24) lead to the relations

$$\frac{M^2 \mathcal{A}^2(\kappa)}{b^2} = 1 - \frac{1}{\kappa^2}, \quad \mathcal{M} \mathcal{A}(\kappa) = b \sqrt{1 - \frac{1}{\kappa^2}} \leq b, \quad \kappa \geq 1 \quad (28)$$

In order to avoid complex values for $\mathcal{M}\mathcal{A}(\kappa)$, one must choose $\kappa \geq 1$. A value of $\kappa = 1$ yields $\mathcal{M}\mathcal{A}(\kappa = 1) = 0$. The condition $\mathcal{M}\mathcal{A}(\kappa) \leq b$ in eq-(28) is also a sign of consistency such that the maximal upper bound of b is not exceeded. In the limit $\kappa \rightarrow \infty$ one has $\mathcal{M}\mathcal{A}(\kappa) \rightarrow b$ and the upper bound b is saturated.

The solutions (26,27) describe a one-parameter family of elliptic-hyperboloids in four dimensions defined by the algebraic equation

$$x^2 + \frac{p^2}{b^2} - t^2 - \frac{E^2}{b^2} = \frac{1}{\mathcal{A}(\kappa)^2} = \frac{\mathcal{M}^2}{b^2(1 - 1/\kappa^2)} \quad (29)$$

The above algebraic equation for the elliptic-hyperboloid is $U(1,1)$ -invariant.

Note the similarity of the family of solutions (26) with the family of Rindler hyperbolic trajectories if one replaces g, τ with \mathcal{A}, ω , respectively. The parameter $\kappa \geq 1$ is arbitrary and depicts the infinite family of trajectories. The similarity with the family of Rindler trajectories which span the Rindler coordinate-grid is more clear when κ is recast as $\kappa = \exp(\mathcal{A}_o \sigma)$, with \mathcal{A}_o constant, with $0 \leq \sigma \leq \infty$, so when $\sigma = 0 \Rightarrow \kappa = 1$, and $\sigma = \infty \Rightarrow \kappa = \infty$. The actual Rindler grid extends all the way to $\sigma = -\infty$ which corresponds to the origin $x = 0, t = 0$.

The initial positions of the trajectories in eq-(26) is described by the functions

$$x(\omega = 0; \kappa) = x_o(\kappa) = \frac{\kappa}{\mathcal{A}(\kappa)} = \frac{\kappa \mathcal{M}}{\mathcal{M}\mathcal{A}(\kappa)} = \frac{\kappa \mathcal{M}}{b \sqrt{1 - (1/\kappa^2)}} \quad (30)$$

$x_o(\kappa)$ is proportional to the throat-sizes of the elliptic-hyperboloids.

When $\kappa = 1$ and $\infty \Rightarrow x_o = \infty$. The minimum value of x_o is obtained by solving $\frac{dx_o}{d\kappa} = 0$. After some algebra one arrives at a quadratic equation for κ : $1 - \frac{2}{\kappa^2} = 0 \Rightarrow \kappa = \sqrt{2} > 1$. Inserting this value of κ into (30) yields

$$(x_o)_{min} = \frac{2\mathcal{M}}{b} = \frac{2G\mathcal{M}}{Gb} \quad (31)$$

If one sets the maximal proper force b to be equal to the Planck mass-squared $b = m_P^2$, in units of $\hbar = c = 1$, then $Gb = Gm_P^2 = L_P^2 m_P^2 = 1$, with L_P being the Planck scale in $4D$, such that the minimal initial position turns out to be $2G\mathcal{M}$ which coincides precisely with the numerical value of the horizon radius of the Schwarzschild black hole in four spacetime dimensions (which must not be confused with the four dimensions of the phase space we have been working with signature $(2, 2)$).

To be strictly rigorous, one should have worked directly with a 4-dim spacetime of signature $(1, 3)$, and an 8-dim phase space of signature $(2, 6)$ and restrict the solutions to a $2 + 2$ -dim subspace by choosing $y = z = p_y = p_z = 0$, which is compatible with the fact that $U(1, 1) \subset U(1, 3)$. In this way, one can appropriately assign $G = L_P^2 = (m_P)^{-2}$ in $\hbar = c = 1$ units and claim that $(x_o)_{min}$ matches the Schwarzschild horizon radius in $4D$.

To sum up, the salient features of this work are (i) : the hyperbolic family of trajectories in phase space automatically satisfy both the maximal speed of light $|dx/dt| \leq 1$, and the maximal proper force $\mathcal{M}\mathcal{A}(\kappa) \leq b$ condition. $\mathcal{M}\mathcal{A}(\kappa)$ is $U(1, 1)$ invariant, which in turn implies $SO(1, 1)$ -invariance; the converse is not true, as we saw with mg_o and τ *not* being invariant under force-boost transformations [5], [6]. (ii) one arrives at $(x_o)_{min} = \frac{2\mathcal{M}}{b}$ which establishes a direct link between the maximal proper force b and the minimum value of the initial position, a minimal distance from the origin given by eq-(31). Furthermore, (iii) : if the maximal proper force is $b = m_P^2$ then $(x_o)_{min} = 2G\mathcal{M}$ happens to coincide with the numerical value of the horizon radius of the Schwarzschild black hole in four spacetime dimensions.

The minimal value of the initial positions is a result which is in sharp contrast with what occurs in special relativity (inertial relativity). The initial positions of the family of Rindler hyperbolic trajectories in $D = 1 + 1$ can range between 0 and ∞ depending on the range of values of the uniform proper spacetime acceleration g_o given by $g_o = \infty$ and $g_o = 0$, respectively. The asymptotes to the Rindler hyperbolas are the null lines going through the origin $x_o = 0$ (zero minimal distance) corresponding to the massless photons trajectories which can be viewed as having an infinite proper acceleration $g_o = \infty$ and zero mass $m = 0$, and such that the double scaling limit is $mg_o = b$ [6], [5].

Hence we have shown how the behavior in phase space is very different and one has found a non-zero minimal value for the initial positions of the family of hyperbolic trajectories and which is determined in terms of the maximal proper force b as shown by eq-(31). It remains to explore the solutions of eqs-(23,24) for a *variable* proper phase space acceleration $\mathcal{A}(\omega; \kappa)$. However, to find solutions to eqs-(23,24) in this case when there is an explicit ω -dependence of \mathcal{A} is far more difficult than finding the solutions to eqs-(1,2) in spacetime and which were displayed by eqs-(4,5).

To finalize let us point out the connection [12] between open strings and the one-parameter family of Rindler hyperbolic trajectories in $D = 1 + 1$ given by

$$t(\eta, \sigma) = \frac{\exp(g_o \sigma)}{g_o} \sinh(g_o \eta), \quad x(\eta, \sigma) = \frac{\exp(g_o \sigma)}{g_o} \cosh(g_o \eta) \quad (32)$$

$$(d\tau)^2 = (dt)^2 - (dx)^2 = \exp(2g_o \sigma) (d\eta)^2 - (d\sigma)^2 \quad (33)$$

Each Rindler hyperbolic trajectory corresponds to setting $\sigma = \text{constant}$ in eq-(33) such that the proper time along each one of these trajectories becomes $(d\tau) = \exp(g_o \sigma) d\eta \Rightarrow \tau = \exp(g_o \sigma) \eta$, and one can write

$$g_o \eta = [\exp(-g_o \sigma) g_o] [\exp(g_o \sigma) \eta] = g(\sigma) \tau; \quad g(\sigma) = \exp(-g_o \sigma) g_o \quad (34)$$

such that the expressions in eq-(32) can be rewritten as

$$t(\tau, \sigma) = \frac{1}{g(\sigma)} \sinh[g(\sigma) \tau], \quad x(\tau, \sigma) = \frac{1}{g(\sigma)} \cosh[g(\sigma) \tau] \quad (35)$$

Note that the functional form of eq-(35) is *almost* similar to eq-(26) with the key difference that there are extra factors of $\kappa = \exp(\mathcal{A}_o \sigma)$ in eq-(26). The solutions (32) obey the string equations of motion²

$$\frac{\partial^2 t}{\partial \eta^2} - \frac{\partial^2 t}{\partial \sigma^2} = 0, \quad \frac{\partial^2 x}{\partial \eta^2} - \frac{\partial^2 x}{\partial \sigma^2} = 0 \quad (36)$$

whereas the solutions in eq-(35) do *not*

$$\frac{\partial^2 t}{\partial \tau^2} - \frac{\partial^2 t}{\partial \sigma^2} \neq 0, \quad \frac{\partial^2 x}{\partial \tau^2} - \frac{\partial^2 x}{\partial \sigma^2} \neq 0, \quad (37)$$

The variables η, σ appearing in eq-(36) are the appropriate temporal and spatial variables associated with the two-dimensional world-sheet, whereas τ in eq-(37) is the proper time affine parameter associated with the world-line of a point mass and whose spatial σ -dependent proper acceleration $g(\sigma)$ obeys the relation

$$\left(\frac{\partial^2 t}{\partial \tau^2}\right)^2 - \left(\frac{\partial^2 x}{\partial \sigma^2}\right)^2 = -(g(\sigma))^2, \quad g(\sigma) = g_o \exp(-g_o \sigma) \quad (38)$$

The physical interpretation of eq-(36) is the following : Imagine an infinite open string stretching from x_o all the way to $x = \infty$ and comprised of a continuum of point masses m where each one of these point masses is subjected to a σ -dependent proper force given by $F(\sigma) = mg(\sigma) = mg_o \exp(-g_o \sigma)$ endowing each one of these point masses with a hyperbolic trajectory such that the σ -dependent proper acceleration $g(\sigma)$ is $g_o \exp(-g_o \sigma)$. One could rewrite the expression for $F(\sigma)$ as $m(\sigma)g_o = [m_o \exp(-g_o \sigma)]g_o$, and in turn, interpret the proper force at each value of σ as that experienced by a σ -dependent mass distribution of points given by $m(\sigma)$ where each one of these mass points is experiencing the *same* proper acceleration g_o .

²The Dirichlet boundary conditions $t = x = 0$, and Neumann boundary conditions $\partial_\sigma t = \partial_\sigma x = 0$ are obeyed at $\sigma = -\infty$, but not at $\sigma = \infty$

Because the physical units of a proper force F are the same as a tension when $c = 1$: $[mc^2/L] = [Tc^2]$ (the tension T is just mass per unit length m/L) one could view this continuum arrangement of accelerated point masses as belonging collectively to an infinite open string with a variable tension $T(\sigma)$ which sweeps a two-dim world-sheet lying inside a region of the Rindler wedge [12]. The view of a string as a collection of point particles has been studied earlier on by [13].

Equating the tension T with the mass density $\frac{dm(\sigma)}{d\sigma}$ gives a variable but negative tension

$$T(\sigma) = \frac{dm(\sigma)}{d\sigma} = -m_o g_o \exp(-g_o \sigma) \quad (39)$$

Strings with a dynamical (variable) tension (both in space and time) $T(\eta, \sigma)$ have been extensively studied by Guendelman [11] over the years. In the modified measure formulation of strings/ branes, the tension appear as an additional dynamical degree of freedom. There are many important physical consequences of these variable tension models of strings and branes. Recently, Guendelman has reviewed how the model avoids the Swampland constraints making treatments for Dark energy and inflation more realistic and how strings with a different tension appear as Dark Matter to us. We refer to [11] and the many references therein for specific details. One should note that the dynamical tension description of Guendelman is based on the modified measure formulation which differs from the mechanism described in this work.

We have discussed above how to introduce a variable tension of the form $T(\sigma)$. The next question is how to introduce an additional temporal dependence. Because the string was comprised of a continuum of point masses that are being accelerated, if they emit gravitational radiation, there will be a loss of mass, and in this fashion, one could attain the sought after temporal dependence leading to a dynamical tension as well $T(\eta, \sigma)$.

To finalize this section, one could explore the maps

$$t(\eta, \sigma), \quad x(\eta, \sigma), \quad E(\eta, \sigma), \quad p(\eta, \sigma) \quad (40)$$

from the two-dim world-sheet of an open string, comprised of the coordinates (η, σ) , to the flat 2 + 2-dim phase background described by the coordinates t, x, E, p . Namely, by embedding a string in a flat phase space background and which is closely related to the metastring proposed by [8].

The simplest open string action one can construct is associated to a 2+2-dim phase space background with coordinates $Z^M = (X^\mu, b^{-1}P^\mu)$, equipped with a flat phase space metric $\eta_{MN} = \text{diag}(1, -1, 1, -1)$, and a constant symplectic form Ω_{MN} . The latter replaces the role of the antisymmetric tensor Kalb-Ramond field in ordinary string backgrounds. The action is

$$S = \frac{T}{2} \int d\eta d\sigma \sqrt{|g|} \left(g^{ab} (\partial_a Z^M) (\partial_b Z^N) \eta_{MN} + \frac{\epsilon^{ab}}{\sqrt{|g|}} (\partial_a Z^M) (\partial_b Z^N) \Omega_{MN} \right) \quad (41)$$

where g_{ab} is the world-sheet auxiliary metric, $a, b = 1, 2$, and $\epsilon^{ab} = -\epsilon^{ba}$.

The action (41) admits the following symmetries :

- (i) invariance under the $U(2, 2)$ transformations of the phase space background coordinates Z^M which include $SO(1, 1)$ Lorentz transformations and force-boost transformations eqs-(15a-15d).
- (ii) invariance under translations $X_\mu \rightarrow X_\mu + r_\mu$ and $P_\mu \rightarrow P_\mu + q_\mu$ with constant vectors r_μ, q_μ .
- (iii) invariance under world-sheet diffeomorphisms $\eta \rightarrow \eta'(\eta, \sigma)$; $\sigma \rightarrow \sigma'(\eta, \sigma)$.
- (iv) invariance under Weyl scalings $g_{ab} \rightarrow e^\phi g_{ab}$.

After fixing the world-sheet diffeomorphism and Weyl scale invariance the action (41) becomes

$$S = \frac{T}{2} \int d\eta d\sigma \left(\eta^{ab} (\partial_a Z^M) (\partial_b Z^N) \eta_{MN} + \epsilon^{ab} (\partial_a Z^M) (\partial_b Z^N) \Omega_{MN} \right) \quad (42)$$

The equations of motion of the variables Z^M (given a constant η_{MN}, Ω_{MN}) are

$$\frac{\partial^2 Z^M}{\partial \eta^2} - \frac{\partial^2 Z^M}{\partial \sigma^2} = 0 \quad (43)$$

Eqs-(43) admit the following solutions in terms of the modes expansion

$$X^\mu(\eta, \sigma) = x^\mu + l_s \pi_x^\mu \eta + i l_s \sum_{n \neq 0} \frac{1}{n} \alpha_n^\mu e^{-in\eta/l_s} \cos(n\sigma/l_s) \quad (44a)$$

$$P^\mu(\eta, \sigma) = p^\mu + \frac{b}{l_s} \pi_p^\mu \eta + i b l_s \sum_{n \neq 0} \frac{1}{n} \beta_n^\mu e^{-in\eta/l_s} \cos(n\sigma/l_s) \quad (44b)$$

The requirement that X^μ, P^μ are real functions require that the constants $x^\mu, p^\mu, \pi_x^\mu, \pi_p^\mu$ are real, while the positive and negative modes are complex conjugate to each other $\alpha_{-n}^\mu = (\alpha_n^\mu)^*$; $\beta_{-n}^\mu = (\beta_n^\mu)^*$. The constants x^μ, π_x^μ represent the string's center of mass position and its canonical conjugate (whose units are those of momentum), respectively. While p^μ, π_p^μ represent the string's center of mass momentum and its canonical conjugate (whose units are those of length), respectively. A new parameter, the string length scale l_s , has been introduced which is related to the string tension $T \sim \frac{1}{l_s^2}$, for details see [14].

The solutions (44) obey the open string Neumann boundary conditions at $\sigma = 0, \pi l_s$

$$\partial_\sigma Z^M(\sigma = 0) = 0, \quad \partial_\sigma Z^M(\sigma = \pi l_s) = 0 \quad (45)$$

The closed string solutions are given in terms of the sum of right and left movers [14] $X^\mu = X_R^\mu(\eta - \sigma) + X_L^\mu(\eta + \sigma), \dots$ obeying periodic boundary conditions $X^\mu(\eta, \sigma) = X^\mu(\eta, \sigma + \pi l_s), \dots$ as follows

$$X_R^\mu(\eta, \sigma) = \frac{1}{2} x^\mu + \frac{1}{2} l_s \pi_x^\mu (\eta - \sigma) + \frac{i}{2} l_s \sum_{n \neq 0} \frac{1}{n} \alpha_n^\mu e^{-2in(\eta - \sigma)/l_s} \quad (46a)$$

$$P_R^\mu(\eta, \sigma) = \frac{1}{2}p^\mu + \frac{b}{2l_s}\pi_p^\mu(\eta - \sigma) + \frac{i}{2}bl_s \sum_{n \neq 0} \frac{1}{n} \beta_n^\mu e^{-2in(\eta - \sigma)/l_s} \quad (46b)$$

$$X_L^\mu(\eta, \sigma) = \frac{1}{2}x^\mu + \frac{1}{2}l_s\pi_x^\mu(\eta + \sigma) + \frac{i}{2}l_s \sum_{n \neq 0} \frac{1}{n} \tilde{\alpha}_n^\mu e^{-2in(\eta + \sigma)/l_s} \quad (46c)$$

$$P_L^\mu(\eta, \sigma) = \frac{1}{2}p^\mu + \frac{b}{2l_s}\pi_p^\mu(\eta + \sigma) + \frac{i}{2}bl_s \sum_{n \neq 0} \frac{1}{n} \tilde{\beta}_n^\mu e^{-2in(\eta + \sigma)/l_s} \quad (46d)$$

where the right moving and left moving oscillator modes are $\alpha_n^\mu, \tilde{\alpha}_n^\mu$, and $\beta_n^\mu, \tilde{\beta}_n^\mu$.

The quantization of a Born-reciprocally-invariant string is beyond the scope of this work and is more subtle than the quantization of the ordinary string. For instance, the authors [17] studied the constraint quantization of the world-line system of a point particle that is invariant under the principle of Born reciprocity. The reciprocal transformations are not spin-conserving in general. The physical state space is vastly enriched as compared with the covariant approach, and contains towers of integer spin massive states (like the string theory spectrum), as well as unconventional massless representations, with continuous Euclidean momentum and arbitrary integer helicity.

Note that the solutions

$$t(\eta, \sigma) = \frac{\exp(g_o\sigma)}{g_o} \sinh(g_o\eta), \quad x(\eta, \sigma) = \frac{\exp(g_o\sigma)}{g_o} \cosh(g_o\eta) \quad (47a)$$

$$E(\eta, \sigma) = \mathcal{M} \exp(g_o\sigma) \cosh(g_o\eta), \quad p(\eta, \sigma) = \mathcal{M} \exp(g_o\sigma) \sinh(g_o\eta) \quad (47b)$$

with $T = \mathcal{M}^2$, obey the open string equations of motion (43) but do *not* obey the open string Neumann boundary conditions (45).

Despite that the boundary conditions are not obeyed by eqs-(47) one could still extract a variable tension function as follows. By setting $\kappa = \exp(\mathcal{A}_o\sigma)$ in the expression for the phase space analog of the proper force found in eq-(28), it yields

$$\mathcal{F}(\sigma) = \mathcal{M} \mathcal{A}(\sigma) = b \sqrt{1 - \exp(-2\mathcal{A}_o\sigma)}, \quad \kappa = \exp(\mathcal{A}_o\sigma) \quad (48)$$

and leads to a σ -dependent mass distribution $\mathcal{M}(\sigma)$ after equating

$$\mathcal{F}(\sigma) = \mathcal{M} \mathcal{A}(\sigma) = \mathcal{M}(\sigma) \mathcal{A}_o \Rightarrow \mathcal{M}(\sigma) = \frac{b}{\mathcal{A}_o} \sqrt{1 - \exp(-2\mathcal{A}_o\sigma)} \quad (49)$$

such that

$$T(\sigma) = \frac{d\mathcal{M}(\sigma)}{d\sigma} = b \frac{\exp(-2\mathcal{A}_o\sigma)}{\sqrt{1 - \exp(-2\mathcal{A}_o\sigma)}} \quad (50)$$

is a *positive* variable tension given in terms of the constant acceleration parameter \mathcal{A}_o and the σ variable.

Because the above tension *blows* up at $\sigma = 0$ one needs to introduce a cut-off σ_{min} that can be chosen to coincide with the value of $\kappa = \sqrt{2}$ associated with the minimal initial position $(x_o)_{min} = 2\mathcal{M}/b$ found in eq-(31). In doing so one arrives at $\sqrt{2} = \exp(\mathcal{A}_o\sigma_{min}) \Rightarrow \sigma_{min} = \ln(\sqrt{2})/\mathcal{A}_o$, and the tension at σ_{min} becomes $T(\sigma_{min}) = \frac{b}{\sqrt{2}} < b$. One could choose the cutoff value for σ to be that point $0 < \sigma_* < \sigma_{min}$ such that $T(\sigma_*) = b$ is precisely the maximal proper force. The tension at $\sigma = \infty$ is zero.

3 Concluding Remarks

In this work we found a family of hyperbolic trajectories of particles living in a four-dim phase space of signature (2, 2) obeying the maximal speed of light and the maximal proper force conditions. A direct link between the maximal proper force b and the minimum value of the initial positions $(x_o)_{min}$ of these hyperbolic paths is found. Furthermore, when the maximal proper force is set to be equal to the Planck-mass squared $b = m_P^2$ then $(x_o)_{min} = 2G\mathcal{M}$ happens to coincide with the numerical value of the horizon radius of the Schwarzschild black hole in four spacetime dimensions.

We proceeded with a rigorous analysis of how open strings sweeping a two-dim world-sheet region inside a Rindler wedge in a $D = 1 + 1$ spacetime can be seen as a continuum of point masses that are being collectively accelerated along the Rindler hyperbolic trajectories. The upshot of this analysis reveals that these strings can naturally acquire a *variable* tension which is consistent with the dynamical tension model of strings and branes proposed by Guendelman in the past years [11]. And finally, we constructed the action of a string moving in a flat target phase space background and found solutions to the equations of motion for the open and closed string. These results differ from the metastring [8] and the double field theories [9]. For these reasons, the quantization program of a string invariant under the principle of Born reciprocity deserves to be investigated further.

So far we have studied the *flat* Born geometry. A curved geometry of the phase space (cotangent space) requires the tools of Hamilton-Cartan geometry (Lagrange-Finsler geometry in the case of tangent space). The Born interval in an 8-dim curved phase space (cotangent space) is given by

$$(d\omega)^2 = g_{\mu\nu}(x, p) dx^\mu dx^\nu + h_{ab}(x, p) (dp^a + A_\mu^a(x, p) dx^\mu) (dp^b + A_\nu^b(x, p) dx^\nu) \quad (51)$$

$g_{\mu\nu}(x, p)$ is the horizontal base spacetime metric; $\mu, \nu = 0, 1, 2, 3$. $h_{ab}(x, p)$ is the vertical space (fiber) metric; $a, b = 0, 1, 2, 3$. $A_\mu^a(x, p)$ is the *nonlinear* connection. The flat space limit occurs when $g_{\mu\nu} = \eta_{\mu\nu}$; $h_{ab} = \frac{1}{b^2}\eta_{ab}$; $A_\mu^a = 0$. See [10] and the many references therein for more details.

A curved geometry of the phase space has many other applications like in resolving the problem of UV divergences in QFT. Recently, an intrinsic regularization via curved momentum space as a geometric solution to divergences in Quantum Field Theory has been proposed by [15]. The notion of curved momentum space, as it arises in the discussion of noncommutative spacetime coordinates or Doubly Special relativity theories, and the nontrivial connections with spectral geometry and Hopf algebras has been analyzed by [16].

To conclude, besides point masses it is warranted to explore, further, strings and branes in phase space backgrounds within the context of Born Reciprocal (Non-inertial) Relativity, and to explore the novel physical consequences of the classical and quantum theory.

Acknowledgments

We thank M. Bowers for assistance; to Eduardo Guendelman for many insightful discussions of his work on strings/branes with dynamical tension and to Igor Kanatchikov for providing reference [2].

References

- [1] M. Born, “Elementary Particles and the Principle of Reciprocity”. *Nature* **163** (1949), p. 207
- [2] E.Prugovečki, *Stochastic Quantum Mechanics and Quantum Spacetime*, Kluwer Academic Publishers, 1984, ISBN 978-9027716170. (Section 4.5 Reciprocity Theory and Born’s Quantum Metric Operator)
- [3] S.Low, “ Canonically relativistic quantum mechanics: Casimir field equations of the Quaplectic group”, arXiv : math-ph/0404077
- [4] S. Low, “ $U(3, 1)$ Transformations with Invariant Symplectic and Orthogonal Metrics” , *Il Nuovo Cimento* **B 108** (1993) 841.
S. Low, “Representations of the canonical group, (the semi-direct product of the unitary and Weyl-Heisenberg groups, acting as a dynamical group on noncommutative extended phase space”, *J. Phys. A : Math. Gen.*, **35** (2002) 5711.
- [5] C. Castro Perelman, “On Born Reciprocal Relativity Theory, The Relativistic Oscillator and the Fulling-Davies-Unruh effect”, to appear in the proceedings of the 9th European Congress of Mathematics (9ECM), the quadrennial Congress of the European Mathematical Society, Seville (Spain) 15 -19 July., 2024.
- [6] C. Castro Perelman, “Born Reciprocal Relativity Theory Redefines the notion of Mass : How Massless photons appear Massive in Accelerated Frames”, submitted to the IJGMMP, 2025.

- [7] E.R. Caianiello, “Is there a Maximal Acceleration”, *Il. Nuovo Cim.*, **32** (1981) 65.
H. Brandt, “Finslerian Fields in the Spacetime Tangent Bundle” *Chaos, Solitons and Fractals* **10** (2-3) (1999) 267.
H. Brandt, “Maximal proper acceleration relative to the vacuum” *Lett. Nuovo Cimento* **38** (1983) 522.
M. Toller, “The geometry of maximal acceleration” *Int. J. Theor. Phys* **29** (1990) 963.
- [8] L. Freidel, R. Leigh, D. Minic, “Born reciprocity in string theory and the nature of spacetime”, *Phys. Letters* **B 730**, (2014) 302.
L. Freidel, R. Leigh, D. Minic, “Modular Spacetime and Metastring Theory”, *Journal of Physics: Conf. Series* **804** (2017) 012032
L. Freidel, R. Leigh, D. Minic, “Quantum Gravity, Dynamical Phase Space and String Theory”, arXiv : 1405.3949
- [9] O. Hohm, C. Hull, and B. Zwiebach, “Generalized metric formulation of double field theory”. *Journal of High Energy Physics*. 2010 (8). arXiv:1006.4823.
B. Zwiebach, “Doubled Field Theory, T-Duality and Courant-Brackets”. *Strings and Fundamental Physics. Lecture Notes in Physics Vol. 851* (2012) pp. 265-291, Springer.
- [10] C. Castro Perelman, “Born Reciprocal Relativity Theory, Curved Phase Space, Finsler Geometry and the Cosmological Constant”, *Annals of Physics* **416**, (May 2020) 168143.
C. Castro Perelman, “On Born Deformed Reciprocal Complex Gravitational Theory and Noncommutative Gravity”, *Phys. Lett.* **B 668** (2008) 442-446.
C. Castro Perelman, “On Born Reciprocal Relativity, Algebraic Extensions of the Yang and Quaplectic Algebra, and Noncommutative Curved Phase Spaces”, *Universe* **9** (3) (2003) 144.
- [11] E. I. Guendelman, “Dynamical Tension Strings with Target Scale Symmetry producing DE, DM and why 4D?” arXiv : 2508.17333.
E.I. Guendelman, “Strings and branes with a modified measure”, *Class.Quant.Grav.* **17**, 3673-3680 (2000)
E.I. Guendelman, A.B. Kaganovich, E.Nissimov, S. Pacheva, “String and brane models with spontaneously or dynamically induced tension”, *Phys.Rev* **D66** :046003 (2002)
E.I. Guendelman, “Superstrings and superbranes with a modified measure”, *Phys.Rev.* **D 63** (2001) 046006

- [12] C. Castro Perelman, “On Maximal Acceleration, Strings with Dynamical Tension, and Rindler Worldsheets”. *Phys. Letts. B*. Published online <https://doi.org/10.1016/j.physletb.2022.137102>.
- [13] M. Pavsic, “A New Approach to the Classical and Quantum Dynamics of Branes”, *Int.J.Mod.Phys. A* **31** (2016) 1650115
M. Pavsic, “Branes and Quantized Fields”. *J.Phys.Conf.Ser.* **845** (2017) 1, 01201.
M. Pavsic, *Stumbling Blocks Against Unification* (World Scientific 2020). Chapter 6, “Brane space and branes as conglomerates of quantum fields”.
- [14] K. Becker, M. Becker, and J. Schwarz, *String Theory and M-Theory* (Cambridge University Press, 2007).
- [15] D. Ketels, “Intrinsic Regularization via Curved Momentum Space: A Geometric Solution to Divergences in Quantum Field Theory”, arXiv : 2502.14443.
- [16] S. A. Franchino-Viñas, S. Mignemi, J. J. Relancio, “The beauty of curved momentum space”, *Proc. Sci.* **436** (2023) 340.
- [17] J. Govaerts, P. D. Jarvis, S. O. Morgan, S. G. Low, “World-line Quantization of a Reciprocally Invariant System”. arXiv : 0706.3736
P.D. Jarvis and S.O Morgan, “Constraint quantization of a world-line system invariant under reciprocal relativity. II” arXiv : 0806.4794.