

# Quantum-elastic geometry: a unified theory of fields and constants of nature

Juan Moreno Borrillo

(Dated: September 15, 2025)

**Abstract.** *We present a unified field theory, termed Quantum-Elastic Geometry (QEG), wherein spacetime is modeled as a fundamental, physical substrate with quantum, elastic, and dissipative properties. The state of this medium is described by a single, symmetric rank-2 tensor field,  $\mathcal{G}_{\mu\nu}$ , whose dynamics are governed by a generally covariant action. Known physical interactions are shown to emerge as distinct, irreducible deformation modes of this unified field: gravity, electromagnetism, and a new field—denominated “thermo-entropic field”—that gives rise to irreversible thermodynamics. Furthermore, fundamental constants of nature are shown to be uniquely determined and interrelated by the substrate’s properties through a principle of self-consistency, and predicts a scale-dependent gravitational coupling that resolves key cosmological tensions, including the Hubble crisis. In summary, the developed framework offers a coherent origin for fields and constants, unifying them as emergent properties of a single, dynamic spacetime substrate.*

“*Entia non sunt multiplicanda praeter necessitatem*”  
— Ockham’s Razor

“*Padre, Señor del cielo y de la tierra, te doy gracias porque has ocultado todo esto a los sabios y entendidos y se lo has revelado a los que son como niños.*”  
— Matthew 11:25

## I. INTRODUCTION

The quest for a unified theoretical framework capable of describing all fundamental interactions from a common origin remains a central theme in contemporary physics [1]. Despite the tremendous success of the Standard Model of particle physics in unifying electromagnetic, weak and strong forces [2], and of General Relativity (GR) in geometrizing gravity [3, 4], a conceptual schism persists between the quantum field theories (QFTs) of the former and the geometric description of the latter [5]. Moreover, observational puzzles such as dark energy and dark matter [6–8], together with the inability to quantize gravity in a conventional QFT framework [9], underscore the need for a deeper, unifying structure.

Thermodynamic and emergent-gravity approaches have hinted at such a structure. Jacobson’s derivation of Einstein’s equations from local entropy balance [10], Verlinde’s entropic gravity proposal [11], and the striking analogies between black-hole thermodynamics and vacuum fluctuations [12] point toward an intimate link between entropy, quantum vacuum dynamics, and spacetime geometry, and suggest that disparate forces may be just different manifestations of a single underlying field.

Recent advances in the paradigm of emergent gravity suggest an even deeper connection, positing that spacetime geometry itself arises from the entanglement structure of an underlying quantum system [13–15]. In this view, entanglement acts as the “glue” of spacetime, a concept that resonates deeply with the QEG model of an “elastic substrate” whose properties are governed by the collective state of its constituent oscillators. The core idea of QEG—that the physics of curved spacetime emerges from an underlying medium—also finds powerful conceptual and experimental support in the field of analogue gravity [16]. In these systems, excitations like sound waves in a fluid can be described by an effective metric analogous to that of a curved spacetime, demonstrating that phenomena like event horizons are more general than gravity itself and can emerge from the collective behavior of a substrate [17].

In this work, we formalize this notion by proposing a theory of *Quantum-Elastic Geometry (QEG)*, which posits that spacetime itself is a dynamic, physical substrate with quantum, elastic, and dissipative properties. We elevate the standard QFT description of fields as collections of harmonic oscillators [18] from a mathematical tool to a physical model, postulating the vacuum as a *quantum oscillator lattice*. This lattice structure introduces a natural discreteness at the most fundamental scale, aligning our model with approaches such as Loop Quantum Gravity [19] and Causal Set Theory [20].

Postulating a physical “substrate” for spacetime inevitably raises the question of a preferred reference frame, or “aether,” and potential violations of Lorentz invariance, one of the most rigorously tested principles in physics. Modern effective field theories, such as the Einstein-Aether framework [21], provide a consistent language for exploring such concepts within a generally covariant context. The QEG model is consistent with the extremely stringent experimental limits on Lorentz violation [22], a condition met because the substrate is fully dynamic and covariant, preserving local Lorentz invariance for all propagating excitations.

The collective state of this oscillatory medium is described by a single, symmetric rank-2 tensor field,  $\mathcal{G}_{\mu\nu}(x)$ , which represents the local strain or deformation of the vacuum. All observed particles and force fields become emergent, large-scale manifestations of the different vibrational, shear, or torsional modes of this underlying lattice. Within this framework, fundamental constants lose their arbitrary character and become effective parameters describing the substrate’s material properties (e.g., stiffness, damping). This leads to a necessary *Geometro-dynamic Equivalence* between the dimensions of mass, length, and time, a concept whose formal justification can be found in modern gauge-theoretic approaches to gravity. Recent work by Partanen and Tulkki, for instance, has shown that founding gravity upon a  $U(1)^4$  gauge symmetry leads to a dimensionless coupling constant and a renormalizable quantum theory [23], providing a rigorous basis for the dimensional collapse explored here.

The aim of this paper is to construct the formal foundations of QEG. We begin by establishing a minimal set of physically-motivated axioms for the spacetime substrate. From these, we demonstrate how the established laws of General Relativity and Electromagnetism emerge as low-energy effective descriptions, along with the necessary emergence of a third fundamental interaction: a new field—which we coin “*thermo-entropic field*”—responsible for irreversible thermodynamics, which is required to account for the substrate’s remaining spatial degrees of freedom. The framework’s predictive power is then revealed by showing how the values of fundamental constants are neither postulated nor derived empirically but are uniquely constrained by the theory’s internal self-consistency. This leads to novel and falsifiable predictions, including the emergence of a scale-dependent gravitational coupling that provides a parameter-free resolution to key cosmological tensions. This specific prediction places the theory within the broader context of varying constant theories [24], which are subject to tight observational constraints from various astrophysical and cosmological probes, providing a direct path to falsify or validate the predictions of QEG [25]. Ultimately, this work presents a complete and coherent framework for the origin of fields and constants from the quantum-elastic properties of spacetime.

## Part I: The Foundations of Quantum-Elastic Geometry

### II. FROM FIRST PRINCIPLES TO SPACETIME EQUIVALENCE

The pursuit of a unified physical description has historically oscillated between two guiding intuitions: *simplicity*—which demands that the universe be described in terms of the fewest possible fundamental entities—and *structural inevitability*—whereby the mathematical form of natural laws emerges as the unique consistent framework rather than as an arbitrary convention—. The central ambition of this work is to bring these two intuitions together, building a unified substrate model governed by the *Principle of Parsimony* (*Ockham’s Razor*) and structural inevitability.

We proceed systematically from a minimal set of axioms: a single elastic substrate described by a deformation field  $\mathcal{G}_{\mu\nu}$ , whose dynamics are governed by a Lorentz-invariant Lagrangian with a single stiffness constant. This structure directly leads to the *Inertial Equivalence Principle*, unifying mass-like and current-like inertias. Furthermore, by showing that the 3D Laplacian dictates the inevitable and unique  $1/r$  static response of the medium, we establish the *Coupling Equivalence Principle*, which equates the dimensional nature of the gravitational and electrostatic coupling constants. Taken together, these two principles force an *Spacetime Equivalence Principle*: the collapse of the dimensions  $[M]$ ,  $[L]$ , and  $[T]$  into a single geometric entity. This unification is an unavoidable consequence of first principles, setting the stage for reinterpreting physical sources as endogenous excitations of spacetime itself.

### A. The Principle of a Unified Physical Substrate

Applying the *Principle of Parsimony* (*Ockham’s Razor*), we begin with a single foundational assumption:

*The universe, at its most fundamental level, consists of a single, unified physical entity or substrate.*

We will show that the identity of this substrate is not arbitrary. General Relativity has revealed that spacetime is a dynamic field,  $g_{\mu\nu}$ , while Quantum Field Theory describes the vacuum as a plenum of fluctuating fields. The most parsimonious conclusion is that *these are one and the same: spacetime itself*. Throughout this Section, we will show how the foundational assumption leads precisely to the conclusion expressed above on the nature of the unified physical entity or substrate.

### B. The Unified Lagrangian

A single substrate must be governed by a single, self-consistent dynamical principle. In modern physics, this is encoded in a Lorentz-invariant Lagrangian, from which all equations of motion are derived. The simplest such Lagrangian for a deformation field  $\mathcal{G}_{\mu\nu}$  consists of a kinetic term, representing the substrate’s inertial resistance to changes in deformation, and a potential term  $V(\mathcal{G})$ , representing its self-interaction:

$$\mathcal{L}_{\text{QEG}} = \frac{1}{2}\kappa(\partial^\alpha\mathcal{G}^{\mu\nu})(\partial_\alpha\mathcal{G}_{\mu\nu}) - V(\mathcal{G}), \quad (1)$$

where  $\kappa$  is a universal constant representing the intrinsic *stiffness* of the substrate. The principle of unification demands that this constant governs the inertial response to *all* possible modes of deformation.

It is crucial to note that the Lagrangian presented in 1 is a simplified formulation, valid in the weak field limit on a flat background spacetime (Minkowski). Its purpose at this stage is to establish the fundamental principles of substrate stiffness ( $\kappa$ ) and the inertia of its different modes of deformation in a clear and direct manner. A fully covariant and background-independent formulation, where the field  $\mathcal{G}_{\mu\nu}$  defines its own dynamic geometry, will be developed in Section VIII A. The present simplified model allows us to deduce the dimensional consequences of unification before addressing the complexity of the complete nonlinear dynamics.

#### *Minimal Scalar Field Model for Modal Excitations*

To provide a dynamical description of the modal excitations of the unified tensor  $\mathcal{G}_{\mu\nu}$ , we can construct a minimal model that captures their essential propagation features. This model serves as an effective description for a single projected mode, which we can denote as a scalar field  $\Phi(x)$ . The dynamics of this effective field are not arbitrary, but are a direct consequence of the general Lagrangian.

Recall that the kinetic part of the unified field Lagrangian is governed by a single, fundamental stiffness constant  $\kappa$ :

$$\mathcal{L}_{\text{kinetic}} = \frac{1}{2}\kappa(\partial^\alpha\mathcal{G}^{\mu\nu})(\partial_\alpha\mathcal{G}_{\mu\nu})$$

We can project these general dynamics onto a single scalar mode  $\Phi(x)$ , which will naturally contain terms for the kinetic and potential energy, analogous to a harmonic oscillator, and the two couplings (or "rigidities") usual for harmonic oscillatory systems,  $\kappa_1$  and  $\kappa_2$ , which are effective combinations of physical constants (e.g.,  $\varepsilon_0$ ,  $\mu_0$ ,  $G$ ,  $k_B$ , etc.) under different substitutions depending on the mode of the elastic-oscillatory manifestation of the common field. Then, one obtains the following *minimal* Lagrangian density:

$$\mathcal{L}(\Phi) = \frac{1}{2} \kappa_1 (\partial_t \Phi)^2 - \frac{1}{2} \kappa_2 (\nabla \Phi)^2 \quad (2)$$

Here,

- $\kappa_1$  controls the inertial" or kinetic response of the field mode,
- $\kappa_2$  represents the elastic/spatial rigidity of the field mode.

The corresponding *action*  $S$  is given by the integral

$$S[\Phi] = \int d^4x \mathcal{L}(\Phi, \partial\Phi; \kappa_1, \kappa_2) \quad (3)$$

Applying the principle of least action,  $\delta S = 0$ , yields the Euler–Lagrange equation:

$$\kappa_1 \frac{\partial^2 \Phi}{\partial t^2} - \kappa_2 \nabla^2 \Phi = 0. \quad (4)$$

This *single* partial differential equation governs the field  $\Phi$ . Depending on how we identify  $\kappa_i$  with physical constants and  $\Phi$  with different spacetime deformations (e.g., mass, charge, temperature), we recover the different field modes.

### C. Relativistic Compatibility and Spacetime Formalism

To ensure compatibility with special relativity, we introduce a four-dimensional spacetime coordinate:

$$X^\mu = \begin{pmatrix} t \\ x \\ y \\ z \end{pmatrix}, \quad \text{with metric } \eta^{\mu\nu} = \text{diag}(-1, 1, 1, 1). \quad (5)$$

We define the spacetime derivatives:

$$\partial_\mu = \frac{\partial}{\partial X^\mu} = \begin{pmatrix} \frac{\partial}{\partial t} \\ \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{pmatrix} \quad (6)$$

The standard Lorentz-invariant kinetic structure is encoded in the d'Alembertian operator:

$$\square\Phi \equiv \eta^{\mu\nu} \partial_\mu \partial_\nu \Phi = -\frac{\partial^2 \Phi}{\partial t^2} + \nabla^2 \Phi. \quad (7)$$

The equation of motion from our minimal field Lagrangian reads:

$$\kappa_1 \frac{\partial^2 \Phi}{\partial t^2} - \kappa_2 \nabla^2 \Phi = 0. \quad (8)$$

To write this in a Lorentz-invariant form proportional to  $\square\Phi = 0$ , we compare:

$$\kappa_1 \frac{\partial^2 \Phi}{\partial t^2} = \kappa_2 \nabla^2 \Phi \quad \implies \quad \frac{\partial^2 \Phi}{\partial t^2} = \frac{\kappa_2}{\kappa_1} \nabla^2 \Phi. \quad (9)$$

Rewriting the d'Alembertian as:

$$\square\Phi = -\frac{\partial^2 \Phi}{\partial t^2} + \nabla^2 \Phi = 0,$$

we see that Lorentz invariance requires:

$$\frac{\kappa_2}{\kappa_1} = 1 \quad \implies \quad \kappa_1 = \kappa_2. \quad (10)$$

Therefore, the field theory is manifestly Lorentz-invariant if and only if the rigidity constants match:  $\kappa_1 = \kappa_2$ . This is a profound physical statement: it reflects that in a Lorentz-covariant theory built upon a single field, the inertial and elastic responses to variations must originate from the same isotropic, fundamental stiffness  $\kappa$  of the vacuum substrate. This ensures the action transforms as a scalar and the field equation becomes the standard wave equation:

$$\square\Phi = 0. \quad (11)$$

This scalar field model, though minimal, shows how each modal projection of the unified field tensor  $\mathcal{G}_{\mu\nu}$  can be described in terms of a wave-like field obeying Lorentz-invariant dynamics. This prepares the ground for more detailed relativistic formulation in terms of Klein–Gordon dynamics and projection structures consistent with general relativity.

### Generalization to the Full Tensor Field

The above result –the necessary equality of temporal (inertial) and spatial (elastic) stiffness coefficients– is not limited to the pedagogical scalar model. It is a fundamental requirement for any relativistic field theory derived from a Lagrangian of the form of Eq. (1). For the full tensor field  $\mathcal{G}_{\mu\nu}$ , the kinetic term  $(\partial^\alpha \mathcal{G}^{\mu\nu})(\partial_\alpha \mathcal{G}_{\mu\nu})$  contains both time derivatives ( $\alpha = 0$ ) and spatial derivatives ( $\alpha = 1, 2, 3$ ). A Lorentz transformation mixes these temporal and spatial components. For the action to remain a Lorentz scalar, the universal coefficient  $\kappa$  multiplying this term must treat both types of derivatives equally. Any fundamental distinction between the substrate's response to temporal changes and spatial changes would break Lorentz invariance at the level of the action itself. Thus, the conclusion  $\kappa_1 = \kappa_2$  from the scalar model is a specific instance of a general principle dictated by the covariant structure of the unified Lagrangian.

### D. The Inertial Equivalence Principle

As we have seen in the previous subsection, the requirement of a unified Lagrangian has a profound and unavoidable consequence deeply rooted in Lorentz covariance: the single stiffness constant  $\kappa$  must consistently describe the inertia of every deformation mode. Analyzing the kinetic term reveals that  $\kappa$  plays a dual role:

- For *compressive* (scalar) modes, associated with mass-like excitations,  $\kappa$  acts analogously to mass in mechanical kinetic energy  $\frac{1}{2} m \dot{x}^2$ . Thus  $[\kappa] \equiv [\text{Mass}]$ .

- For *torsional* (vector) modes, associated with currents,  $\kappa$  acts as a coefficient of rotational inertia, mathematically analogous to inductance in electrodynamics. Thus  $[\kappa] \equiv [\text{Inductance}]$ .

Since Lorentz covariance and the principle of a single unified field demand that a single constant  $\kappa$  governs the inertial properties of all components of  $\mathcal{G}_{\mu\nu}$ , it follows that the dimensional character of mass and inductance must be identical:

$$\boxed{[\text{Mass}] \equiv [\kappa] \equiv [\text{Inductance}]}. \quad (12)$$

It is crucial to emphasize that *this is not merely a physical analogy but a structural requirement of the unified Lagrangian*. The single kinetic term,  $\frac{1}{2}\kappa(\partial^\alpha\mathcal{G}^{\mu\nu})(\partial_\alpha\mathcal{G}_{\mu\nu})$ , governs the inertia of *all* possible deformation modes of the substrate. Because this one mathematical object must simultaneously describe the resistance to compressive changes (a mass-like inertia) and the resistance to torsional changes (an inductance-like inertia), the dimensional characters of Mass and Inductance are forced to be identical. *They are not just analogous; they are two different phenomenological manifestations of the single, unified stiffness parameter  $\kappa$ .*

#### Dimensional Consequences of the Inertial Equivalence Principle

Taking the SI units of inductance  $L$  as

$$[L] = [ML^2I^{-2}T^{-2}],$$

the Inertial Equivalence Principle leads directly to

$$[M] \equiv [ML^2I^{-2}T^{-2}].$$

For dimensional consistency, this equivalence requires that the combination  $[L^2I^{-2}T^{-2}]$  must be dimensionless. Solving for the dimension of current  $[I]$  yields

$$[I^2] \equiv [L^2T^{-2}] \implies \boxed{[I] \equiv [LT^{-1}]}. \quad (13)$$

As a sanity check, in the same way that mass  $M$  in a mechanical oscillator is analogous to inductance  $L$  in an RLC circuit, the resistance  $R$  in an RLC circuit is analogous to the damping coefficient  $b$  in a mechanical oscillator. Establishing the dimensional equivalence between them, we find that

$$[MT^{-1}] \equiv [ML^2T^{-3}I^{-2}],$$

which again implies that  $[L^2I^{-2}T^{-2}]$  is dimensionless, consistent with the result (13).

Subsequently, from the Inertial Equivalence Principle, it follows that electric current acquires the dimensions of velocity. Expressed abstractly,

$$[I] \equiv [LT^{-1}], \quad (14)$$

so that current is no longer an independent unit (as in SI), but a kinematical manifestation of deformation flow. In physical terms, current is the propagation velocity of a torsional wave of the substrate, while mass is the inertial coefficient of a compressive deformation. Both are modes of the same elastic field.

## E. The Geometro-Elastic Principle and the Inevitability of the Laplacian Operator

The central thesis of QEG is that the vacuum is a continuous elastic medium. To describe the static configuration of its deformation field,  $\mathcal{G}_{\mu\nu}$ , we must identify the mathematical operator that governs equilibrium in such a substrate. This operator can be uniquely and rigorously determined by the most fundamental principles of symmetry and simplicity.

#### Justification from Symmetry: The Mandate of Isotropy

The most foundational property we can assume about the vacuum substrate is that it is *isotropic*: it has no preferred direction in space. Any fundamental law governing it must respect this symmetry. For a continuous elastic medium in static equilibrium, the operator describing its response must be invariant under rotations and translations.

The Laplacian ( $\nabla^2 = \sum_i \partial_i^2$ ) is the *unique* second-order linear differential operator that possesses this property. Any other operator of the same order, such as a weighted sum like  $(\partial_x^2 + 2\partial_y^2 + \dots)$ , would explicitly break rotational invariance, arbitrarily selecting certain axes as special. This would imply that the fabric of spacetime is intrinsically anisotropic, a claim that would require extraordinary justification and contradicts large-scale observation. Therefore, the *principle of isotropy* alone is sufficient to select the Laplacian as the necessary mathematical structure for describing the substrate's static response.

#### Justification from Simplicity: Ockham's Razor and Locality

The principle of parsimony (Ockham's Razor) demands that we use the simplest possible mathematical structure that adequately describes the physics. The Laplacian operator describes the simplest form of *local elastic tension*: it relates the state of the field at a point to the average state in its immediate infinitesimal vicinity.

One could postulate more complex, higher-order operators like the biharmonic operator ( $\nabla^4$ ). However, this would imply that the fundamental interactions of the substrate are non-local, depending not just on the immediate neighborhood but on its neighbors' neighbors. Without any physical or empirical necessity for such complexity at the most fundamental level, Ockham's Razor compels the choice of the lowest-order, non-trivial operator capable of describing elastic deformation: the Laplacian.

#### Mathematical Consequence: The Universal 1/r Potential

Once the principles of symmetry and simplicity force the selection of the Laplacian, the geometric form of all long-range static interactions is no longer a choice, but a *mathematical inevitability*. The static equilibrium equation for the deformation field  $\Phi$  in the presence of a point-like source at the origin (a Dirac delta function,  $\delta(\vec{r})$ ) must take the form of Poisson's equation:

$$\nabla^2\Phi(\vec{r}) = -J \cdot \delta(\vec{r}) \quad (15)$$

The unique solution to this equation in three-dimensional space is its Green's function, which has the universal form:

$$\nabla^2 \Phi(\mathbf{r}) = -\delta^{(3)}(\mathbf{r}) \implies \Phi(\mathbf{r}) = \frac{1}{4\pi r}. \quad (16)$$

This establishes the  $1/r$  geometric modulus as the *universal mathematical structure for all long-range static interactions* of an isotropic, elastic, 3D substrate. Consequently, both Newton's law of gravitation and Coulomb's law must appear as macroscopic limits of this same underlying Laplacian response. This forces a crucial conclusion: *if the substrate is truly unified, the constants that couple sources to the field in these laws must share the same dimensional essence*, a requirement that will be formalized as the Coupling Equivalence Principle in the next subsection.

#### Note on Short-Range Interactions

It is important to emphasize that the Laplacian operator and the resulting  $1/r$  potential are characteristics of the theory's limit for massless, static modes. This framework thus recovers the universal structure of long-range forces such as gravity and electromagnetism. Short-range interactions, such as the strong and weak nuclear forces, would be described within this framework as excitations of massive or non-linear modes of the field  $\mathcal{G}_{\mu\nu}$ . For these modes, the static field operator would take a different form (e.g., the Yukawa operator,  $\nabla^2 - m^2$ ), giving rise to short-range potentials (such as  $e^{-mr}/r$ ) and breaking the universality of the  $1/r$  behavior.

## F. The Coupling Equivalence Principle

We can extract another fundamental principle from the universal dynamics imposed by the unified Lagrangian and Lorentz covariance. Variation of the action leads to a field equation of the form:

$$\kappa \square \mathcal{G}_{\mu\nu} - \frac{\partial V}{\partial \mathcal{G}^{\mu\nu}} = J_{\mu\nu},$$

where the d'Alembertian operator,  $\square$ , governs the dynamics of all massless excitations of the substrate. This universal, hyperbolic operator is the unique source of the unified response.

The familiar empirical laws for static fields arise as the low-energy limit ( $\partial_t \rightarrow 0$ ) of these fundamental wave dynamics, where the d'Alembertian reduces to the Laplacian ( $\square \rightarrow \nabla^2$ ):

- For **Gravity**:  $\nabla^2 \Phi_g = 4\pi G \rho_m$
- For **Electrostatics**:  $\nabla^2 V = -(4\pi K_e) \rho_e$

Since both of these static laws are different projections of the *same* underlying dynamic operator ( $\square$ ), the constants that scale their respective sources— $G$  and  $K_e$ —cannot be dimensionally distinct. In a unified framework where mass density ( $\rho_m$ ) and charge density ( $\rho_e$ ) are manifestations of the same source tensor  $J_{\mu\nu}$ , it is a mathematical necessity that their couplings share the same dimensional character. To postulate otherwise would violate the covariant nature of the theory.

This leads us to state the *Coupling Equivalence*

*Principle* as a direct consequence of covariant dynamics:

$$\boxed{[G] \equiv [K_e]}. \quad (17)$$

This principle asserts that gravitational and electromagnetic couplings share the same dimensional essence, being merely modal projections of a single underlying rigidity of spacetime.

#### Dimensional Consequences of the Coupling Equivalence Principle

Let us now derive the dimensional consequences of this principle combined with our previous findings. The conventional dimensions for  $G$  and  $K_e$  are

$$[G] = [M^{-1}L^3T^{-2}], \quad [K_e] = [ML^3T^{-4}I^{-2}].$$

Setting  $[G] \equiv [K_e]$ , we obtain

$$[M^{-1}L^3T^{-2}] \equiv [ML^3T^{-4}I^{-2}].$$

Rearranging for  $[M]$ , and recalling that we have established the Inertial Equivalence Principle,  $[I] \equiv [LT^{-1}]$  (13), we substitute  $I^{-2}$  as

$$[M^{-1}L^3T^{-2}] \equiv [ML^3T^{-4}(L^{-2}T^2)].$$

Simplifying, we find

$$[M^{-1}L^3T^{-2}] \equiv [MLT^{-2}] \rightarrow$$

$$[M^{-1}L^3] \equiv [ML] \rightarrow$$

$$[L^2] \equiv [M^2] \rightarrow$$

$$\boxed{[M] \equiv [L]}. \quad (18)$$

This outcome, contingent on the Inertial Equivalence Principle (12) and the Coupling Equivalence Principle (IF), signifies that mass and length share the same fundamental dimension within this theoretical structure.

Moreover, substituting  $[M]$  and  $[I]$  into the previous equivalence between resistance  $R$  and damping coefficient  $b$  based on the Inertial Equivalence Principle,  $[MT^{-1}] \equiv [ML^2T^{-3}I^{-2}]$ , we find that  $[T^{-4}L^4]$  becomes dimensionless, which in turn implies the fundamental equivalence

$$\boxed{[M] \equiv [L] \equiv [T]}. \quad (19)$$

This result does not arise from a choice of units, but from geometric and dynamical consistency: the Laplacian enforces the  $1/r$  law for static responses, Lorentz invariance equates time and space, and the unified Lagrangian requires a single stiffness constant  $\kappa$  for all inertial modes. The collapse of mass, length, and time into a single dimensional entity is therefore an inevitable structural feature of a unified substrate.

## G. The Endogenous Nature of Sources

Finally, if mass, charge, and current share the same dimensional character as length and time, they cannot be external properties imposed on the substrate. Rather, they are *endogenous excitations*: localized, stable topological configurations of the deformation field  $\mathcal{G}_{\mu\nu}$ . Mass-energy and charge are not causes of deformation, but particular *modes of deformation* themselves. Thus particles and fields, matter and geometry, are unified as different aspects of the same physical substrate: *spacetime itself*.

### III. MODAL DECOMPOSITION OF THE UNIFIED FIELD

The deformation field  $\mathcal{G}_{\mu\nu}$  encodes the state of the unified substrate. To connect this framework with known physical interactions, it is natural to decompose  $\mathcal{G}_{\mu\nu}$  into modes according to their tensorial character and dynamical role. The decomposition mirrors the standard separation of elastic media into compressive and shear responses, but is here generalized to a covariant, field-theoretic setting.

#### A. Scalar, Vector, and Tensorial Sectors

We classify the components of  $\mathcal{G}_{\mu\nu}$  into three categories:

- **Scalar (compressive) mode:**  $\mathcal{G}_{00}$ , associated with localized compressions of the substrate. This sector corresponds to inertial mass and gravitation, as compressive deformations propagate radially and source the familiar Newtonian potential.
- **Vector (torsional) modes:**  $\mathcal{G}_{0i}$ , associated with shear-like deformations oriented along spatial directions. These modes are naturally linked to currents and electromagnetic phenomena, as they propagate transversely and encode the transport of torsional strain.
- **Tensorial modes:** The symmetric tensor  $\mathcal{G}_{ij}$  can be uniquely decomposed into its trace (an isotropic scalar mode) and its traceless part (an anisotropic shear mode). The traceless part is naturally identified with gravitational waves and anisotropic stresses. The trace,  $\text{Tr}(\mathcal{G}_{ij})$ , transforms as a scalar under spatial rotations and thus must couple to isotropic sources like pressure. As this components  $\mathcal{G}_{ij}$  represent the remaining degrees of freedom of the unified field, for the QEG framework to be complete, these modes must also correspond to a physical interaction. As we will demonstrate rigorously in Part IV, the dynamics of this scalar mode, when subjected to the substrate's intrinsic dissipation, inevitably lead to the laws of irreversible thermodynamics. Therefore, we identify this sector with a fundamental **thermo-entropic field**, not by analogy, but as a necessary consequence of the theory's completeness and internal consistency.

We now present the explicit projectors for the spatial tensor  $\mathcal{G}_{ij}$  due to its rich internal structure, which hosts multiple physical modes. The projection of the modes  $\mathcal{G}_{00}$  and  $\mathcal{G}_{0i}$  is trivial in comparison, as it reduces to the direct selection of those components, which already transform as a scalar and a vector under spatial rotations, respectively.

*Explicit projectors on the spatial tensor  $\mathcal{G}_{ij}$*

In a homogeneous/isotropic background (or in Fourier space on  $\mathbb{R}^3$ ), define  $k_i$  and the transverse projector  $P_{ij} = \delta_{ij} - \hat{k}_i \hat{k}_j$ ,  $\hat{k}_i = k_i/k$ . The symmetric

tensor  $\mathcal{G}_{ij}$  decomposes as

$$\mathcal{G}_{ij} = \underbrace{\left( \psi \delta_{ij} + (\partial_i \partial_j - \frac{1}{3} \delta_{ij} \nabla^2) E \right)}_{\text{scalar}} + \underbrace{\partial_{(i} F_{j)}}_{\text{vector, } \partial_i F_i = 0} + \underbrace{h_{ij}^{\text{TT}}}_{\text{tensor, } \partial_i h_{ij}^{\text{TT}} = 0, h_{ii}^{\text{TT}} = 0}. \quad (20)$$

Corresponding projectors in Fourier space read

$$\begin{aligned} (P^{\text{T}})_{ij,kl} &= \frac{1}{2} (P_{ik} P_{jl} + P_{il} P_{jk}) - \frac{1}{3} P_{ij} P_{kl}, \\ (P^{\text{V}})_{ij,kl} &= \frac{1}{2} (\hat{k}_i P_{jl} \hat{k}_k + \hat{k}_j P_{il} \hat{k}_k + \dots), \\ (P^{\text{S}})_{ij,kl} &= \frac{1}{3} P_{ij} P_{kl} + \hat{k}_i \hat{k}_j \hat{k}_k \hat{k}_l \end{aligned} \quad (21)$$

so that  $P^{\text{T}} + P^{\text{V}} + P^{\text{S}} = \mathbb{I}$  and  $P^A P^B = \delta^{AB} P^A$ .

*Justification from Symmetry and Lorentz Covariance*

The assignments of the tensor components of the deformation tensor  $\mathcal{G}_{\mu\nu}$  to the physical phenomena that we have made above can be justified as inevitable and physically necessary choices by appealing to the principles of symmetry and Lorentz covariance. The mapping is dictated by the transformation properties of sources under rotations and Lorentz boosts. Each physical source (scalar mass-energy, vector current, tensor stress/flux) must couple to the component of  $\mathcal{G}_{\mu\nu}$  with matching transformation properties. *Any other assignment would violate covariance.* Here's the breakdown:

*$G_{00}$  and Scalar Sources (Mass-Energy)*

- **Physical Source:** The primary source of Newtonian gravity is **mass**, a scalar quantity (it does not change under spatial rotations). In relativity, this is the energy density component,  $T_{00}$ , which also behaves as a scalar under spatial rotations.
- **Tensor Component:** The  $G_{00}$  component of the tensor is the only component that transforms as a **scalar** under rotations.
- **Conclusion:** By the principle of covariance, a scalar source must couple to a scalar mode of the field. The identification of mass-energy with  $G_{00}$  is the simplest and most direct way to achieve this. Associating a scalar source with a vector ( $G_{0i}$ ) or tensor ( $G_{ij}$ ) component would introduce an arbitrary direction into the physics, which is not observed.

*$G_{0i}$  and Vector Sources (Currents)*

- **Physical Source:** The source of magnetism is **electric current**,  $J_i$ , which is a 3-vector. It has a direction and magnitude that change predictably under rotation.
- **Tensor Component:** The three  $G_{0i}$  components are the only parts of the  $G_{\mu\nu}$  tensor that transform together as a **3-vector** under rotations.
- **Conclusion:** A vector source must couple to a vector-like part of the unified field. The identification of currents with the  $G_{0i}$  modes is therefore almost inevitable. This ensures that the laws of physics (like Ampère's Law) maintain their vector form in all rotated reference frames.

### *G<sub>ij</sub> and Tensor Phenomena (Stress and Gravitational Waves)*

- **Physical Phenomena:** Phenomena like pressure, shear stress ( $T_{ij}$ ), and the propagation of gravitational waves are inherently **tensorial**. They describe how space itself is stretched and sheared in different directions.
- **Tensor Component:** The  $G_{ij}$  components form a  $3 \times 3$  symmetric tensor that directly represents the spatial strain (stretching and shearing) of the physical substrate.
- **Conclusion:** It is physically necessary to map tensor phenomena onto the tensor components of the unified field. The  $G_{ij}$  components are the only ones that have the correct mathematical structure to describe the complex, directional stresses and strains within the spacetime fabric.

In summary, the assignment is a direct mapping of physical symmetries onto the mathematical symmetries of the tensor  $G_{\mu\nu}$ . The “shape” of the source or phenomenon (scalar, vector, or tensor) must match the “shape” of the deformation mode it couples to. This assignment is essentially unique under Lorentz invariance: the shape of the source dictates the shape of the field component it excites. This makes the chosen identification not just a convenient choice, but the only one that is fully consistent with the observed Lorentz symmetry of our universe.

This way, we have shown how gravitation, electromagnetism, and other tensor spatial deformations are interpreted as distinct *modal projections* of the same underlying substrate field.

### B. Unified Source Structure

In conventional field theory [26, 27], each interaction couples to a distinct current: the stress-energy tensor  $T^{\mu\nu}$  for gravitation, the electromagnetic four-current  $J^\mu$  for electrodynamics, and thermodynamic fluxes for entropy. In the present framework, all these appear as different components of a single unified source tensor  $J_{\mu\nu}$ , which couples universally to  $\mathcal{G}_{\mu\nu}$  via the action principle. Explicitly:

$$\begin{aligned} J_{00} &\longleftrightarrow T_{00} \quad (\text{mass-energy density component}) \\ J_{0i} &\longleftrightarrow J_{\text{em}}^i \quad (\text{spatial components of 4-current}) \\ \text{Tr}(J_{ij}) &\longleftrightarrow p \quad (\text{pressure, thermodynamic state}) \end{aligned}$$

In addition, the traceless spatial part acts as the natural source for shear modes:

$$\begin{aligned} (J_{ij} - \frac{1}{3}\delta_{ij} \text{Tr}(J)) &\longleftrightarrow \\ \pi_{ij} \quad (\text{anisotropic stress / shear source}) & \end{aligned} \quad (22)$$

This identification unifies the sources of interaction as manifestations of the same underlying conservation principle, arising from the symmetries of the substrate.

### C. Modal Dynamics from the Unified Lagrangian

The universal field equation derived earlier,

$$\kappa \square \mathcal{G}_{\mu\nu} - \frac{\partial V}{\partial \mathcal{G}^{\mu\nu}} = J_{\mu\nu},$$

when projected onto different modes, reproduces the macroscopic laws of physics:

- **Gravitational mode ( $\mathcal{G}_{00}$ ):** In the static, massless limit the equation reduces to

$$\nabla^2 \Phi_g = 4\pi G \rho_m,$$

i.e. Newton’s law of gravitation. At higher order, we will show later in the Paper how the full dynamics recovers the Einstein field equations.

- **Electromagnetic modes ( $\mathcal{G}_{0i}$ ):** The corresponding equations reduce to Maxwell’s laws, with current  $J^\mu$  acting as the source of transverse torsional excitations.

- **Other tensor deformation modes ( $\mathcal{G}_{ij}$ ):** The isotropic component describes large-scale volumetric expansion, while the traceless component propagates as shear-like excitations. In Section XX we will deduce how in the hydrodynamic limit these reproduce diffusion-type equations and the thermodynamic identities associated with the entropy flow.

Thus, all known field equations are recovered as modal reductions of the same substrate dynamics, validating the unification scheme.

*Clarification.* It is important to stress that gravitational phenomena split into two complementary aspects in this framework. The Newtonian potential and the coupling to mass-energy density ( $T_{00}$ ) arise from the scalar sector  $\mathcal{G}_{00}$ . In contrast, *gravitational radiation* is not encoded in this scalar but in the traceless tensorial excitations of  $\mathcal{G}_{ij}$ . Thus, while  $\mathcal{G}_{00}$  governs the scalar (compressive) aspect of gravitation, the tensorial shear modes  $\Sigma_{ij}$  account for the full propagation of gravitational waves. Consistently, only the traceless spatial components couple to the anisotropic stress  $\pi_{ij}$ , while the scalar potential couples to the rotationally invariant source  $T_{00}$ , preserving the matching of tensorial ranks between sources and fields required by Lorentz covariance.

### D. Lagrangian framework for modal excitations

Having established the physical identity of each deformation mode, we now construct the canonical Lagrangian framework required to describe their dynamics and interaction with sources. Building on the minimal scalar field formulation—which established that any modal projection must obey Lorentz-invariant dynamics—we transition from the fundamental Lagrangian governed by the stiffness constant  $\kappa$  to an effective, canonical Lagrangian for each mode. This is achieved via field renormalization, a standard procedure in field theory where the physical constant  $\kappa$  is absorbed into the field’s definition (e.g.,  $\mathcal{G}' = \sqrt{\kappa} \mathcal{G}$ ) to simplify the kinetic term.

This formulation allows the modal excitations of the unified tensor  $\mathcal{G}_{\mu\nu}$  to be described in terms of renormalized, scalar-like fields  $\mathcal{G}^{(X)}$ . The dynamics of these modes, including potential mass terms that arise from the full theory’s potential term  $V(\mathcal{G}_{\mu\nu})$ , can be described by the following canonical Klein–Gordon Lagrangian:

$$\mathcal{L}_X = \frac{1}{2} \partial^\mu \mathcal{G}^{(X)} \partial_\mu \mathcal{G}^{(X)} - \frac{1}{2} m_X^2 (\mathcal{G}^{(X)})^2, \quad (23)$$

where  $\mathcal{G}^{(X)}$  represents a renormalized projection of the full tensor  $\mathcal{G}_{\mu\nu}$  along mode  $X$ .<sup>1</sup> The term  $m_X$  is not a new fundamental parameter, but an *effective mass* that a mode acquires due to the self-interaction potential of the underlying unified field. This Klein–Gordon-type Lagrangian provides a Lorentz-invariant basis for describing both massless ( $m_X = 0$ ) and massive field modes within the unified elastic framework.

The Euler–Lagrange equation yields:

$$\square \mathcal{G}^{(X)} + m_X^2 \mathcal{G}^{(X)} = 0, \quad (24)$$

which reduces to the Klein–Gordon equation for free scalar propagation in Minkowski space.

Although the Lagrangian  $\mathcal{L}_X$  describes free fields, coupling to physical sources can be incorporated via minimal interaction terms of the form:

$$\mathcal{L}_{\text{int}}^{(X)} = \mathcal{G}^{(X)}(x) \cdot J^{(X)}(x), \quad (25)$$

where  $J^{(X)}$  is an effective source density corresponding to charge, mass, entropy flux, etc. These terms play an analogous role to the coupling  $A_\mu J^\mu$  in electrodynamics. Since  $\mathcal{G}^{(X)}$  represents a modal projection of the unified tensor  $\mathcal{G}_{\mu\nu}$ , the coupling is assumed to act only on the relevant scalarized or vectorial component associated with the physical mode.

A more general coupling scheme could link the full tensor to the energy–momentum content of matter via:

$$\mathcal{L}_{\text{int}} = \mathcal{G}_{\mu\nu}(x) T^{\mu\nu}(x), \quad (26)$$

from which each modal interaction  $\mathcal{G}^{(X)} J^{(X)}$  would arise as a projection or contraction. This formulation ensures full compatibility with general relativistic coupling schemes.

*Cross-check.* In Secs. XXE and XXG we show that the universal dissipative invariant, implemented via a Rayleigh functional, yields a hyperbolic–parabolic (telegrapher) dynamics at the effective level. This mirrors causal relativistic hydrodynamics (Israel–Stewart), guaranteeing finite signal speeds and positive entropy production in the gravito-entropic sector.

## E. Static solutions

In the static limit and for massless modes ( $m_X = 0$ ), the sourced Euler–Lagrange equation derived from  $\mathcal{L}_X + \mathcal{L}_{\text{int}}^{(X)}$  reduces to the Poisson equation

$$\nabla^2 \Phi_X(\vec{r}) = -J^{(X)}(\vec{r}), \quad (27)$$

where we denote by  $\Phi_X$  the static potential associated to the mode  $X$ . For a point-like unit source located

at the origin,  $J^{(X)}(\vec{r}) = \delta^{(3)}(\vec{r})$ , the Green’s function solution is

$$\Phi_X(r) = \frac{1}{4\pi r}. \quad (28)$$

Multiplying this fundamental response by an effective charge  $C_X$  yields the modal potential:

$$\Phi_X(r) = C_X \frac{1}{4\pi r} \quad (29)$$

which matches the expressions summarized in Sec. XXIII.

## IV. THE UNIFIED DIMENSIONAL FRAMEWORK

### A. Geometrization of Physical Quantities

The Spacetime Equivalence Principle ( $[M] \equiv [L] \equiv [T]$ ) forces a complete rewriting of the dimensional structure of physics. All fundamental physical quantities can be expressed in terms of a single geometric unit,  $[L]$ . The consequences, summarized in Table I, are profound: physical sources like mass and charge acquire the dimension of length, while all dynamical and coupling constants become dimensionless pure numbers.

In this new framework, one constant stands apart: **Planck’s constant**,  $h$ . Its dimensions of action,  $[E \cdot T]$ , do not collapse to  $[L]$  or  $[1]$ , but instead become:

$$[h] = [E \cdot T] \equiv [L \cdot L] = [L^2]. \quad (30)$$

This is a profound result. It reveals that the quantum of action is not an abstract parameter, but a fundamental quantum of **spacetime area**. This geometrization of quantum mechanics is a cornerstone of the QEG framework, linking the discreteness of the quantum world to the geometric fabric of the substrate.

### B. Dimensional Equivalence and Its Implications for Fundamental Units

Through the dimensional collapse, we establish that all fundamental physical quantities share a common geometric basis:

$$[L] \equiv [T_{\text{ime}}] \equiv [M] \equiv [E] \equiv [Q] \equiv [T_{\text{emp}}]. \quad (31)$$

This does not negate the operational distinction among meters, seconds, or kilograms in experimental practice, but clarifies the role of universal constants. *In this framework, any constant not itself a source of deformation (mass-energy, charge, thermal or entropic sources, action) becomes dimensionless and functions as a conversion factor.* Hence, SI units, once normalized by these constants, are numerically equivalent expressions of a single underlying geometric quantity:

$$1 \text{ m} \equiv 1 \text{ s} \equiv c^2 \text{ kg} \equiv 1 \text{ J} \equiv 1 \text{ C} \equiv k_B \cdot 1 \text{ K}. \quad (32)$$

This equivalence is not a matter of notation, but of geometry: universal constants act as the *metric coefficients of physical space*, establishing the calibration between nominal units. Their role is thus

<sup>1</sup> While this Klein–Gordon form correctly captures the universal mass and propagation dynamics for any mode, a full description for vector or tensor modes would require a more structured Lagrangian (e.g., a Proca Lagrangian for massive vector modes) to account for all degrees of freedom. For the purpose of describing the fundamental dynamics of sourced and source-free propagation, this canonical form is sufficient.

TABLE I. Dimensional collapse of physical quantities in the QEG framework.

Quantity	Standard SI Dimensions	QEG Dimension [L]
<i>Physical Sources</i>		
Mass	[M]	[L]
Energy	[ML <sup>2</sup> T <sup>-2</sup> ]	[L]
Charge	[IT]	[L]
Temperature	[K]	[L]
<i>Dynamical/Coupling Constants</i>		
Velocity / Current	[LT <sup>-1</sup> ]	[1] (Dimensionless)
Resistance	[ML <sup>2</sup> T <sup>-3</sup> I <sup>-2</sup> ]	[1]
Permittivity, $\epsilon_0$	[M <sup>-1</sup> L <sup>-3</sup> T <sup>4</sup> I <sup>2</sup> ]	[1]
Permeability, $\mu_0$	[MLT <sup>-2</sup> I <sup>-2</sup> ]	[1]
Gravitational C., $G$	[M <sup>-1</sup> L <sup>3</sup> T <sup>-2</sup> ]	[1]
Boltzmann C., $k_B$	[ML <sup>2</sup> T <sup>-2</sup> K <sup>-1</sup> ]	[1]

not arbitrary, but structural, ensuring that all units reduce consistently to a unified geometric basis.

While the theory itself is scale-free, connecting it to empirical quantities requires a fiducial reference, naturally chosen as the SI system since CODATA values are defined within it. This calibration should not be mistaken for introducing a free parameter: setting a reference length, e.g.  $L_{\text{ref}} = 1$  m, serves as a **coherence test**. By anchoring the theoretical unit to an experimental one, the entire system of SI constants emerges consistently from the predicted relations. The significance lies not in predicting numerical values, but in validating that the structure of physical laws is compatible with a unified geometric origin.

## Part II: General Properties of Geometro-elastic Excitations

Building on the foundational framework of Part I, this Part focuses on the *general, mode-independent* consequences that any excitation of the unified substrate must satisfy. We begin by deriving the operator form of the substrate's elastic response from the unified Lagrangian. Projecting this onto its normal modes, we obtain a universal Hooke-type constitutive law where the geometric deformation (action) is proportional to the applied source. This yields a single, powerful source law,  $Q = kS$ , valid for mass, charge, and any fundamental deformation mode. This structure then provides the basis for understanding the duality of stiffness and compliance and the dissipative nature of the substrate.

### V. A UNIFIED CONSTITUTIVE LAW FROM THE GENERAL ELASTIC EQUATION

#### A. Operator form of the elastic response

Starting from the unified dynamics of Part I,

$$\kappa \square \mathcal{G}_{\mu\nu} - \frac{\partial V}{\partial \mathcal{G}^{\mu\nu}} = J_{\mu\nu}, \quad (33)$$

the linear, long-wavelength (massless) regime relevant for static and weakly time-dependent fields is obtained by neglecting the non-linear and massive parts of  $V$ . In this regime one may write, schematically,

$$\mathbb{K} \mathcal{G} = J, \quad \mathbb{K} := \kappa \square \xrightarrow{\text{static}} -\kappa \nabla^2, \quad (34)$$

where  $\mathbb{K}$  is the *stiffness operator* of the substrate. The solution is expressed via the inverse (Green operator)  $\mathbb{C} := \mathbb{K}^{-1}$ , which plays the role of a *compliance operator*:

$$\mathcal{G} = \mathbb{C} J, \quad \mathbb{C} \equiv \mathbb{K}^{-1}. \quad (35)$$

Equations (34)–(35) are the covariant, field-theoretic generalization of Hooke's law (stress = stiffness  $\times$  strain; strain = compliance  $\times$  stress) and hold for *any* tensorial component of the deformation field  $\mathcal{G}_{\mu\nu}$  and source  $J_{\mu\nu}$ .

#### B. Modal projection: from fields to lumped constitutive law

In homogeneous/isotropic backgrounds the operator  $\mathbb{K}$  diagonalizes on normal modes. Let  $\Pi_i$  denote the projector onto mode  $i$  (scalar, vectorial, or tensorial). Projecting (34) we obtain the algebraic response for each mode amplitude,

$$k_i \mathcal{G}_i = J_i, \quad \Rightarrow \quad \mathcal{G}_i = \frac{1}{k_i} J_i, \quad (36)$$

where  $k_i > 0$  is the *modal stiffness*. Equation (36) is precisely a Hooke law at the modal level: *deformation equals compliance times source*.

#### C. Action as geometric deformation and the universal constitutive law

From Part I we established the dimensional collapse  $[M] \equiv [L] \equiv [T]$  and that action has the character of a spacetime area,  $[S] = [L^2]$ . For a normalized unit 4-volume (the fundamental cell of the substrate), the scalar measure of deformation carried by mode  $i$  is thus naturally identified with a *modal action*  $S_i$  proportional to  $\mathcal{G}_i$ . Similarly, the projected source  $J_i$  defines the *modal charge*  $Q_i$  (mass, electric charge, thermal/entropic source, *etc.*) through the same normalization cell. With this identification, (36) becomes the *unified Hooke law* for sources:

$$\boxed{S_i = \frac{1}{k_i} Q_i}, \quad \Leftrightarrow \quad \boxed{Q_i = k_i S_i}. \quad (37)$$

That is, *the deformation (action/area) required to stabilize a mode equals the applied source times the modal compliance*. Equivalently, *the source equals the modal*

*stiffness times the action.* This is the sought-for generalization: the law holds verbatim for compressive (mass-like), torsional (electromagnetic), and tensorial (thermo-entropic) modes.

#### D. Why this form is inevitable

The operator identity  $\mathcal{G} = \mathcal{C}J$  is fixed by covariance and linearity around equilibrium. Its modal reduction  $\mathcal{G}_i = J_i/k_i$  is forced by homogeneity/isotropy. With action identified as the geometric measure of deformation in the normalized cell,  $S_i \propto \mathcal{G}_i$ , the constitutive law (37) is not an ansatz but the *unique linear, covariant map* between sources and deformations compatible with the unified Lagrangian of Part I. Any alternative would either break covariance, violate linear superposition near equilibrium, or introduce extra dimensional constants contrary to the Spacetime Equivalence Principle.

#### E. Sign conventions and stability

The minus signs familiar from restoring forces (e.g. Hooke's law  $F = -kx$ ) are absorbed here in the definition of  $J$  as the *applied* source that balances the internal restoring response. Stability requires  $k_i > 0$  for all propagating modes, ensuring that the quadratic energy is bounded below and that  $\mathbb{K}$  is (elliptic/hyperbolic) invertible in the corresponding regime.

### VI. THE PRINCIPLE OF MODAL RECIPROcity: STATIC STABILITY AND DYNAMIC CAUSALITY IN THE SUBSTRATE

#### A. Static Reciprocity: The Condition of Stability and the Hierarchy of Forces

For a single, unified substrate to be both stable (not collapsing) and excitable (not infinitely rigid), its elastic properties across orthogonal modes cannot be independent. We elevate this physical requirement to a fundamental *Principle of Modal Reciprocity*. This principle states that *the stiffness of the substrate in a given mode is inversely related to its stiffness in an orthogonal mode.*

For the primary compressive (longitudinal,  $\parallel$ ) and torsional (transverse,  $\perp$ ) modes, this principle is formalized as a stability condition:

$$k_{\parallel} \cdot k_{\perp} = \mathcal{C}_{\text{geom}} \cdot \kappa^2, \quad (38)$$

where  $\kappa$  is the universal stiffness constant of the substrate from Part I, and  $\mathcal{C}_{\text{geom}}$  is a dimensionless geometric factor of order unity reflecting the isotropy of three-dimensional space. As will be shown in Sec. XXVIII, this factor can be derived from the geometric integration of self-energy in spherical coordinates.

Equation (38) guarantees the stability of the geometro-elastic medium. It expresses the fact that the medium's resistance to longitudinal strain is reciprocally balanced by its resistance to transverse strain. A stiffer electromagnetic response (large  $k_{\perp}$ ) necessarily entails a more compliant, or "softer," gravitational response (small  $k_{\parallel}$ ). Any deviation from this reciprocity would render the substrate either

undeformable (if both stiffnesses are large) or unstable (if both are small).

#### Physical interpretation

Equation (38) captures the observed hierarchy of interactions without arbitrary assumptions:

- The electromagnetic field corresponds to torsional excitations with extremely high stiffness  $k_{\perp}$ , consistent with its strength and short-range rigidity.
- Gravity corresponds to compressive excitations with extremely low stiffness  $k_{\parallel}$ , consistent with its weakness and long-range compliance.

Thus, the empirical disparity between electromagnetic and gravitational strengths is not accidental but a direct manifestation of the reciprocal balance demanded by a single geometro-elastic substrate.

#### Inevitability of the duality

The duality between stiffness and compliance follows inevitably from three principles already established:

1. The *Spacetime Equivalence Principle* ( $[M] \equiv [L] \equiv [T]$ ) reduces all dimensional assignments to geometry.
2. The *unified constitutive law*  $Q_i = k_i S_i$  requires each source to be stabilized by a finite stiffness.
3. The *single-substrate hypothesis* ensures that all modal stiffnesses derive from the same  $\kappa$ , which enforces the reciprocity relation (38).

Any violation of this reciprocity would break energy balance in the substrate: if both  $k_{\parallel}$  and  $k_{\perp}$  were simultaneously large, the medium would forbid deformation altogether (contradicting the existence of excitations); if both were small, the medium would collapse under any source (contradicting stability). The hierarchy of forces is therefore *not contingent but structurally inevitable*, rooted in the duality of stiffness and compliance in a unified elastic spacetime.

#### Conclusion

We conclude that the extreme weakness of gravitation relative to electromagnetism arises as the elastic complement of the substrate: one mode is strong because the other is weak. This stiffness-compliance duality is not an optional feature but the only configuration that allows a single geometro-elastic substrate to remain both deformable and stable. In the next section we extend this framework by introducing the concept of *dissipation*, showing how the intrinsic "viscosity" of the substrate gives rise to the dimensionless fine-structure constant  $\alpha$ .

#### B. Dynamic Reciprocity: The Causal Universality Condition

Beyond the static stability condition, the Principle of Modal Reciprocity finds a profound dynamic manifestation rooted in a core axiom of the theory:

**Lorentz Invariance.** For the QEG framework to be self-consistent, all emergent massless, wave-like excitations of the unified substrate must propagate at the same universal speed,  $c$ . This *principle of Causal Universality* imposes a strict constraint on the constitutive properties of any given mode.

Let us define the effective inertial (permeability-like) and compliant (permittivity-like) constants for an arbitrary mode ' $i$ ' as  $\mu_i$  and  $\varepsilon_i$ , respectively. The propagation speed of a wave in this modal sector is given by  $v_i = 1/\sqrt{\mu_i\varepsilon_i}$ . The Causal Universality Condition therefore requires that for every wave-like mode  $i$ :

$$\mu_i\varepsilon_i = \frac{1}{c^2}. \quad (39)$$

Furthermore, the theory establishes a hierarchy of interactions, where the inertial response of any mode  $i$  can be expressed as a scaling of a baseline inertia,  $\mu_{\text{base}}$ . Let this be described by a dimensionless scaling factor  $s_i$ , such that:

$$\mu_i = s_i \cdot \mu_{\text{base}}. \quad (40)$$

By combining these two conditions, the dynamic nature of reciprocity becomes explicit. Substituting Eq. (40) into Eq. (39) and solving for the modal compliance  $\varepsilon_i$ , we find:

$$(s_i \cdot \mu_{\text{base}})\varepsilon_i = \frac{1}{c^2} \implies \varepsilon_i = \frac{1}{s_i} \left( \frac{1}{\mu_{\text{base}}c^2} \right). \quad (41)$$

Recognizing that the term in parenthesis is simply the baseline compliance,  $\varepsilon_{\text{base}}$ , we arrive at the general law of dynamic reciprocity:

$$\boxed{\varepsilon_i = \frac{1}{s_i} \cdot \varepsilon_{\text{base}}} \quad (42)$$

This result is a cornerstone of the theory. It rigorously demonstrates that if the inertial response of a mode is scaled by a factor  $s_i$ , its compliant response must be scaled by the exact inverse factor,  $1/s_i$ . This dynamic reciprocity is not an isolated coincidence but the second core manifestation of the Principle of Modal Reciprocity, complementing the static relationship between stiffnesses. It confirms that the vacuum's elastic properties are deeply interconnected through both stability and causality, with the capacity to yield in one mode being intrinsically linked to its resistance in another.

## VII. THE DISSIPATIVE NATURE OF THE SUBSTRATE AND THE ORIGIN OF THE FINE-STRUCTURE CONSTANT

The Lagrangian presented in Part I,  $\mathcal{L}_{\text{QEG}}$ , describes an ideal, perfectly elastic medium. However, no physical medium can be perfectly elastic: stability under real excitations requires a mechanism of *dissipation*. Without it, oscillations would grow without bound and localized excitations could not stabilize, contradicting the observed persistence of particles and fields.

Indeed, the necessity of a dimensionless dissipative parameter follows inevitably from three prior requirements:

1. The *unified constitutive law* ( $Q_i = k_i S_i$ ) demands finite, stable deformations.

2. The *principle of modal reciprocity* requires complementary stiffnesses across orthogonal modes, but does not by itself suppress divergences in excitations.
3. The *single-substrate hypothesis* forbids the introduction of independent damping constants for each mode: dissipation must be controlled by a unique, universal invariant.

Taken together, these conditions leave no freedom: the substrate must carry a single, dimensionless measure of dissipation. This is entirely analogous to the *damped harmonic oscillator*, where the addition of a velocity-proportional term is the unique way to stabilize oscillations without altering equilibrium. By covariance, the same logic applies at the field level: the equation of motion must include a dissipative contribution proportional to the "velocity" of the deformation field.

The most general covariant modification of the field equations is

$$\kappa \square \mathcal{G}_{\mu\nu} - \frac{\partial V}{\partial \mathcal{G}^{\mu\nu}} - \zeta_{\mu\nu}^{\alpha\beta} \partial_0 \mathcal{G}_{\alpha\beta} = J_{\mu\nu},$$

where  $\zeta_{\mu\nu}^{\alpha\beta}$  is a dimensionless tensor encoding the intrinsic damping of the substrate. Its presence is unavoidable: it is the unique covariant structure that introduces dissipation without violating Lorentz invariance or adding new dimensional scales.

Because the Spacetime Equivalence Principle collapses  $[M]$ ,  $[L]$ , and  $[T]$  into geometry, no new dimensional constants can enter consistently. Dissipation must therefore be governed by a *dimensionless invariant*. We define its scalar norm as

$$\alpha \equiv \|\zeta_{\mu\nu}\|, \quad (43)$$

### 1. Physical Origin of the Damping Coefficient

The introduction of a dissipative term into the substrate's dynamics requires the identification of a dimensionless damping constant. We show that this role is uniquely filled by the *fine-structure constant*,  $\alpha$ . This is not an arbitrary choice. Within the Standard Model,  $\alpha$  governs the fundamental vertex of Quantum Electrodynamics (QED), which describes the quantum interaction between light and matter. More fundamentally, QED is the theory of the vacuum's own self-interaction, mediated by the continuous creation and annihilation of virtual particle-antiparticle pairs. In the QEG framework, dissipation is precisely the mechanism by which a coherent deformation (a field excitation) loses energy to the underlying quantum fluctuations of the substrate. It is therefore natural to identify the universal damping ratio of the medium with the very constant that quantifies its fundamental quantum self-interaction. Thus, in QEG,  $\alpha$  is elevated from a purely electromagnetic coupling constant to a universal property of the spacetime substrate: its intrinsic dissipative coefficient.

### A. Confirmation from the Vacuum's Oscillatory Response

This theoretical identification is powerfully confirmed by analyzing the vacuum as an oscillatory medium, analogous to an RLC circuit. The fine-structure constant  $\alpha$  [28] can be defined as the ratio of two energies:

- the energy needed to overcome the electrostatic repulsion between two electrons a distance of  $d$  apart
- the energy of a single photon of wavelength  $\lambda = 2\pi d$  (or of angular wavelength  $d$ )

Therefore, we have that

$$\begin{aligned}\alpha &= \left( \frac{e^2}{4\pi\epsilon_0 d} \right) / \left( \frac{\hbar c}{\lambda} \right) \\ &= \frac{e^2}{4\pi\epsilon_0 d} \times \frac{2\pi d}{\hbar c} = \frac{e^2}{4\pi\epsilon_0 d} \times \frac{d}{\hbar c} \\ &= \frac{e^2}{4\pi\epsilon_0 \hbar c}\end{aligned}\quad (44)$$

Other hand, in the context of an RLC circuit, the quality factor or Q factor [29] is a dimensionless parameter that describes how underdamped an oscillator or resonator is. It is defined as the ratio of the initial energy stored in the resonator to the energy lost in one radian of the cycle of oscillation. Therefore, we have that

$$\begin{aligned}Q &\stackrel{\text{def}}{=} 2\pi \times \frac{\text{Energy stored}}{\text{Energy dissipated per cycle}} \\ &= 2\pi f_r \times \frac{\text{Energy stored}}{\text{Power loss}} \\ &= \omega_0 \times \frac{\text{Energy stored}}{\text{Power loss}}\end{aligned}\quad (45)$$

Where  $f_r$  is the resonance frequency.

The fine-structure constant can be written as

$$\alpha = \frac{1}{2} Z_0 \sigma,$$

where  $Z_0 = \mu_0 c = \frac{1}{\epsilon_0 c}$  is the vacuum impedance and  $\sigma = \frac{e^2}{\hbar}$  is the conductance quantum [30, 31]. It follows that

$$Z_0 \sigma = 2\alpha,$$

which can be interpreted as the intrinsic energy loss characteristic per radian for the vacuum medium itself. Thus, one can naturally define a vacuum quality factor as

$$Q = \frac{1}{Z_0 \sigma} = 2\pi \times \frac{\epsilon_0 \hbar c}{e^2} = \frac{1}{2\alpha}.$$

where we can identify  $E_{\text{stored}} = \frac{\hbar c}{\lambda}$  and  $E_{\text{dissipated}} = \frac{e^2}{\epsilon_0 \lambda}$ .

The appearance of these two energies follows directly from well-established properties of a single electromagnetic vacuum mode of wavelength  $\lambda$  and the framework we have discussed throughout this Paper. On the one hand, the most fundamental quantum excitation of an electromagnetic mode is a single photon. The energy of this photon, which represents the total energy “stored” in the oscillating mode for this process, is given by the Planck-Einstein relation:

$$E_{\text{stored}} = \hbar\omega = \frac{\hbar c}{\lambda}$$

where the energy of a single photon of wavelength  $\lambda = 2\pi d$  is used. This interpretation defines  $E_{\text{stored}}$  as the energy of the fundamental quantum of light that populates and defines the oscillation within that

vacuum mode.

On the other hand, within our elastic vacuum formalism, it is natural to propose a dissipated energy analogous to Hooke’s law. As we will demonstrate below with a rigorous calculation based on Larmor’s formula, this simple analogy predicts with great accuracy the energy dissipated by a quantum dipole.

- Using Hooke’s Law, we can identify the dissipated energy using the formula  $|E| = |kx^2|$  where  $k$  is the elasticity constant, and  $x$  the displacement [32]. As in the context of the unified field we have identified  $k = \frac{1}{C} = \frac{1}{\epsilon_0 \lambda}$ , and the displacement  $x$  with the elementary charge  $e$ , we get that  $E_{\text{dissipated}} = \frac{e^2}{\epsilon_0 \lambda}$ .

- Alternatively, consider an oscillating dipole  $p(t) = p_0 \cdot \cos(\omega t)$  with amplitude  $p_0 = e \cdot d$ , where  $d = \lambda/(2\pi)$  is the separation and  $\omega = c/d$  is the angular frequency. Using Larmor formula for time-averaged power radiated by an oscillating dipole [33, 34]:

$$\langle P \rangle = \frac{p_0^2 \omega^4}{12\pi\epsilon_0 c^3}\quad (46)$$

we can derive the energy dissipated per cycle (period  $T = 2\pi/\omega$ ):

$$E_{\text{dissipated}} = \langle P \rangle \times T = \left( \frac{p_0^2 \omega^4}{12\pi\epsilon_0 c^3} \right) \left( \frac{2\pi}{\omega} \right)\quad (47)$$

And one finally gets that

$$E_{\text{dissipated}} = \frac{\pi e^2}{3\epsilon_0 \lambda} \approx 1.05 \cdot \frac{e^2}{\epsilon_0 \lambda}\quad (48)$$

Thus, both energies can be derived from first-principles and lead directly to

$$Q = 2\pi \times \frac{E_{\text{stored}}}{E_{\text{dissipated}}} = 2\pi \times \frac{\hbar c/(\lambda)}{e^2/(\epsilon_0 \lambda)} = \frac{1}{2\alpha},$$

For an underdamped oscillator, the damping ratio is defined as

$$\zeta = \frac{1}{2Q},$$

which leads directly to

$$\boxed{\zeta = \alpha}$$

This perfect correspondence between the first-principles requirement for a dissipative invariant and the result from the vacuum’s oscillatory dynamics provides strong evidence that  $\alpha$  is the universal damping ratio of the geometro-elastic medium.

## B. Interpreting $c$ in terms of the Damped Resonant Frequency of the System

In a standard underdamped oscillator model [35–37], the damped frequency  $\omega_d$  is given by

$$\omega_d = \omega_0 \sqrt{1 - \zeta^2},\quad (49)$$

where  $\omega_0$  is the *undamped* resonant (natural) frequency of the system, and  $\zeta$  is the damping ratio.

We can associate these frequencies with propagation speeds by multiplying each angular frequency by the reference length within our framework, of *one meter*, yielding speeds in units of  $\text{m s}^{-1}$ . Denoting:

$$v_{\text{damped}} = \omega_d \times 1 \text{ m}, \quad v_{\text{undamped}} = \omega_0 \times 1 \text{ m},$$

we can identify  $v_{\text{damped}}$  with the *measured* speed of light, conventionally denoted by  $c$ . In other words,

$$c = v_{\text{damped}} = \omega_d \times 1 \text{ m}.$$

From Eq. (49), we thus have

$$c = \omega_0 \cdot 1 \text{ m} \sqrt{1 - \zeta^2}$$

Or, equivalently,

$$c_{\text{measured}}^2 = c_{\text{real}}^2 (1 - \alpha^2) \quad (50)$$

Which, solving for  $\zeta$ , can be rewritten as

$$\zeta = \alpha = \sqrt{1 - \frac{c_{\text{measured}}^2}{c_{\text{real}}^2}} \quad (51)$$

Note the similarity of the above expression with the reciprocal of the Lorentz factor formula [38]. Thus, the fine-structure constant  $\alpha$  can be regarded as the reciprocal of a ‘‘Lorentz-like’’ factor via

$$\zeta = \alpha = \frac{1}{\gamma} = \sqrt{1 - \frac{c_{\text{measured}}^2}{c_{\text{real}}^2}}. \quad (52)$$

These two views—the *damped oscillator* analogy for electromagnetic propagation and the *Lorentz-like* factor interpretation for  $\alpha$ —are not only compatible, but in fact reinforce each other:  $\alpha$  emerges as a geometric or relativistic ‘‘scaling factor’’ that governs attenuation in the oscillatory unified field, connecting electromagnetic propagation and the quantum vacuum’s dissipative properties.

### C. Dissipative duality: roles of the damping factor $\zeta = \alpha$ and the reciprocal of the quality factor $\frac{1}{Q} = 2\alpha$

*Physical Origin: The Fermionic g-factor*

The theoretical framework that we will develop reveals an apparent duality in the vacuum’s dissipative response. On one hand, the damping that affects wave propagation manifests as a relativistic damping ratio  $\zeta = \alpha$ . On the other hand, the attenuation of energy in quantum interactions, quantified by the quality factor  $Q$ , is governed by  $1/Q = 2\alpha$ .

We propose that this duality is not a contradiction but a reflection of a deeper physical principle: *the factor of 2 originates from the Landé g-factor ( $g_e$ ) of a fundamental, spin-1/2 fermionic excitation of the vacuum*, whose value predicted by the Dirac Equation is precisely  $g_e = 2$ . This hypothesis establishes a fundamental distinction between two dissipative regimes:

- **Bosonic Propagation Damping ( $\zeta$ ):** The factor  $\zeta = \alpha$  represents the intrinsic damping experienced by a bosonic excitation (such as a spin-1 photon) as it propagates through the elastic vacuum medium. It is a pure measure of spacetime’s ‘‘viscosity.’’

- **Fermionic Interaction Damping ( $1/Q$ ):**

The factor  $1/Q = g_e \alpha \approx 2\alpha$  governs processes that involve the fermionic nature of the vacuum’s excitations. This includes the ‘‘dressing’’ of a bare charge to form a stable particle (like the electron) or the energy transfer that gives rise to a quantum excitation. In these cases, the dissipation depends not only on the base viscosity ( $\alpha$ ) but is also amplified by the intrinsic spin-1/2 response ( $g_e = 2$ ) of the excitation.

Far from being an electron-centric hypothesis, this interpretation posits that  $g_e = 2$  is a topological property of the fermionic modes of the vacuum lattice itself. Particles like the electron simply inherit this fundamental characteristic. This idea is consistent with the treatment of the anomalous magnetic moment ( $a_e = (g_e - 2)/2$ ) as a higher-order correction to the same dissipative mechanism.

#### *Analogy with the RLC Circuit: Transient Damping vs. Resonant Dissipation*

This physical distinction between  $\zeta$  and  $Q$  finds a powerful analogue in the behavior of a classical RLC circuit, which strengthens its conceptual justification:

- The **damping ratio ( $\zeta$ )** in an RLC circuit determines the system’s *transient response* (i.e., how a free oscillation decays). Analogously,  $\zeta = \alpha$  in our model governs the decay of a wave propagating freely through the vacuum.
- The **quality factor ( $Q$ )**, in contrast, describes the *steady-state or resonant response*. It is defined as the ratio of reactive impedance to dissipative resistance, quantifying the energy loss *per cycle* in an established oscillation. Analogously,  $1/Q = g_e \alpha$  in our model governs the energy dissipation in a stable interaction, such as that which constitutes a ‘‘dressed’’ particle, where vacuum energy is continuously stored and dissipated.

Therefore, the vacuum is interpreted as a medium possessing a base friction for wave propagation ( $\zeta = \alpha$ ) and a distinct, amplified dissipative resistance for fermionic interactions ( $1/Q = g_e \cdot \alpha$ ). This interpretation resolves the apparent duality by unifying the vacuum’s relativistic properties with its quantum, spin-dependent interactions.

### D. Interpretation and discussion on Vacuum Damping

Throughout this Paper, we are proposing that the quantum vacuum itself acts as a structured, elastic-dissipative medium. In this picture, the vacuum consists of fluctuating virtual excitations with internal degrees of freedom that collectively endow it with both stiffness (reactive elasticity) and finite relaxation time (viscosity). A propagating electromagnetic wave then loses energy—not into real particles, but into the hidden structure of the vacuum—via coupling to these degrees of freedom. This loss manifests macroscopically as a damping ratio  $\zeta$ , which we identify with the dimensionless fine-structure constant via  $\zeta = \alpha$ . Such a damping constant is natural if one treats the vacuum as an ensemble of coupled oscillators or as an emergent condensed-matter system, as suggested by various approaches to quantum gravity and emergent spacetime [39–41]. The resulting reduction in propagation speed is then not a kinematic effect, but a

first-principles consequence of quantum back-reaction. This allows us to interpret the measured speed of light  $c$  as a damped, effective velocity arising from the underlying dissipative structure of the vacuum.

Importantly, this view does not violate local Lorentz invariance. All local observers measure the same effective speed  $c = c_{\text{measured}}$ , and all physical laws remain Lorentz-invariant in that frame. The distinction between  $c_{\text{real}}$  and  $c$  thus becomes a global, geometric feature of the vacuum — akin to how curvature encodes gravitational effects in general relativity. In this case, however, the “curvature” is not geometric but modal: a manifestation of the vacuum’s internal damping modes, whose excitation state defines a preferred frame only at a topological level, not at the level of measurable kinematics. Analogous to an RLC circuit, where the “natural” frequency  $\omega_0$  is never directly observed but rather only inferred through modeling, the proposed  $c_{\text{real}} > c$  does not admit superluminal information transfer, and thus poses no contradiction to special relativity or experiment.

### E. Final notes

Higher-order radiative effects — e.g. the electron’s anomalous magnetic moment  $a_e = \alpha/2\pi$  — can be viewed as additional layers of the same dissipative mechanism. The damping encoded in the fine-structure constant  $\alpha$  would be the first-order manifestation of how the vacuum’s oscillator lattice “bleeds” energy back into itself through quantum fluctuations, and the anomalous magnetic moment can be viewed, in our framework, as the simplest radiative attenuation of a bare “undamped” coupling by the lattice’s elastic resistance. Higher-order Feynman diagrams then correspond to more intricate couplings among modes of the vacuum, each contributing successive powers of  $\alpha$ .

More generally, it implies that any relation among fundamental constants—whether in electromagnetism, gravitation or thermodynamics—must be dressed by a universal, dimensionless form factor  $\Xi_{\text{eff}}(\alpha)$  that encodes the accumulated effect of loop-induced damping within the vacuum lattice. In this way, radiative corrections are not mere perturbative afterthoughts, but the fingerprint of the same elastic and dissipative structure that unifies all fields at their quantum origin.

In this view, *the fine-structure constant  $\alpha$  becomes not merely a coupling constant, but a unifying signature of modal attenuation across all field interactions* — electromagnetic, gravitational, and thermo-entropic alike. Any consistent geometro-elastic substrate must contain a dimensionless dissipative invariant; its identification with the fine-structure constant  $\alpha$  is therefore not optional, but logically enforced by the unified framework.

## Part III: Dynamic of fields and sources of deformation in quantum-elastic geometry

### VIII. EMERGENCE OF THE CLASSICAL FIELD EQUATIONS

Having established the foundational principles of the QEG substrate, we now demonstrate how the established equations of General Relativity and Electromagnetism emerge as consistent, low-energy effective descriptions of the substrate’s dynamics. This section

provides the explicit derivations, bridging the microscopic model of  $\mathcal{G}_{\mu\nu}$  with the macroscopic physics of  $g_{\mu\nu}$  and  $F_{\mu\nu}$ .

#### A. From Substrate Dynamics to General Relativity

The unified field  $\mathcal{G}_{\mu\nu}$  contains the full microscopic description of the spacetime substrate. The smooth, classical geometry described by GR emerges as its macroscopic, coarse-grained average. We therefore define the effective metric  $g_{\mu\nu}$  as the statistical expectation value of the fundamental field:

$$g_{\mu\nu}(x) \equiv \langle \mathcal{G}_{\mu\nu}(x) \rangle \quad (53)$$

This *background-independent* definition is a cornerstone of the theory. It posits that what we perceive as “curved spacetime” is the cumulative, large-scale effect of the substrate’s deformation. This identification stems as a necessary consequence of the framework’s axioms: in IIB we established that all coupling constants reduce dimensionally to the same stiffness  $\kappa$ , and in IIG we showed that mass, charge, and entropy are endogenous modes of  $\mathcal{G}_{\mu\nu}$ . Therefore, *any macroscopic geometric descriptor must emerge from  $\mathcal{G}_{\mu\nu}$  itself*. No independent background is admissible without violating the Equivalence Principles already established. Moreover, Lorentz invariance requires that the coarse-grained substrate be represented by a symmetric rank-2 tensor. *The only covariant candidate is the statistical average of  $\mathcal{G}_{\mu\nu}$ , which we identify as the effective metric*. This ensures a direct correspondence between the microscopic modes and macroscopic curvature: scalar compressions ( $\langle \mathcal{G}_{00} \rangle$ ) manifest as time-time curvature, torsional modes ( $\langle \mathcal{G}_{0i} \rangle$ ) as frame-dragging, and tensorial strains ( $\langle \mathcal{G}_{ij} \rangle$ ) as spatial curvature.

#### The Emergence of the Einstein-Hilbert Action

The fundamental action  $S_{\text{QEG}}$  is a generally covariant functional of the microscopic symmetric field  $\mathcal{G}_{\mu\nu}$  of Lorentzian signature, which induces a valid four-volume form. We adopt the natural invariant measure  $\sqrt{-\det(\mathcal{G})} d^4x$  and a kinetic term built from the Levi-Civita connection of  $\mathcal{G}_{\mu\nu}$ . Explicitly,

$$S_{\text{QEG}} = \int d^4x \sqrt{-\det(\mathcal{G})} \left[ \frac{\kappa}{2} \mathcal{G}^{\alpha\beta} \mathcal{G}^{\mu\nu} \nabla_\alpha \mathcal{G}_{\mu\nu} \nabla_\beta \mathcal{G}_{\rho\sigma} \mathcal{G}^{\rho\sigma} - V(\mathcal{G}) \right] \quad (54)$$

where the schematic notation  $(\nabla\mathcal{G})^2$  used earlier is here understood as the positive-definite quadratic form built by contracting indices with  $\mathcal{G}^{\mu\nu}$ . Other equivalent kinetic choices related by field redefinitions or addition of covariant total derivatives lead to the same infrared (IR) effective theory after coarse-graining.<sup>2</sup>

*a. Effective Action and the Low-Energy Limit.* The classical action  $S_{\text{eff}}$  for the macroscopic metric  $g_{\mu\nu} = \langle \mathcal{G}_{\mu\nu} \rangle$  is obtained by integrating out the microscopic, high-frequency fluctuations of  $\mathcal{G}_{\mu\nu}$ . As  $S_{\text{QEG}}$

<sup>2</sup> One may equivalently write the kinetic term as  $\frac{\kappa}{2} \nabla_\alpha \mathcal{G}_{\mu\nu} \nabla^\alpha \mathcal{G}^{\mu\nu}$  up to covariant surface terms; the present form makes index contractions transparent and keeps manifest general covariance.

is generally covariant,  $S_{\text{eff}}$  must also be. In the low-energy limit, its Lagrangian density,  $\mathcal{L}_{\text{eff}}$ , can be expressed as a derivative expansion in powers of curvature. The leading terms are:

$$\mathcal{L}_{\text{eff}} = \sqrt{-g} (C_0 + C_1 R + \mathcal{O}(R^2)) \quad (55)$$

The coefficients are determined by the properties of the QEG substrate. The constant term  $C_0$  arises from the potential  $V(\mathcal{G})$  and corresponds to a cosmological constant,  $C_0 = -2\Lambda$ . The term linear in the Ricci scalar,  $C_1 R$ , must emerge from the kinetic term of the QEG action, which is governed by the universal stiffness  $\kappa$ . For the theory to be consistent, we establish a *correspondence principle* linking the microscopic damping to the macroscopic gravitational coupling:

$$C_1 = \frac{1}{16\pi G} \propto \kappa \quad (56)$$

This identification is central: it elevates the gravitational constant from an empirical parameter to a direct measure of the substrate's elastic stiffness. The effective Lagrangian thus naturally takes the Einstein-Hilbert form.

*b. Consistency with Uniqueness Theorems.* This result, derived from the internal logic of QEG, is strongly supported by external consistency arguments. The classic Weinberg-Deser argument shows that any consistent, universally coupled low-energy theory for the massless spin-2 excitations of  $\mathcal{G}_{\mu\nu}$  inevitably reconstructs the Einstein-Hilbert action. Furthermore, Lovelock's Theorem guarantees that in  $3+1$  dimensions, this action is unique for equations of motion with at most second derivatives of the metric. These theorems confirm that the form derived from our physical correspondence principle is not only natural within QEG, but is the only mathematically consistent possibility.

Including a matter action  $S_{\text{matter}}$  that couples to the effective metric, we have:

$$S_{\text{eff}}[g, \Psi_{\text{matter}}] = \int d^4x \sqrt{-g} \left( \frac{1}{16\pi G} R - 2\Lambda \right) + S_{\text{matter}} \quad (57)$$

Varying this with respect to  $g^{\mu\nu}$  yields the Einstein Field Equations:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G T_{\mu\nu}, \quad (58)$$

where  $G_{\mu\nu}$  is the Einstein tensor and the stress-energy tensor is defined as  $T_{\mu\nu} \equiv -\frac{2}{\sqrt{-g}} \frac{\delta S_{\text{matter}}}{\delta g^{\mu\nu}}$ . The coefficient  $G$  is fixed by requiring that the theory reproduces Newtonian gravity in the weak-field, static limit.<sup>3</sup> The  $g_{00}$  component of the equation becomes  $\nabla^2 g_{00} \approx 8\pi G T_{00}$ . Comparing this with the Poisson equation,  $\nabla^2 \Phi = 4\pi G \rho$ , and the definition  $g_{00} \approx -(1 + 2\Phi)$ , confirms the normalization chosen in the action.

*Newtonian check.* In the weak-field, static limit  $g_{00} \simeq -(1 + 2\Phi)$  and  $T_{00} \simeq \rho$ , the 00-component yields  $\nabla^2 \Phi = 4\pi G \rho$ , fixing the overall normalization of  $G$  in the action.

---

<sup>3</sup> We will demonstrate in subsequent parts of this work (Parts IV and V) that  $G$  is not a free parameter, but is instead determined by the fundamental elastic and dissipative properties of the substrate.

*Consistency Check: Symmetries and Conservation.* The general covariance of the effective theory leads to a crucial consistency check. The Einstein tensor is mathematically constructed to be divergenceless ( $\nabla^\mu G_{\mu\nu} \equiv 0$ ), which, via the field equations, enforces the covariant conservation of stress-energy:  $\nabla^\mu T_{\mu\nu} = 0$ . This aligns with the deep insight, articulated by Weinberg and others, that any theory of a massless, interacting spin-2 particle must inevitably lead to the principle of equivalence and the structure of General Relativity.

### Geodesics and the Principle of Least Deformation

In GR, the motion of free particles follows geodesics that extremize the spacetime interval. Within the substrate model, we reinterpret this principle dynamically: free trajectories minimize the integrated deformation of the substrate along their path.

Let  $\mathcal{D}[\gamma]$  denote the deformation functional associated with a worldline  $\gamma$ . A natural and covariant choice for this functional, representing the total integrated strain along a path, is the line integral measured by the fundamental field itself:

$$\mathcal{D}[\gamma] = \int \sqrt{-\mathcal{G}_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}} d\lambda \quad (59)$$

where  $\lambda$  is an arbitrary path parameter. Then the principle of least deformation is

$$\delta \mathcal{D}[\gamma] = 0, \quad (60)$$

which seeks worldlines that are extremal with respect to the microscopic geometry. Under the coarse-graining operation where  $\mathcal{G}_{\mu\nu}$  is replaced by its macroscopic average  $g_{\mu\nu}$ , this variational principle becomes  $\delta \int \sqrt{-g_{\mu\nu} dx^\mu dx^\nu} = 0$ , which is precisely the standard definition of a geodesic in General Relativity:

$$\frac{d^2 x^\mu}{d\tau^2} + \Gamma^\mu_{\nu\lambda} \frac{dx^\nu}{d\tau} \frac{dx^\lambda}{d\tau} = 0, \quad (61)$$

with  $\Gamma^\mu_{\nu\lambda}$  the Christoffel symbols of  $g_{\mu\nu}$ .

This correspondence is also inevitable. Since motion sources deformation in  $\mathcal{G}_{\mu\nu}$ , the least-deformation condition is the unique covariant generalization of free motion consistent with the Inertial Equivalence Principle. Under coarse-graining, it reduces to geodesic motion in the emergent metric  $g_{\mu\nu}$ .

Thus, the classical geodesic principle is reinterpreted as a manifestation of the substrate's elastic tendency to minimize cumulative strain, reinforcing the view of geometry as emergent rather than fundamental.

### Conclusion

We have shown that the effective spacetime metric  $g_{\mu\nu}$  emerges naturally from the deformation field  $\mathcal{G}_{\mu\nu}$ , as a macroscopic descriptor of substrate dynamics. The Einstein-Hilbert action appears as the large-scale limit of the unified Lagrangian, and geodesic motion follows from the principle of least deformation.

Hence, General Relativity is not a competing framework but an emergent approximation: the continuum geometry of GR arises from the coarse-grained behavior of the underlying quantum-elastic substrate.

## B. Emergence of Maxwell's Equations

The derivation for electromagnetism follows from identifying the physical fields with the torsional modes of the substrate. The primary challenge is to construct an antisymmetric field strength tensor and its associated 4-vector potential from the dynamics of the fundamental *symmetric* tensor  $\mathcal{G}_{\mu\nu}$ .

*a. Field Identification via Geometric Construction.* To construct an effective  $U(1)$  gauge potential  $A_\mu$  from the underlying symmetric tensor  $\mathcal{G}_{\mu\nu}$ , we require a construction that satisfies several key principles. The resulting 1-form must: (i) be built from the covariant derivatives of the fundamental field, as gauge potentials are related to field gradients; (ii) isolate the purely torsional (vector) modes of the substrate's deformation; and (iii) respect Lorentz covariance. The following form is the simplest and most direct construction that satisfies these physical requirements.

$$A_\mu(x) \equiv C_{EM} u^\alpha P_{\mu\alpha}^{(EM)\beta\gamma} \nabla_\beta \mathcal{G}_{\gamma\delta} u^\delta \quad (62)$$

where  $u^\mu$  is the substrate's local rest 4-velocity,  $C_{EM}$  is a normalization constant, and  $P_{\mu\alpha}^{(EM)\beta\gamma}$  is a symmetry-respecting projector that selects the appropriate torsional (vector) modes of the deformation gradient.<sup>4</sup> The physical, gauge-invariant electromagnetic field strength tensor  $F_{\mu\nu}$  is then naturally defined as the exterior derivative of this emergent potential:

$$F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu. \quad (63)$$

This construction provides the natural bridge from the symmetric tensor  $\mathcal{G}_{\mu\nu}$  to the antisymmetric field strength  $F_{\mu\nu}$ . It guarantees the two source-free Maxwell equations ( $\nabla \cdot \vec{B} = 0$  and  $\nabla \times \vec{E} + \partial \vec{B} / \partial t = 0$ ) through the Bianchi identity  $\partial_{[\lambda} F_{\mu\nu]} \equiv 0$ , while the invariance of  $F_{\mu\nu}$  under the transformation  $A_\mu \rightarrow A_\mu + \partial_\mu \lambda$  ensures the emergence of a local  $U(1)$  gauge symmetry.

*b. Derivation of the Field Equations and Coefficient Matching.* Having constructed the field  $F_{\mu\nu}$  from first principles, gauge and Lorentz invariance uniquely fix the lowest-order effective Lagrangian density for the torsional sector to be quadratic in  $F$ :

$$S_{EM}[A; g] = \int d^4x \sqrt{-g} \left( -\frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} - A_\mu J^\mu \right). \quad (64)$$

The coefficient  $\mu_0$  is not a new fundamental constant, but an effective parameter emerging from the substrate's stiffness  $\kappa$  in the torsional sector; its precise relationship will be derived in Part IV. The term  $F_{\mu\nu} F^{\mu\nu}$  is the only parity-even, gauge- and Lorentz-invariant scalar at two derivatives. Varying Eq. (64) with respect to  $A_\nu$  yields the inhomogeneous Maxwell equations:

$$\nabla_\mu F^{\mu\nu} = \mu_0 J^\nu. \quad (65)$$

In the static, weak-field limit, this becomes  $\nabla^2 \Phi = -\rho/\epsilon_0$ , which fixes the normalization  $\epsilon_0 \mu_0 = 1/c^2$  and recovers Coulomb's law. Thus, the full set of Maxwell equations is recovered.

<sup>4</sup> While other, more complex constructions might be possible, this form stands as the most natural and parsimonious expression for an emergent gauge potential. Its validity is ultimately confirmed by its success in reproducing the full structure of Maxwell's equations and the subsequent consistency of the entire network of derived constants.

*c. Consistency Check: Symmetries and Conservation.* The emergent  $U(1)$  gauge symmetry of the action implies, via Noether's theorem, the conservation of the electric current,  $\nabla_\mu J^\mu = 0$ . The structure is thus fully self-consistent. The ability to choose a gauge, such as the Lorenz gauge condition ( $\partial_\mu A^\mu = 0$ ), to simplify calculations without affecting the physical content is preserved, ensuring full compatibility with the standard formalism of electrodynamics.

## C. Conceptual Closure

The derivations above establish a crucial bridge between the microscopic dynamics of the QEG substrate and the macroscopic language of classical field theory. We have demonstrated that the *mathematical structures* of General Relativity and Electromagnetism—specifically, a symmetric rank-2 tensor field for gravity and an antisymmetric rank-2 field strength for electromagnetism—emerge as inevitable descriptions of the substrate's compressive and torsional modes, respectively.

However, a full demonstration of emergence requires closing the logical loop by showing that the *coupling constants* ( $G, \mu_0$ ) are not free parameters to be fitted to experiment, but are themselves determined by the fundamental properties of the substrate (stiffness  $\kappa$  and dissipation  $\alpha$ ). This section has established the *form* of the laws; the subsequent Parts of this work are dedicated to deriving their *substance*—the values and interrelations of the constants themselves. This will complete the unification, showing that both the field equations and the constants that populate them arise from the same, single underlying reality.

## IX. UNIVERSAL SELF-INTERACTION AND THE DUALITY OF COUPLINGS

A cornerstone of this framework is that sources are stable, extended deformations of the substrate. To understand their properties, we must analyze their self-energy—the energy required to sustain such a deformation. We will demonstrate that this energy is not a uniquely defined quantity, but can be calculated in two physically distinct ways. This duality in the definition of self-energy, as we will show, is the fundamental origin of a scale-dependent effective coupling constant for all physical interactions.

### A. The duality of Self-Interaction Energy

The principle of superposition dictates that the total energy of a system of deformations is the sum of the energies from the mutual interaction of each pair. For a distribution of a modal source specified by a density  $\rho_A(\mathbf{x})$ , the total self-interaction energy  $U_A$  is given by the integral over all pairs of points in the distribution:

$$U_A[\rho_A] = \frac{1}{2} \int d^3x d^3y \rho_A(\mathbf{x}) \mathcal{K}_A(\mathbf{x}, \mathbf{y}) \rho_A(\mathbf{y}) \quad (66)$$

where the kernel  $\mathcal{K}_A(\mathbf{x}, \mathbf{y}) = \lambda_A G_A(\mathbf{x}, \mathbf{y})$  contains the mode-dependent coupling strength  $\lambda_A$  and the Green's function  $G_A$  for the interaction (e.g.,  $1/|\mathbf{x} - \mathbf{y}|$  for long-range forces). This general form can be evaluated through different physical procedures, which reveal its inherent duality:

a. 1. *Global Self-Energy ( $U_{\text{glob}}$ ): The Work of Formation.* The first method calculates the total energy by computing the work done to assemble the deformation from its infinitesimal constituents brought from infinity. This represents the *total work of formation*, including all internal binding energy.

Let us imagine assembling a spherical deformation of final radius  $a$  and uniform source density  $\rho_A$  by building it up in successive thin spherical layers, as illustrated in [42]. At any intermediate stage, the sphere has a radius  $r$  and contains a total modal source  $Q_A(r)$ . The potential at its surface is  $\Phi_A(r) = \lambda_A Q_A(r)/r$ . The work  $dU_A$  required to bring an additional infinitesimal layer of source  $dQ_A$  from infinity to the surface at radius  $r$  is:

$$dU_A = \Phi_A(r) dQ_A = \left( \frac{\lambda_A Q_A(r)}{r} \right) dQ_A \quad (67)$$

For a uniform density  $\rho_A$ , the source contained within radius  $r$  is  $Q_A(r) = \rho_A \cdot (\frac{4}{3}\pi r^3)$ . The source in the next layer of thickness  $dr$  is  $dQ_A = \rho_A \cdot (4\pi r^2 dr)$ . Substituting these into the expression for  $dU_A$ :

$$dU_A = \frac{\lambda_A}{r} \left( \frac{4}{3}\pi \rho_A r^3 \right) (4\pi \rho_A r^2 dr) = \frac{16\pi^2 \lambda_A \rho_A^2}{3} r^4 dr \quad (68)$$

The total energy required to assemble the full sphere is the integral of  $dU_A$  from  $r = 0$  to  $r = a$ :

$$U_{\text{glob}} = \int_0^a dU_A = \frac{16\pi^2 \lambda_A \rho_A^2}{3} \int_0^a r^4 dr = \frac{16\pi^2 \lambda_A \rho_A^2 a^5}{15} \quad (69)$$

Expressing this in terms of the total source  $Q_A = \rho_A \cdot (\frac{4}{3}\pi a^3)$ , we arrive at the definitive expression for the Global Self-Energy:

$$U_{\text{glob}} = \frac{3}{5} \frac{\lambda_A Q_A^2}{a} \quad (70)$$

The geometric factor  $C_g = 3/5$  is the universal signature of the total formation energy of a uniform, volume-filling deformation.

b. 2. *Local Interaction Energy ( $U_{\text{loc}}$ ): The External Field Energy.* The second method calculates the energy by considering the source to be fully assembled and residing on a conducting surface, as in a capacitor. This procedure calculates the energy stored exclusively in the field *external* to the source's boundary, representing the energy available for interaction with other distant sources [42].

The energy  $U_A$  required to place a source  $Q_A$  on a configuration with "modal capacitance"  $C_A$  is given by:

$$U_A = \frac{1}{2} \frac{Q_A^2}{C_A} \quad (71)$$

For a spherical shell of radius  $a$ , which models a source distribution with no internal structure, the modal capacitance is  $C_A = a/\lambda_A$ . Substituting this yields the energy stored in the external field:

$$U_{\text{loc}} = \frac{1}{2} \frac{Q_A^2}{a/\lambda_A} = \frac{1}{2} \frac{\lambda_A Q_A^2}{a} \quad (72)$$

Here, the geometric factor is  $C_g = 1/2$ . This Local Interaction Energy correctly describes the energy of a thin spherical shell source and represents the quasi-linear regime where internal, non-linear binding energies are excluded.

## B. Consequence: The Duality of Effective Couplings

The rigorous derivation above shows that the self-energy of a deformation is not single-valued, but depends on the physical question being asked: are we calculating the total work of formation ( $U_{\text{glob}}$ ) or the energy available for external interaction ( $U_{\text{loc}}$ )? *The difference is captured by a purely geometric factor.*

This duality in energy implies an unavoidable duality in the effective coupling constant,  $\lambda_A$ . We can therefore define two distinct couplings, whose applicability depends on the physical regime:

- **The Local Coupling ( $\lambda_{\text{loc}}$ ):** This coupling describes systems dominated by non-linear self-interaction, such as dense, clumpy structures. The relevant energy is the total work of formation,  $U_{\text{glob}}$ . It is called "global" energy because the calculation integrates over the *entire global volume of the source itself*, a procedure physically relevant for understanding a **local** object's internal structure. It is therefore associated with the geometric factor  $C_g = 3/5$ .
- **The Global Coupling ( $\lambda_{\text{glob}}$ ):** This coupling describes systems in the quasi-linear regime, such as the homogeneous universe on **global** scales. The relevant energy is that of the external field,  $U_{\text{loc}}$ , where internal non-linearities are averaged out. It is called "local" energy because the calculation considers only the field from the *local boundary of the source outwards*. It is therefore associated with the geometric factor for a surface-like source,  $C_g = 1/2$ .

This section has thus established, from a rigorous generalization of classical field theory principles, a necessary duality of effective couplings. This provides the fundamental basis for the scale-dependent nature of gravity, which will be explicitly developed and applied in Sec. XXVIII.

## Part IV: From geometro-elastic properties to Fundamental Constants of nature

To connect our previous abstract framework to physics as observed, we must now explore how the substrate's structure manifests itself in concrete, measurable constants. The transition requires a new step: moving from the *structural level* of laws and symmetries to the *modal level*, where specific excitations of the medium give rise to quantized actions and effective responses.

This Part is devoted to that bridge. We show that once the substrate is treated as a network of oscillators, its modal excitations naturally yield:

1. The geometric nature of action, establishing  $\hbar$  and other quanta as area-like invariants of space-time.
2. Electromagnetic and thermo-entropic modal actions, which connect directly to the observed values of  $\hbar$  and  $\Lambda$ .
3. The reinterpretation of  $\mu_0$ ,  $K_e$ , and  $k_B$  as effective electromotive responses, i.e. modal resonances of the same elastic substrate under different scaling regimes.
4. The emergence of the fine-structure constant  $\alpha$  as a universal damping ratio, and its dual role in both bosonic propagation and fermionic dressing.

5. A synthesis equation linking  $e$ ,  $k_B$ ,  $\mu_0$ ,  $\alpha$ , and  $h$ , which condenses the thermal, electromagnetic, and quantum domains into a single relation.

The guiding principle of this Part is the same as in the previous one: *nothing is postulated that is not required by the substrate's geometry and dynamical consistency*. The constants of physics emerge here as inevitable modal invariants of a single oscillatory medium.

In the next sections we develop these results step by step, beginning with the geometric interpretation of action itself.

## X. THE GEOMETRIC NATURE OF ACTION

The first step in the modal analysis is to determine the dimensional and geometric meaning of the action,  $S$ . In conventional physics,  $S$  is defined as the integral of the Lagrangian  $L$  over time, or equivalently, of the Lagrangian density  $\mathcal{L}$  over spacetime. In our framework of dimensional unification, this definition acquires a precise geometric interpretation.

### A. Action as a Geometric Deformation Area

By construction, the action carries the dimensions of energy  $\times$  time:

$$[S] = [E] \cdot [T]. \quad (73)$$

Within the unified dimensional framework,  $[E] \equiv [L]$  and  $[T] \equiv [L]$ , so

$$[S] = [L] \cdot [L] = [L^2]. \quad (74)$$

Thus, the action is not an abstract bookkeeping device but a measure of *spacetime area*. Every physical process corresponds to the sweeping out of a geometric deformation area within the substrate.

### B. The Dimension of the Lagrangian Density

The action is defined as the integral of the Lagrangian density over spacetime:

$$S = \int \mathcal{L} d^4x. \quad (75)$$

Since  $[S] = [L^2]$  and  $[d^4x] = [L^4]$ , it follows that

$$[\mathcal{L}] = \frac{[S]}{[d^4x]} = \frac{[L^2]}{[L^4]} = [L^{-2}]. \quad (76)$$

This is a profound and unavoidable result: *any* consistent Lagrangian density in this framework must have the dimensions of inverse area. Crucially,  $[L^{-2}]$  is precisely the dimensional signature of Gaussian or Ricci scalar curvature, which means that the identification  $\mathcal{L} \propto R$  in the Einstein–Hilbert action is not arbitrary but a necessary consequence of dimensional consistency within a geometro-elastic substrate.

## C. Interpretation

The equivalence

$$[\mathcal{L}] \equiv [R] \quad (77)$$

shows that the Einstein–Hilbert Lagrangian,  $\mathcal{L}_{EH} \propto R$ , is not an empirical postulate but an intrinsic geometric requirement. This elevates the principle of least action into a *principle of extremal geometry*: the dynamics of fields and particles emerge from spacetime's intrinsic tendency to minimize integrated curvature. In this view, the action is literally the spacetime area swept out by a deformation, and the Lagrangian density is the local curvature cost of that deformation.

## D. Normalization and the Fiducial Scale

As already advanced in Sec. IV, while the theory is intrinsically scale-free, connecting it to measurable quantities requires the introduction of a fiducial reference scale. This is a standard normalization procedure that anchors the abstract geometry of the theory to the operational units used in empirical physics, such as the SI system.

Recalling from Sec. IV that all physical magnitudes collapse to a single geometric unit  $[L]$ , the choice of a reference length, time, or mass is formally equivalent. We therefore adopt the meter as our fiducial unit to maintain consistency with established conventions—any other fiducial choice (second, kilogram, etc.) would yield an equivalent identity by the Spacetime Equivalence Principle—. Therefore, *from this point forward, the fiducial unit of one meter,  $L_{ref} = 1 \text{ m}$ —or its equivalents as defined by the conversion scheme in Sec. IV—will serve as the baseline scale for all subsequent derivations of physical constants.*

## E. The baseline Lagrangian density

We introduce the baseline Lagrangian density as the curvature associated with the unit area:

$$\mathcal{L}_{\text{base}} := \frac{1}{(1 \text{ m})^2}. \quad (78)$$

Integrating this density over a unit 4-volume,  $d^4x = (1 \text{ m})^4$ , yields the fiducial action  $S_{\text{base}}$ :

$$S_{\text{base}} = \int \mathcal{L}_{\text{base}} d^4x = \frac{1}{(1 \text{ m})^2} \cdot (1 \text{ m})^4 = 1 \text{ m}^2. \quad (79)$$

This normalization forces a consistency condition between the geometric and dynamical descriptions of action. The Spacetime Equivalence Principle requires this geometric area to be equivalent to the standard unit of physical action, leading to the identity:

$$1 \text{ m}^2 \equiv 1 \text{ J} \cdot \text{s},$$

This is not a new physical law, but the bridge that ensures the theory's predictions are numerically consistent with experimental reality. It confirms that the energy needed to sustain a deformation over time physically corresponds to a literal area swept out in the spacetime substrate.

Having established the geometric meaning of action and anchored it to a measurable scale, we can

now address its quantization. In the next section, we will show that the Planck constant,  $\hbar$ , emerges as the *fundamental quantum of this spacetime area*, unifying the continuous geometry of the substrate with the discrete nature of quantum mechanics.

## XI. THE QUANTIZATION OF ACTION AND THE EMERGENCE OF PLANCK'S CONSTANT

Having established that action is geometrically a spacetime area, we must address whether this area is continuous or discrete. The answer is dictated by the physical stability of the substrate. A continuous spectrum of deformations, allowing for arbitrarily small action, would lead to an instability analogous to the ultraviolet catastrophe: excitations could fragment into an infinite number of infinitesimal modes, preventing the formation of the stable, localized structures observed in nature (i.e., particles).

Therefore, *a consistent and stable geometro-elastic substrate requires a fundamental regulator: a minimal, non-zero unit of deformation*. We identify this quantum of spacetime area with Planck's constant,  $\hbar$ .

### A. Planck's Constant as the Minimal Geometric Quantum

The stability of the substrate thus imposes the quantization of action. Any physical action  $S$  must be an integer multiple of this fundamental geometric unit:

$$S = n \hbar, \quad n \in \mathbb{Z}^+, \quad (80)$$

where  $\hbar$  is the elementary quantum of spacetime area. Its dimensions,  $[\hbar] = [S] = [L^2]$ , confirm its nature as a fundamental geometric area, not merely a dynamical parameter.

In this framework,  $\hbar$  is not an external constant imported from quantum mechanics, but an intrinsic property of the geometry of the substrate itself. Just as a crystal lattice enforces a minimum cell size, the finite deformability of the geometro-elastic substrate enforces a minimum unit of action.

### B. Geometric Consequences of Quantization

The identification of  $\hbar$  with a minimal area has three immediate consequences:

1. **Resolution of the continuity/discreteness tension:** Quantum discreteness is the tessellation of the continuous substrate into fundamental patches of area  $\hbar$ .
2. **Universality:** Every physical process, regardless of its nature, must be built from integer multiples of the same minimal geometric unit of deformation.
3. **Necessity of quantization:** The discrete spectra of quantum mechanics are not axioms but inevitable consequences of a stable substrate with finite deformability.

### C. Conclusion

We conclude that Planck's constant is the minimal area of spacetime deformation permitted by the stable, geometro-elastic substrate. This result completes the bridge between classical geometry and quantum discreteness. The next logical step is to derive the numerical value of this fundamental area from the substrate's properties. In the following sections, we will show how the interplay between the substrate's modal dynamics, its principle of reciprocity, and its intrinsic dissipation determines the precise value of  $\hbar$ .

## XII. MODAL SCALES AND THE ELECTROMAGNETIC QUANTUM OF AREA

The unified substrate supports distinct orthogonal modes (compressive, torsional, tensorial), all governed by the same geometro-elastic law but with different modal responses. While unified in origin, each mode is characterized by a natural *length scale* at which its dynamics become dominant. In this section, we *derive*—rather than postulate—the characteristic length of the torsional (electromagnetic) mode directly from the constitutive properties of the vacuum, and use it to obtain the electromagnetic quantum of area,  $\hbar$ .

### A. The characteristic length of the electromagnetic mode

The vacuum's electromagnetic response is encoded by its constitutive parameters,  $(\varepsilon_0, \mu_0)$ . We now construct the *unique* length scale compatible with the following principles:

1. **Scale covariance:** All modal scales must ultimately be built from the fiducial reference length,  $L_{\text{ref}}$ , introduced in Sec. XD, without introducing independent dimensional parameters.
2. **Lorentz and Duality Covariance:** The scale must be invariant under Lorentz transformations and electromagnetic duality rotations. This restricts its dependence on the constitutive parameters to the unique invariant combination  $\varepsilon_0 \mu_0 = \frac{1}{c^2}$ .<sup>5</sup>
3. **Minimality / structural simplicity:** In the absence of further physical principles, the relationship must be the simplest possible function consistent with the symmetries. This avoids introducing unmotivated complexity into the framework.

These requirements uniquely determine the form of  $\ell_{\text{EM}}$ . By (i), any candidate must be of the form

$$\ell_{\text{EM}} = L_{\text{ref}} f(\varepsilon_0, \mu_0).$$

By (ii),  $f$  can only depend on the duality-symmetric, Lorentz-invariant product  $\varepsilon_0 \mu_0$ ; by (iii), the simplest non-trivial relationship is linear:

$$\ell_{\text{EM}} = L_{\text{ref}} (\varepsilon_0 \mu_0). \quad (81)$$

<sup>5</sup> The vacuum is symmetric under duality rotations that mix electric and magnetic fields. This symmetry is broken by quantities like the impedance  $Z_0 = \sqrt{\mu_0/\varepsilon_0}$  which cannot, therefore, define a fundamental geometric scale of the vacuum itself.

Using the vacuum identity  $\varepsilon_0\mu_0 = 1/c^2$  and the SI-anchoring  $L_{\text{ref}} = 1\text{ m}$  of Sec. X, we obtain the *electromagnetic modal length*:

$$\ell_{\text{EM}} = \frac{1\text{ m}}{c^2}. \quad (82)$$

This result is not an ansatz, but the only possible construction compatible with the axiomatic foundations of the theory.

### B. Derivation of the electromagnetic quantum of area $\hbar$

Since  $[S] = [L^2]$ , the action carried by a mode of characteristic length  $\ell$  must scale as  $\ell^2$ . More precisely, by homogeneity and the fiduciary normalization of Sec. X,

$$S_{\text{mode}}(\ell) = C \left( \frac{\ell}{L_{\text{ref}}} \right)^2 S_{\text{base}} \quad \text{with} \quad S_{\text{base}} = L_{\text{ref}}^2. \quad (83)$$

Requiring continuity to the baseline configuration when  $\ell = L_{\text{ref}}$ , i.e.  $S_{\text{mode}}(L_{\text{ref}}) = S_{\text{base}}$ , fixes  $C = 1$ . Hence

$$S_{\text{mode}}(\ell) = \ell^2. \quad (84)$$

Applying (84) to the torsional (electromagnetic) mode and substituting the modal length derived in Eq. (82),

$$\ell_{\text{EM}} = \frac{1\text{ m}}{c^2},$$

we obtain the first-order electromagnetic quantum of area, which we identify with the reduced Planck constant:

$$\hbar \equiv S_{\text{EM}} = (\ell_{\text{EM}})^2 = \left( \frac{1\text{ m}}{c^2} \right)^2 = \frac{1\text{ m}^2}{c^4}. \quad (85)$$

Using the fiduciary identity  $1\text{ m}^2 \equiv 1\text{ J s}$  (Sec. X), Eq. (85) yields  $\hbar \approx 1.23 \times 10^{-34}\text{ J s}$ , in leading-order agreement with the measured value  $\hbar_{\text{exp}} = 1.054 \times 10^{-34}\text{ J s}$ .

*On higher-order corrections.* The residual discrepancy is expected: the torsional mode is not perfectly elastic. As shown in Secs. VII and IX, dissipation and self-interaction dress the first-order result by a universal, dimensionless factor controlled by  $\alpha$ , yielding

$$\hbar = \frac{1\text{ m}^2}{c^4} \Xi_{\text{EM}}(\alpha), \quad \Xi_{\text{EM}}(\alpha) = 1 + \mathcal{O}(\alpha),$$

with the leading radiative dressing consistent with the vacuum's quality factor and one-loop-like form factors discussed later. Thus  $\hbar$  emerges *predictively* from the substrate's modal scale (82) together with the dissipative structuring of Sec. IX.

## XIII. THE GEOMETRIC ORIGIN OF CHARGE AND ELECTROMAGNETIC STIFFNESS

In Sec. XII we derived the characteristic length of the torsional (electromagnetic) mode,

$$\ell_{\text{EM}} = \frac{1\text{ m}}{c^2},$$

from axiomatic principles of scale covariance, Lorentz/duality covariance, and structural simplicity. We now show that this length is not an auxiliary construct but the very *structural unit of electric charge*. This identification not only grounds charge geometrically but also reveals that the electromagnetic stiffness of the substrate is encoded in  $c^2$ .

### A. The Structural Charge as a Geometric Length

A central conclusion of the unified framework is the *endogeneity of sources*: mass, charge, and entropy are not external inputs but stable excitations of the substrate itself (Sec. II A). Accordingly, the torsional deformation supported by  $\ell_{\text{EM}}$  must manifest as the minimal quantum of charge. We therefore identify this *structural charge*,  $q_s$ , with the modal length:

$$q_s \equiv \ell_{\text{EM}} = \frac{1}{c^2}\text{ m}, \quad (86)$$

This reinterprets the fundamental unit of charge as a geometric length scale intrinsic to the substrate.

### B. The Universal Elastic Law and Electromagnetic Stiffness

This identification can be validated against the universal constitutive relation established in Sec. V,

$$Q = k S.$$

For the electromagnetic mode,

$$q_s = k_{\text{EM}} S_{\text{EM}}, \quad (87)$$

with  $S_{\text{EM}} = \hbar$  the modal action. Substituting the previously derived values yields

$$k_{\text{EM}} = \frac{q_s}{S_{\text{EM}}} = \frac{\frac{1\text{ m}}{c^2}}{\frac{1\text{ m}^2}{c^4}} = \frac{c^2}{1\text{ m}}. \quad (88)$$

This stiffness is not a new fundamental constant but the speed of light squared, normalized by the fiducial meter introduced in Sec. X D. In other words,

$$q_s = \frac{c^2}{1\text{ m}} \cdot \hbar. \quad (89)$$

*Physical interpretation.* This law stands in precise analogy with Einstein's mass-energy equivalence:

$$\begin{aligned} E &= m c^2 && \text{(compressive mode, gravitational)} \\ q_s &= \frac{\hbar}{1\text{ m}} c^2 && \text{(torsional mode, electromagnetic)}. \end{aligned} \quad (90)$$

Whereas  $E = m c^2$  interprets mass as “frozen energy,” the above relation interprets charge as sourcing a deformation whose action is proportional to  $c^2$ . In both,  $c^2$  functions as a *universal elastic modulus of space-time*, relating substrate latent content (energy or action density) to observable sources (mass or charge).

### C. Conclusion

We conclude that the elementary electric charge is the quantized torsional source of the substrate, arising inevitably from the modal scale  $\ell_{\text{EM}}$  and the universal constitutive law  $Q = kS$ . Its stiffness is encoded in  $c^2$ , in full analogy with the role of  $c^2$  in mass–energy equivalence.

The physically observed elementary charge,  $e$ , arises when this bare excitation is *dressed* by the dissipative and self-interaction effects of the substrate. As will be formalized in later sections, these corrections are governed universally by the fine–structure constant  $\alpha$ .

## XIV. THE THERMO-ENTROPIC MODAL SCALE AND THE COSMOLOGICAL ACTION

Having analyzed the torsional (electromagnetic) mode, we now turn to the compressive/tensorial modes of the substrate, associated with large-scale, isotropic phenomena such as thermodynamics and cosmology. Our aim is to derive the natural length scale of this mode,  $\ell_{\text{th}}$ , and the corresponding action it supports, revealing a direct link between the quantum-elastic structure of the vacuum and the cosmological constant.

### A. The Thermo-Entropic Length Scale

The thermo-entropic mode represents the lowest-energy, largest-scale collective excitations of the substrate. Its characteristic length scale cannot be a new fundamental parameter but must emerge from the intrinsic properties of the framework already established:

1. **Quantum Discreteness:** encoded by  $\hbar$ , the minimal unit of action/area.
2. **Relativistic Structure:** encoded by  $c$ , which governs causal propagation.
3. **Reference Normalization:** the fiducial length  $L_{\text{ref}}$  XD, which anchors all scales.

By structural necessity, the only non-trivial length that can be constructed from these three ingredients is the **quantum-relativistic reference length**,  $L_q$ :

$$L_q = \frac{\hbar c}{L_{\text{ref}}}. \quad (91)$$

For the lowest-energy mode of the substrate, there is no other available scale to define its characteristic length. We therefore must identify  $\ell_{\text{th}}$  with this unique quantum-relativistic construct:

$$\ell_{\text{th}} \equiv L_q = \frac{\hbar c}{1 \text{ m}}. \quad (92)$$

Substituting the first-order expression for  $\hbar$  from Eq. (85),  $\hbar = 1 \text{ m}^2/c^4$ , yields:

$$\ell_{\text{th}} = \frac{(1 \text{ m}^2/c^4) c}{1 \text{ m}} = \frac{1 \text{ m}}{c^3}. \quad (93)$$

Thus, the characteristic length of the thermo-entropic mode is uniquely determined by the theory’s structure:

$$\boxed{\ell_{\text{th}} = \frac{1 \text{ m}}{c^3}} \quad (94)$$

### B. The Thermo-Entropic Modal Action

As established by the universal scaling law  $S_{\text{mode}}(\ell) = \ell^2$ , the action of the thermo-entropic mode is given by:

$$S_{\text{th}} = (\ell_{\text{th}})^2 = \left(\frac{1 \text{ m}}{c^3}\right)^2 = \frac{1 \text{ m}^2}{c^6}. \quad (95)$$

With the fiducial identification  $1 \text{ m}^2 \equiv 1 \text{ J} \cdot \text{s}$ , this corresponds to the value:

$$\boxed{S_{\text{th}} = \frac{1}{c^6} \text{ J} \cdot \text{s}} \approx 1.37 \times 10^{-51} \text{ J} \cdot \text{s}. \quad (96)$$

### C. Interpretation

This quantity represents the *minimal elastic action* associated with a global, isotropic deformation of the substrate. Its extreme smallness is a direct consequence of the modal hierarchy, where compressive/tensorial modes are suppressed by higher powers of  $c$ . Subsequent sections will demonstrate that this residual action density is naturally identified with the observed cosmological constant  $\Lambda$ , thereby linking quantum discreteness, relativistic structure, and cosmic acceleration within a single geometro-elastic framework.

#### *Conclusion on Modal Actions*

We have demonstrated how the actions associated with the primary modes of the substrate emerge not as empirical inputs, but as geometrically quantized projections of a single underlying vacuum structure. The baseline action ( $S_{\text{base}}$ ), the electromagnetic quantum of action ( $\hbar$ ), and the thermo-entropic action ( $S_{\text{th}}$ ) are not independent parameters but represent the scaled geometric integrals of the same deformable substrate. These results strongly suggest that these modal actions can be interpreted as effective **Noether invariants**, each arising from a specific symmetry projection of the unified geometro-elastic dynamics. In this view, what we traditionally call fundamental constants are, in fact, the geometric signatures of the vacuum’s quantized deformability.

## XV. MODAL LAGRANGIAN DENSITY VERSUS EFFECTIVE ENERGY DENSITY

Given the Spacetime Equivalence Principle and the identification of action as an area,  $[S] = [L^2]$ , it becomes inevitable to distinguish between the *modal Lagrangian density* ( $\mathcal{L}$ ), associated with a single coherent excitation of the substrate, and the *effective vacuum energy density* ( $\rho_{\text{vac}}$ ), which appears as a macroscopic observable after coarse-graining over incoherent modes. Specifically:

- **The Lagrangian Density** ( $\mathcal{L}$ ) describes the dynamics of a single, coherent, fundamental mode of the vacuum oscillator lattice. It is a theoretical, microscopic quantity. Dimensional consistency and the fiduciary normalization  $1 \text{ m}^2 \equiv 1 \text{ J} \cdot \text{s}$  require that the fundamental modal action density takes the form

$$\mathcal{L}_{\text{modal}} = \frac{\hbar c}{1 \text{ m}^4}$$

- **The Measured Energy Density** ( $\rho_{\text{vac}}$ ) is a macroscopic, cosmological observable. It reflects the net effect of an immense number of uncoordinated vacuum oscillators, with their phases and spatial orientations being statistically random. At the macroscopic level, statistical isotropy enforces a coarse-grained average over random phases, entirely analogous to the emergence of  $g_{\mu\nu} = \langle G_{\mu\nu} \rangle$ . The observable energy density is therefore

$$\rho_{\text{eff}} \equiv \frac{1}{2\pi} \int_0^{2\pi} \mathcal{L}_{\text{modal}} d\theta = \frac{\mathcal{L}_{\text{modal}}}{2\pi} = \frac{\hbar c}{2\pi \cdot 1 m^4}$$

Finally, note that:

- By substituting  $k$  with the quantum of angular frequency of the electromagnetic mode  $\frac{c}{\lambda}$ , and Planck's constant  $h$  for the quantum of action, we obtain a quantum expression of mass-energy using 37:

$$m = \frac{\hbar c}{\lambda} \quad (97)$$

This equation directly links the energy of photons (or other quantum excitations) to mass, reinforcing Einstein's mass-energy equivalence from a fundamentally new perspective. Substituting  $\lambda = 1 m$  in (97), corresponding to the characteristic scale of the unit quantum oscillator in our framework, we obtain the quantum of mass-energy for the electromagnetic field  $m = \frac{\hbar c}{1 m}$ .

- Dividing this quantum mass-energy by a volume  $V = 1 m^3$ , and considering the linear momentum  $\hbar = \frac{h}{2\pi}$  we obtain a quantum of mass density  $\rho_{\text{vac}} = \frac{\hbar c}{1 m^4}$  which corresponds with (XV). When transitioning from the description of a single, coherent angular mode to a macroscopic, isotropic average over all possible phases or solid angles (XV), we get  $\rho_{\text{eff}} \equiv \frac{\hbar c}{2\pi \cdot 1 m^4} \approx 5.03 \times 10^{-27} \text{ kg m}^{-3}$  which retrospectively matches the Planck satellite measurements, thereby turning the concordance into a non-trivial prediction of the model rather than an adjustment. [43].

## XVI. MODAL DYNAMICS AND THE EMERGENCE OF COUPLING CONSTANTS

Having established the characteristic scales of the substrate's modes, we now derive the coupling constants governing their dynamics. We will show that the magnetic permeability ( $\mu_0$ ), Coulomb's constant ( $K_e$ ), and Boltzmann's constant ( $k_B$ ) are not independent parameters. They emerge as projections of a single, universal inductive response law, evaluated at the characteristic velocity of each mode. To do so, we first establish the unique scaling of these velocities.

### A. Modal Velocities and the Uniqueness of Scaling

In our framework, current is a dimensionless measure of deformation flow,  $[I] = 1$ . The physical distinction between modes arises from their characteristic propagation velocity,  $v$ . These velocities are not arbitrary but are uniquely determined by the theory's axiomatic principles:

1. **Fiducial Baseline:** The reference mode is defined by the fiducial scales, corresponding to a normalized velocity  $v_{\text{base}} = 1$ .
2. **Relativistic Limit:** Lorentz invariance introduces a single invariant speed,  $c$ . The fastest, torsional (electromagnetic) excitations must propagate at this speed,  $v_{\text{EM}} = c$ .
3. **Modal Reciprocity:** The principle of reciprocity requires that the highly stiff, fast torsional mode ( $v \sim c$ ) be balanced by a highly compliant, slow compressive/torsional (thermo-entropic) mode. The only non-trivial scaling inverse to  $c$  is  $1/c$ . Thus,  $v_{\text{th}} = 1/c$ .

These three velocities,

$$\{1, c, 1/c\},$$

are therefore the only ones consistent with the geometro-elastic substrate. Any alternative assignment would violate either Lorentz invariance, reciprocity, or the fiducial normalization.

### B. The Universal Law of Inductive Response

The substrate's dynamic response to a changing deformation flow is governed by a universal inductive law, analogous to Faraday's law:

$$\mathcal{E} = -L_{\text{vac}} \frac{dI}{dt}, \quad (98)$$

where  $\mathcal{E}$  is the resulting modal potential and  $L_{\text{vac}} = \mu_0 L_{\text{ref}}$  is the fiducial inductance. The term  $dI/dt$  represents the characteristic rate of change, or acceleration, of the modal deformation.

The characteristic rate of change is the ratio of the characteristic velocity to the characteristic timescale:  $dI/dt \sim v_{\text{mode}}/\tau_{\text{mode}}$ . The timescale of a mode is inversely proportional to its velocity (high-velocity modes have short timescales, slow modes have long ones), so  $\tau_{\text{mode}} \sim 1/v_{\text{mode}}$ . This leads to the universal scaling rule:

$$\frac{dI}{dt} \sim \frac{v_{\text{mode}}}{1/v_{\text{mode}}} = (v_{\text{mode}})^2. \quad (99)$$

Hence, the modal potential response is proportional to the square of the characteristic velocity:

$$\boxed{\mathcal{E}_{\text{mode}} \propto \mu_0 (v_{\text{mode}})^2}. \quad (100)$$

This law is covariant, dimensionally consistent, and introduces no new parameters: all modal couplings must be determined by  $\mu_0$  and the velocity hierarchy.

### C. Derivation of the Coupling Constants

Applying the scaling law (100) to the three modal velocities yields the coupling constants directly.

1. **Baseline Potential ( $\mu_0$ ):** For the reference mode,  $v_{\text{mode}} = 1$ . The response is the baseline potential:

$$\mathcal{E}_{\text{base}} \equiv \mu_0, \quad (101)$$

the intrinsic inertial potential of the substrate in the quasi-static limit.

2. *Electromagnetic Potential ( $K_e$ ):* For the torsional mode,  $v_{\text{mode}} = c$ . The response is scaled by  $c^2$ :

$$\mathcal{E}_{\text{EM}} = \mu_0 c^2 = \frac{1}{\varepsilon_0}. \quad (102)$$

This corresponds, up to the geometric factor  $4\pi$ , to Coulomb's constant  $K_e = 1/4\pi\varepsilon_0$ . Thus,  $K_e$  emerges as the relativistically-scaled inductive potential of the substrate.

3. *Thermo-Entropic Potential ( $k_B$ ):* For the compressive/tensorial mode,  $v_{\text{mode}} = 1/c$ . The response is suppressed by  $1/c^2$ :

$$\mathcal{E}_{\text{th}} = \frac{\mu_0}{c^2}. \quad (103)$$

This value coincides, to leading order, with Boltzmann's constant  $k_B$ . The small deviation observed experimentally is interpreted as the radiative dressing of this response by the universal dissipative invariant  $\alpha$ , consistent with Secs. VII and VII.

#### D. Synthesis and Interpretation

We therefore obtain the unified ladder of coupling constants:

$$\boxed{\{\mu_0, 4\pi K_e, k_B\} \approx \mu_0 \{1, c^2, c^{-2}\} [1 + \mathcal{O}(\alpha)]}. \quad (104)$$

This shows that the constants governing magnetostatics, electrostatics, and thermodynamics are not independent. They are modal projections of the same inductive response of the vacuum, evaluated at the three characteristic velocities allowed by the geometro-elastic substrate.

*Conceptual significance.* This result provides a unified operational principle: *electromotive response emerges from the substrate's geometric resistance to deformation.* What experimental physics has treated as three unrelated constants are revealed here as facets of a single property.  $\mu_0$  sets the baseline inductive stiffness;  $K_e$  is its relativistic amplification at high frequency ( $v = c$ ); and  $k_B$  is its reciprocal attenuation in the slow, dissipative regime ( $v = 1/c$ ). In this sense, magnetostatics, electrostatics, and thermodynamics are unified as different *modal responses of the same geometro-elastic medium.*

### XVII. SYNTHESIS OF CONSTANTS AND THE VACUUM'S CONSTITUTIVE EQUATION

Building on the principles derived in the preceding sections, we now demonstrate their ultimate convergence. We will show that the modal invariants of the substrate are not independent but are locked into a single, self-consistent relationship by the theory's structure. This synthesis is achieved by applying the principles of fermionic dressing and damped equipartition.

#### A. The Electro-Thermal Identity

We first establish a direct link between the elementary charge and the thermal potential of the vacuum.

1. **Principle of Fermionic Dressing:** The observed elementary charge,  $e$ , is the bare structural charge of the substrate,  $q_s$ , dressed by the dissipative response of the vacuum's spin-1/2 modes. As established in Sec. VII C, this is governed by the factor  $g_e \alpha \approx 2\alpha$ :

$$e = q_s \cdot (g_e \alpha). \quad (105)$$

2. **The Bare Charge from Constitutive Properties:** The bare charge  $q_s$  can be constructed independently from the vacuum's constitutive properties. In Sec. XVI, we identified  $\mu_0$  with the substrate's baseline electromagnetic potential. The corresponding fiducial capacitance is  $C_{\text{fid}} = \varepsilon_0 \cdot (1 \text{ m})$ . The bare charge is their product:

$$q_s = \mu_0 \cdot C_{\text{fid}} = \mu_0 \varepsilon_0 \cdot (1 \text{ m}) = \frac{1 \text{ m}}{c^2}. \quad (106)$$

3. **Synthesis:** In Sec. XVI, we also derived the thermo-entropic potential as  $k_B \equiv \mu_0/c^2$ , which implies  $1/c^2 \equiv k_B/\mu_0$ . Substituting this into the expression for  $q_s$  gives  $q_s \approx (k_B \cdot 1 \text{ m})/\mu_0$ . Inserting this into the dressing principle yields:

$$e \equiv \left( \frac{k_B \cdot 1 \text{ m}}{\mu_0} \right) (g_e \alpha) \implies \boxed{\mu_0 \cdot e \equiv k_B \cdot 2\alpha}, \quad (107)$$

where the fiducial meter X D (equivalent to 1 K IV) normalizes the relation. This equation forms the **electro-thermal identity** of the vacuum.

#### B. The Thermo-Quantum Identity

We now establish a key thermodynamic principle of the QEG framework: *the effective thermal energy available to a normal mode of the geometro-elastic substrate is suppressed by its quality factor,  $Q$ .* We demonstrate the robustness of this "Damped Equipartition Principle" by outlining two independent derivations from fundamental statistical mechanics.

*a. 1. Energy Balance Approach.* Consider a single, underdamped normal mode of the substrate modeled as a harmonic oscillator with quality factor  $Q_A = 1/(2\zeta_A)$ , in thermal equilibrium with a bath at temperature  $T$ . In thermal equilibrium (classical limit) a single quadratic oscillator has  $\langle E_{\text{kin}} \rangle = \frac{1}{2} k_B T$ ,  $\langle E_{\text{pot}} \rangle = \frac{1}{2} k_B T$ , hence

$$\langle E_{\text{stored}} \rangle = k_B T. \quad (108)$$

From (108) one gets the energy irreversibly lost to the bath per cycle as:

$$\Delta E_{\text{diss}}^{(\text{cycle})} = \frac{2\pi}{Q_A} k_B T \quad (109)$$

At equilibrium the bath must inject the *same* amount of energy back into the mode over a cycle, by detailed balance. It is convenient to quotient out the trivial  $2\pi$  of phase advance and work *per natural radian* (the natural coarse-graining time in a quasi-monochromatic steady state). We therefore define the *available thermal energy per radian*

$$\boxed{E_{\text{eff}}^{(A)} \equiv \frac{\Delta E_{\text{diss}}^{(\text{cycle})}}{2\pi} = (k_B T) \frac{1}{Q_A}} \quad (110)$$

*b. 2. Fluctuation-Dissipation Approach.* The same result is obtained independently from the Fluctuation-Dissipation Theorem (FDT). Consider a single normal mode  $A$  described (in generalized coordinates) by a damped harmonic oscillator with natural frequency  $\omega_0$ , damping ratio  $\zeta_A \ll 1$ , and stiffness scale  $\kappa_A$  (as in Sec. X). The causal susceptibility (coupling Hamiltonian  $-f(t)x$ ) is

$$\begin{aligned}\chi_A(\omega) &= \frac{1}{\kappa_A(\omega_0^2 - \omega^2 - i2\zeta_A\omega_0\omega)}, \\ \chi_A''(\omega) &= \frac{2\zeta_A\omega_0\omega}{\kappa_A[(\omega_0^2 - \omega^2)^2 + (2\zeta_A\omega_0\omega)^2]}.\end{aligned}\quad (111)$$

The fluctuation-dissipation theorem (classical limit) for our conventions reads

$$\begin{aligned}S_{xx}(\omega) &= \frac{2k_B T}{\omega} \chi_A''(\omega) \\ \langle x^2 \rangle &= \frac{1}{2\pi} \int_0^\infty S_{xx}(\omega) d\omega,\end{aligned}\quad (112)$$

which reproduces  $\langle E_{\text{pot}} \rangle = \frac{1}{2}\kappa_A\omega_0^2\langle x^2 \rangle = \frac{1}{2}k_B T$ . The average power dissipated by the mode due to the thermal fluctuations is

$$P_{\text{diss}} = \int_0^\infty d\omega \omega \chi_A''(\omega) \frac{k_B T}{\pi} = \frac{\omega_0}{Q_A} k_B T, \quad (113)$$

where the last equality follows by evaluating the narrow-band Lorentzian integral using (111) and  $Q_A^{-1} = 2\zeta_A$ . Multiplying by the period  $T_0 = 2\pi/\omega_0$  yields exactly (109), hence dividing by  $2\pi$  gives (110). This completes an independent FDT proof of

$$\boxed{E_{\text{eff}}^{(A)} = (k_B T)/Q_A}$$

*Quantum correction.* Beyond the classical limit the factor  $k_B T$  is replaced by  $\frac{\hbar\omega}{2} \coth(\hbar\omega/2k_B T)$ . All results below persist with this replacement; at the fiducial scale we remain in the classical regime by construction.

#### *Application to the Vacuum and the Thermo-Quantum Identity*

We now apply this principle to the vacuum itself. As established in Sec. VII, processes involving the "dressing" of localized, spin-1/2 excitations (i.e., fermions) are governed by a quality factor determined by the fine-structure constant  $\alpha$  and amplified by the Dirac g-factor,  $g_e = 2$ :

$$Q_{\text{ferm}} = \frac{1}{g_e \alpha} \approx \frac{1}{2\alpha} \quad (114)$$

The effective thermal energy available from the vacuum bath to fuel such a fermionic process is therefore:

$$E_{\text{eff}}^{(\text{ferm})} = (k_B T) \frac{1}{Q_{\text{ferm}}} = (k_B T)(g_e \alpha) \quad (115)$$

A stable, resonant quantum of the substrate (i.e., a particle) at a characteristic wavelength  $\lambda$  requires that this available energy matches the quantum of excitation,  $E_q = hc/\lambda$ . This *energy matching condition*,  $E_{\text{eff}}^{(\text{ferm})} = E_q$ , evaluated at the fiducial scale of the

framework ( $T \equiv 1\text{ K}, \lambda = 1\text{ m}$ ), yields one of the central results of our theory:

$$\boxed{(k_B \cdot 1\text{ K}) \cdot (2\alpha) = \frac{hc}{1\text{ m}}} \quad (116)$$

This equation, which we term the **Thermo-Quantum Identity**, forms a cornerstone of the synthesis of constants developed in Sec. XVII. It connects the thermodynamic properties of the vacuum ( $k_B$ ), its dissipative quantum nature ( $\alpha$ ), and its fundamental quantum of action ( $h$ ).

### C. The Vacuum's Constitutive Equation

The electro-thermal and thermo-quantum identities are two sides of the same coin. Combining them yields a single, powerful synthesis that interlocks the three domains of physics:

$$\boxed{\mu_0 \cdot e \equiv k_B \cdot 2\alpha \equiv \frac{hc}{1\text{ m}}} \quad (117)$$

where the second-order terms have been omitted, and a more accurate approximation would be given by

$$\begin{aligned}\mu_0 \cdot e_q(2\alpha + \frac{\alpha}{2\pi} + \dots) \\ \equiv k_B \cdot 1\text{ K}(2\alpha + \dots) \\ \equiv \frac{hc}{1\text{ m}}(1 + 2\alpha + \dots)\end{aligned}\quad (118)$$

This is the **Vacuum's Constitutive Equation**. It is not a numerical coincidence, but the necessary structural closure condition of the Quantum-Elastic Geometry framework. It reveals the fundamental constants of nature not as arbitrary measured values, but as deeply interconnected parameters whose ratios are fixed by the unified, dissipative, and quantized properties of the spacetime substrate itself.

## XVIII. THE NATURE OF THE GRAVITATIONAL CONSTANT $G$

### A. Static Origin: Deriving $G$ from Modal Reciprocity and Self-Energy

The *Principle of Modal Reciprocity*, established in Sec. VI, posits that the gravitational compliance ( $G$ ) is inversely proportional to the electromagnetic stiffness ( $K_e$ ), scaled by a geometric factor:

$$G = C_{\text{geom}} \cdot \frac{1}{K_e} = C_{\text{geom}} \cdot 4\pi\epsilon_0 \quad (119)$$

Our task is to determine  $C_{\text{geom}}$  from first principles. As anticipated in Sec. VIA, this factor is derived from the geometric integration of self-energy.

In Sec. IX, we rigorously derived the universal geometric factor associated with the self-energy of a dense, volume-filling spherical source. We showed that the total work of formation ( $U_{\text{glob}}$ ) for such a configuration is characterized by a geometric factor of  $C_g = 3/5$ . Since the standard Newtonian constant of gravitation,  $G_N$ , is precisely the coupling that governs these localized, self-interacting systems (e.g., stars

and galaxies), it is a requirement of consistency to identify its geometric coefficient with this value:

$$C_{\text{geom}} = \frac{3}{5} \quad (120)$$

Substituting this directly into the reciprocity principle gives the definitive expression for the standard Newtonian constant of gravitation:

$$G_N \equiv G_{\text{loc}} = \left(\frac{3}{5}\right) 4\pi\epsilon_0 \quad (121)$$

Evaluating with physical constants,

$$G = \left(\frac{3}{5}\right) 4\pi\epsilon_0 \approx 6.676 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2},$$

in excellent agreement with the measured Newtonian constant. The numerical success of this derivation confirms the proposed reciprocity principle: the vacuum's longitudinal compliance (gravitational softness,  $G$ ) is inversely proportional to its transverse stiffness (electromagnetic tension,  $K_e$ ). This reveals a profound symmetry where gravity and electromagnetism emerge as orthogonal projections of the same underlying field elasticity. The empirical disparity between their strengths is not accidental but a structural necessity: the vacuum is extremely compliant to gravitational deformation precisely because it is so immensely rigid electromagnetically. In essence, the vacuum acts as a geometric impedance surface, whose tension and compliance must balance, thereby setting the apparent strengths of the fundamental forces.

## B. Dynamic Origin of $G$ : Gravity as a Quadratically-Damped Vacuum Response

Having established the static interpretation of  $G$  as a compliance modulus of the vacuum, we now turn to its *dynamic origin*. We demonstrate that  $G$  is not a fundamental constant, but an *inevitable, emergent consequence* of the dissipative structure of the quantum-elastic substrate. The derivation follows directly from three principles already established in the QEG framework:

1. The vacuum possesses a universal baseline stiffness, quantified by the magnetic permeability  $\mu_0$ , which sets the primordial interaction scale XVIC.
2. Dissipation is encoded in the *structural damping tensor*  $\zeta_{\mu\nu}$ , with scalar norm  $\|\zeta_{\mu\nu}\| = \alpha$ , the fine-structure constant VII.
3. Gravity couples to the energy-momentum tensor  $T_{\mu\nu}$ , i.e. to quadratic field invariants, and therefore must arise as a *second-order dissipative response*.

*From Primary Stiffness to Effective Coupling.* At the bare level, the vacuum stiffness is  $\mu_0$ . The presence of dissipation filters this bare coupling through a *dissipative transfer function*  $\mathcal{T}(\alpha)$ :

$$G = \mu_0 \cdot \mathcal{T}(\alpha).$$

For first-order processes (e.g. charge dressing),  $\mathcal{T}(\alpha) \propto \alpha$ . However, because gravity couples to  $T_{\mu\nu}$ , its response requires *two interactions* with the dissipative medium (mass deformation  $\rightarrow$  dissipative modulation  $\rightarrow$  induced force). Thus,

$$\mathcal{T}^{(2)}(\alpha) \propto \alpha^2.$$

In the absence of other fundamental geometric or normalization factors, the principle of parsimony dictates that this proportionality is direct, setting the constant to unity in a first-order approximation. Therefore, we have that

$$\mathcal{T}^{(2)}(\alpha) \equiv \alpha^2$$

*Final Constitutive Relation.* This uniquely fixes the effective gravitational coupling as

$$G \equiv \mu_0 \alpha^2 \quad (122)$$

which is not a conjecture but a *necessary consequence* of: (i)  $\mu_0$  as primordial stiffness, (ii)  $\alpha$  as damping norm, and (iii) the quadratic nature of gravitational coupling.

*Tensorial Formulation.* Promoting  $\alpha^2$  to its tensorial origin, we write

$$G g_{\mu\nu} = \mu_0 \zeta_{\mu\alpha} \zeta^{\alpha\nu}, \quad (123)$$

which expresses gravity as the effective longitudinal stiffness generated by the quadratic action of the structural damping tensor. In analogy with elasticity, if  $\mathcal{G}_{\mu\nu}$  is the strain, then  $\zeta_{\mu\nu}$  represents the induced internal stress. Gravity is thus the macroscopic manifestation of the vacuum's dissipative geometry.

*Numerical Validation.* Evaluating with physical constants,

$$G = \mu_0 \alpha^2 \approx (4\pi \times 10^{-7}) \cdot (0.007297)^2 \approx 6.69 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2},$$

in excellent agreement with the measured Newtonian constant.

*Conclusion.* We conclude that  $G$  is not a free parameter, but the *quadratically-damped residual of the electromagnetic substrate stiffness*. This interpretation anchors gravity within the same constitutive law as electromagnetism and thermodynamics, completing the triad of elastic responses of the vacuum.

## C. Unified Interpretation of the Fundamental Forces

Having derived the nature of the fundamental constants, we can now reinterpret the force laws they govern, revealing that they are not disparate laws, but complementary modal projections of a single underlying mechanism: *momentum exchange through the elastic and dissipative vacuum*.

### 1. Newton's and Coulomb's Laws as Damped Momentum Transfer

From the previously derived first-order expressions, such as  $G \equiv \mu_0 \cdot \alpha^2$  and  $k_B \equiv \mu_0/c^2$ , it follows that:

$$G \equiv k_B \cdot \alpha^2 \cdot c^2$$

Using this expression, we can rewrite Newton's law to reveal its thermo-entropic nature. This force is mediated by the exchange of *damped, longitudinal momentum*, and expressed in terms of damped relativistic momenta, where  $\zeta = \alpha$  is the norm of the structural damping tensor:

$$F_g \equiv k_B \cdot \frac{(Mc \cdot \zeta) \cdot (mc \cdot \zeta)}{r^2} \quad (124)$$

Thus, the terms in the numerator correspond to effective, damped relativistic momenta, whose propagation is modulated by the damping structure encoded in  $\zeta_{\mu\nu}$ . Gravity emerges as the effective resistance to the coherent alignment of projected momenta through the vacuum's dissipative tensorial geometry.

By symmetry, the Coulomb force can be reformulated as the exchange of *undamped, transverse momentum*:

$$F_e = K_e \cdot \frac{Q_1 Q_2}{r^2} = \frac{\mu_0 \cdot c^2}{4\pi} \cdot \frac{Q_1 Q_2}{r^2} = \mu_0 \cdot \frac{(Q_1 c)(Q_2 c)}{4\pi r^2} \quad (125)$$

This structure mirrors the gravitational expression, with  $Qc$  playing the role of transverse modal momentum and the vacuum magnetic permeability  $\mu_0$  acting as the transverse field stiffness. Thus, both Newton's and Coulomb's laws appear as complementary modal projections of the same unified vacuum tensorial response.

## 2. Physical Interpretation: the Structural Differences

The distinct mathematical forms of Eqs. 124 and 125 are not accidental; they reflect profound differences in the physical nature of the momentum exchange. Comparing the rewritten laws for the gravitational (Eq. 124) and Coulomb (Eq. 125) forces reveals two fundamental structural differences that provide deep insight into their distinct physical natures.

*a. 1. The Selective Role of Damping.* A crucial difference is the explicit presence of the damping factor,  $\zeta$ , in the gravitational force, while it is absent in the static Coulomb force. This does not imply that electromagnetism is an undamped phenomenon—indeed, we have argued that the propagation of light is damped. Rather, it reveals the fundamental character of each interaction:

- The **Coulomb force** is a *conservative interaction* between static charges. Its formulation does not involve dissipation. The dissipative effects of electromagnetism arise in dynamic phenomena, such as radiation.
- The **gravitational force**, when expressed in this thermo-entropic form, is revealed to be an inherently *dissipative and entropic interaction*. It is not a static potential force in the classical sense, but the result of momentum exchange through a dissipative medium. Therefore, it *must* explicitly include the damping factor  $\zeta$  as part of its fundamental definition.

This suggests that damping is a key feature that distinguishes the entropic (longitudinal) modes of the vacuum from the conservative (transverse) static modes.

*b. 2. The Geometry of Propagation ( $r^2$  vs.  $4\pi r^2$ ).* The second key difference lies in the geometry of the denominator.

- The **Coulomb force** includes the factor  $4\pi r^2$ , the surface area of a sphere. This reflects the isotropic, wave-like nature of the interaction, where the influence of a point charge propagates outwards uniformly in all directions, spreading over a spherical surface. This is characteristic of a **transverse wave**.

- The re-expressed **gravitational force**, by contrast, lacks the  $4\pi$  solid angle factor. Its  $r^2$  dependence represents a direct interaction between two points. This supports the interpretation of a **longitudinal interaction**, where the force acts as a direct "pressure" or "tension" along the line connecting the two masses, rather than as a field radiating spherically.

In summary, both interactions reflect momentum exchange across a structured medium, but what differs is the symmetry of the exchange. The immense difference in their strengths arises not from arbitrarily different coupling constants, but from the physical properties of the vacuum itself:

- The gravitational force emerges as a highly suppressed, entropy-weighted longitudinal momentum flow. Gravity is weak not because its coupling is small, but because the vacuum is extremely rigid against this type of compressional deformation, and the Boltzmann constant  $k_B$  quantifies the high entropic cost required.
- The electromagnetic force, in contrast, reflects a much more efficient, transversely mediated momentum exchange, amplified by the vacuum's comparatively soft resistance to shear-like deformations, a process governed by  $\mu_0$ .

## XIX. SYNTHESIS: UNIFYING THE VACUUM'S ELASTIC AND QUANTUM PROPERTIES

We have now established a complete, two-scale framework for the gravitational constant derived from both static and dynamic principles. The ultimate test of this framework is its internal consistency. We will now demonstrate that the relationships derived are not only mutually compatible but also lead to profound connections between the vacuum's elastic, electromagnetic, and quantum characteristics.

### A. A Consistency Condition for the Vacuum

We have derived the gravitational constant  $G$  from two independent perspectives, grounded in the vacuum's elastic properties:

- **Static Origin:** Based on a principle of reciprocity, where  $G$  is the geometrically-scaled compliance of the vacuum:

$$G = \left(\frac{3}{5}\right) 4\pi\epsilon_0$$

- **Dynamic Origin:** Based on a model of quadratic damping, where  $G$  is the primary vacuum stiffness, quadratically suppressed by dissipation:

$$G = \mu_0 \alpha^2$$

For this framework to be self-consistent, these two expressions for  $G$  must be equal. Equating them reveals a profound **consistency condition** that the vacuum's properties must obey:

$$\boxed{\left(\frac{3}{5}\right) 4\pi\epsilon_0 \equiv \mu_0 \alpha^2} \quad (126)$$

This equation is a powerful prediction of the theory. Using the relation  $Z_0^2 = \frac{\mu_0}{\epsilon_0}$ , we can rewrite this condition as:

$$Z_0 \equiv \frac{\sqrt{\frac{3}{5}4\pi}}{\alpha} \approx 376.28 \Omega \quad (127)$$

where one can identify the second-order term and write the more exact expression

$$Z_0 \equiv \frac{\sqrt{\frac{3}{5}4\pi}}{\alpha} \left( 1 + \frac{\alpha}{2\pi} + \dots \right) \approx 376.73 \Omega$$

Another direct and powerful corollary is a relationship between  $\alpha$ ,  $G$ , and  $\mu_0$ . Rearranging Eq. (126) yields:

$$\alpha \equiv \sqrt{\frac{G}{\mu_0}} \quad (128)$$

This simple expression confirms that  $\alpha$  is not an independent constant, but a *geometric attenuation coefficient* derived from the ratio of the vacuum's longitudinal and transverse stiffnesses, and elevates the fine-structure constant from a mere electromagnetic coupling parameter to a *universal measure of vacuum damping topology*, governing how all interactions manifest.

## B. Consistency Check: Emergence of the Static Reciprocity Principle

In Section VIA, we motivated the need for a principle of static reciprocity, where the vacuum's compliance to gravitational deformation (related to  $G$ ) should be inversely related to its stiffness against electromagnetic deformation (related to  $K_e$ ) to ensure stability. Having now derived the gravitational constant  $G$  from dynamic principles, we can check if this physical requirement is automatically satisfied by the theory, rather than being a separate postulate.

One can compute the ratio of the gravitational compliance to the electromagnetic stiffness. This yields:

$$\frac{G}{K_e} = \frac{\mu_0 \alpha^2}{\mu_0 c^2 / 4\pi} = \frac{4\pi \alpha^2}{c^2} \quad (129)$$

This result is profound. It demonstrates that the relationship between  $G$  and  $K_e$  is not a simple inversion. Instead, it is precisely what one would expect from a dissipative elastic substrate. The leading-order effect of dissipation on some system's elastic structure is expected to be the simplest non-trivial functional dependence. A linear relationship implies a directional dependence, violating parity, while a constant offset is trivial. The next simplest and most natural form, consistent with second-order dissipative effects (proportional to power loss), is a quadratic dependence. Finding that the gravitational compliance ( $G$ ) is indeed inversely proportional to the electromagnetic stiffness ( $K_e$ ), and that is also quadratically suppressed by the universal damping invariant of the vacuum ( $\alpha^2$ ), becomes a *successful, non-trivial consistency check* of the QEG framework. The theory, by virtue of its dynamic and dissipative structure, naturally produces the hierarchy of forces required for a stable substrate.

## C. The quality factor as a Quantum-Electrodynamic Property

From the definition of the fine-structure constant, we have:

$$2\alpha = \frac{e^2}{2\pi\epsilon_0\hbar c}$$

Using the key consistency condition derived from our model  $G_{Glob} \equiv 2\pi\epsilon_0 \equiv 2\alpha/c$  (207), and substituting this into the denominator gives:

$$2\alpha \equiv \frac{e^2}{\left(\frac{2\alpha}{c}\right)\hbar c} = \frac{e^2}{2\alpha\hbar} \rightarrow \boxed{4\alpha^2 \equiv \frac{1}{Q^2} \equiv \frac{e^2}{\hbar}} \quad (130)$$

This remarkable result implies that the ability of a channel to conduct charge is determined by the dissipative properties of the spacetime in which it exists.

The Quantum Hall Effect (QHE) shows that resistance is quantized in units of the **von Klitzing constant**,  $R_K$ :

$$R_K = \frac{h}{e^2} = 2\pi \frac{\hbar}{e^2} \equiv 2\pi Q^2$$

This implies that the quantized resistance measured in a 2D electron gas is not an emergent property of the material system alone, but rather a direct manifestation of the vacuum's intrinsic damping. The electrons in a QHE system are, in effect, probing the fundamental dissipative structure of spacetime itself.

## D. A Unified Emergence Mechanism for Gravity and Charge

The internal consistency of our framework reveals a profound structural symmetry between the origin of the local gravitational constant,  $G_{Loc}$ , and the squared elementary charge,  $e^2$ . As derived in Section XVIII B:

$$G_{Loc} \equiv \mu_0 \cdot \zeta^2 \quad (131)$$

We now derive an analogous expression for the squared elementary charge. Starting from our derived first-order equivalence  $e \equiv \frac{2\alpha \cdot 1\text{m}}{c^2}$ , squaring both sides yields:

$$e^2 \equiv 4\alpha^2 \cdot \frac{(1\text{m})^2}{c^4} \rightarrow \frac{1}{4}e^2 \equiv \hbar \cdot \zeta^2$$

The parallel between the two results is striking and reveals a universal mechanism. Both gravity and charge are emergent properties determined by the same squared damping factor,  $\zeta^2 \equiv \alpha^2$ , acting as a "*universal dissipative filter*." The fundamental interactions and charges of our universe become "*echoes*" of more primordial vacuum properties ( $\mu_0$  and  $\hbar$ ), all filtered through the same dissipative process ( $\zeta^2$ ). The difference in their nature and magnitude arises from their distinct origins:

- **Gravity** emerges from the *classical, elastic* properties of the vacuum ( $\mu_0$ ).
- **Charge** emerges from the *fundamental, quantum* properties of the vacuum ( $\hbar$ ), and as an effective electromagnetic field coupling, it adopts the canonical normalization factor of  $\frac{1}{4}$ .

This shared emergence mechanism is one of the most powerful pieces of evidence for the coherence of the proposed framework, suggesting a deep unity in the principles that govern the cosmos.

### E. A Bridge to General Relativity: An Ohm's Law for the Vacuum

Additionally, our framework provides a remarkable consistency check with General Relativity. Substituting  $G \equiv \mu_0 \cdot \alpha^2$  one can check that

$$G \cdot c \equiv \mu_0 \cdot c \cdot \alpha^2 = Z_0 \cdot \alpha^2 \approx \frac{1}{50.13} \Omega \quad (132)$$

The above implies that the product  $G \cdot c$  defines a fundamental 'resistive-like' constant of the vacuum, which we denote as the *natural resistance*  $X_N$ . The relationship  $G \equiv \frac{1}{16\pi c}$  can be interpreted as an Ohm's Law for the vacuum,  $V = I \cdot R$ , where:

- **Potential ( $V$ ):** The gravitational constant  $G$ , within our dimensionally collapsed framework, acquires the role of a fundamental potential or electromotive force, as  $G \equiv \mu_0 \cdot \zeta^2$  and  $[\mu_0] = [V]$  (XVIC).
- **Current ( $I$ ):** The term  $1/c$  represents the natural, scaled current of the vacuum for the thermotropic mode (XVIC).
- **Resistance ( $R$ ):** This implies that the vacuum possesses an intrinsic, dimensionless resistance  $R_{\text{vac}} = 1/(16\pi)$ .

This vacuum resistance is not an arbitrary number, but can be derived from first principles by decomposing it into two meaningful factors:

$$R_{\text{vac}} = \frac{1}{16\pi} = \frac{1}{4} \cdot \frac{1}{4\pi} \quad (133)$$

The two terms have clear physical origins:

1. **The Geometric Resistance ( $1/4\pi$ ):** This factor arises directly from the Green's function of the 3D Laplacian operator (II E). It represents the fundamental geometric impedance of three-dimensional space, quantifying how the influence of a point source is diluted as it spreads over a spherical solid angle.
2. **The Canonical coupling factor ( $1/4$ ):** This factor is the canonical normalization constant required for any standard gauge field theory, as seen in the electromagnetic Lagrangian  $\mathcal{L}_{EM} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$ . Its presence here suggests that gravity, as an *emergent effective field*, must inherit the canonical normalization of the underlying field theory framework.

Therefore, the relationship we propose is not a numerical coincidence, but a profound statement about the structure of the vacuum. Consider the equation

$$G \equiv \frac{1}{c} \cdot \left( \frac{1}{4} \cdot \frac{1}{4\pi} \right) \implies c \equiv \frac{1}{16\pi G} \quad (134)$$

In Sec. VIII A, we established a correspondence principle identifying the macroscopic gravitational stiffness,  $1/(16\pi G)$ , with the microscopic stiffness of the substrate,  $\kappa$ . The identity in Eq. (134) therefore implies a direct proportionality between the speed of light and the substrate's intrinsic stiffness:

$$c \propto \kappa \quad (135)$$

This is a remarkable and intuitive consequence of the theory. It suggests that the speed of light is not an independent, postulated axiom, but rather an emergent property determined by the elastic characteristics of the vacuum. This is the hallmark of

wave propagation in an elastic medium: *the stiffer the medium, the faster the wave*. The QEG framework thus unifies the kinematics of spacetime ( $c$ ) with the substrate's constitutive properties ( $\kappa$ ), providing a mechanical origin for the cosmic speed limit.

Subsequently, this result serves as a fundamental "bridge" between our elastic vacuum model and the geometric formulation of General Relativity. It shows that the pre-factor in the Einstein-Hilbert action,

$$S_{EH} = \frac{c^4}{16\pi G} \int R\sqrt{-g} d^4x,$$

is not a random assortment of constants. Instead, the term  $c^4/(16\pi G)$  can be seen as the "stiffness" of spacetime against curvature, emerging directly from the interplay between the vacuum's fundamental current, its geometric properties, and the canonical structure of field theory (1/4).

### F. The Impedance of Free Space as a Damped Geometric Resistance

Note that we can rewrite eq. 132 just substituting  $\zeta \equiv \alpha$  and  $\frac{1}{50.13} \equiv \frac{1}{16\pi}$ , to have that:

$$\boxed{Z_0 \cdot \zeta^2 \equiv \frac{1}{4} \cdot \frac{1}{4\pi}} \quad (136)$$

where the geometric interpretation of the terms in the left has already been exposed (XIX E). Thus, the physical impedance of the vacuum,  $Z_0$ , can be described as the fundamental geometric resistance modulated by the vacuum's own damping factor,  $\zeta = \alpha$ . This equation unifies geometry, quantum dissipation, and electromagnetism in a single, elegant expression.

### G. First-Principles Prediction of the Fine-Structure Constant

A profound consequence of the vacuum's self-consistency is that the value of the fine-structure constant,  $\alpha$ , becomes uniquely determined by the geometric and topological constraints of the framework. In the preceding sub-subsections, we derived two independent, constitutive laws for the vacuum impedance:

1. From the consistency between the static and dynamic origins of the gravitational constant  $G$ , we obtained (Eq. XIX A):

$$Z_0 \equiv \frac{\sqrt{\frac{3}{5}4\pi}}{\alpha} \quad (137)$$

2. From the interpretation of  $Z_0$  as a damped geometric resistance, we derived (Eq. 136):

$$Z_0 \equiv \frac{1}{4} \frac{1}{4\pi\zeta^2} = \frac{1}{16\pi\alpha^2} \quad (138)$$

For the theoretical framework to be internally consistent, these two expressions for  $Z_0$  must be equal. Equating Eq. 137 and Eq. 138 forces a single, unique value for the fine-structure constant:

$$\frac{\sqrt{\frac{3}{5}4\pi}}{\alpha} \equiv \frac{1}{16\pi\alpha^2} \quad (139)$$

Solving for  $\alpha$  yields:

$$\alpha \equiv \frac{1}{16\pi\sqrt{\frac{3}{5}4\pi}} \quad (140)$$

where one can identify the second-order term and write the more exact expression

$$\alpha \equiv \frac{1}{16\pi\sqrt{\frac{3}{5}4\pi}} \left(1 + \frac{\alpha}{2\pi} + \dots\right)$$

The first-order predicted value matches the experimentally measured CODATA value of  $\alpha \approx 1/137.036$  with remarkable accuracy, and is highly significant. It shows that the value of the fine-structure constant is not an arbitrary input, but rather the precise value required to harmonize the vacuum's fundamental properties: its geometric impedance arising from the 3D Laplacian ( $\frac{1}{4\pi}$ ), its canonical field structure inherited from gauge theories ( $\frac{1}{4}$ ), and the geometry of its self-interaction energy ( $\sqrt{\frac{3}{5}4\pi}$ ). This elevates the status of  $\alpha$  from a mere electromagnetic coupling parameter to the primary geometric and topological constant of the unified vacuum.

As a final note, for the above interpretation to be consistent, then  $\frac{1}{\sqrt{\frac{3}{5}4\pi}}$  must have dimensions of reciprocal of a resistance. We will uncover and check that this is indeed the case in (XIX H).

## H. Impedance vs. Dissipative Resistance and the quality factor $Q$

The robustness of this theoretical framework can be further tested by examining the consistency of the classical RLC oscillator analogy, which has served as a powerful conceptual guide. Consider the standard formula for the quality factor of a series RLC circuit at resonance:

$$Q = \frac{\omega_0 L}{R} \quad (141)$$

Within our framework, we have already established the following vacuum parameters:

- The vacuum quality factor, derived from quantum conductance, is  $Q = 1/(2\alpha)$  (51).
- The characteristic inductive term is  $\omega_0 L = (\frac{c}{1s})(\mu_0 \cdot 1m)$ . Given the spacetime equivalence where  $1s \equiv 1m$ , this simplifies to  $\omega_0 L = \mu_0 c$ , which is precisely the definition of the vacuum's wave impedance,  $Z_0$ .

Substituting these established quantities into the classical Q-factor formula allows us to solve for the effective resistance,  $R_Q$ , that must govern this specific dissipative process:

$$\frac{1}{2\alpha} = \frac{Z_0}{R_Q} \quad (142)$$

This consistency requirement leads directly to a new constitutive law for the vacuum:

$$R_Q = 2\alpha Z_0 \equiv \frac{Z_0}{Q} \equiv 2 \cdot \sqrt{\frac{3}{5}4\pi} \quad (143)$$

This result is profound. It reveals that the vacuum possesses two distinct, conceptually different resistive

properties:

**1. Wave Impedance ( $Z_0$ ):**  $Z_0$  is the vacuum's characteristic impedance to wave propagation. It is a reactive property that governs the ratio of the electric and magnetic field strengths in an electromagnetic wave.

**2. Dissipative Resistance ( $R_Q$ ):** This newly derived quantity,  $R_Q$ , is the vacuum's effective resistance governing the *rate of energy dissipation per cycle* in an oscillation, as quantified by the quality factor. It is a fundamentally dissipative, rather than reactive, property.

Equation 143 provides the explicit relationship between these two properties. It states that the dissipative resistance of the vacuum is its fundamental wave impedance modulated by its own universal quality factor,  $Q$ . This is physically intuitive: the resistance to energy loss should be proportional to both the resistance to wave propagation and the intrinsic friction of the medium.

Note that using Eq. 143, we can express  $\frac{3}{5}4\pi = \frac{1}{4}R_Q^2$ . Substituting this into the left-hand side of the master consistency condition (Eq. 126) yields:

$$\frac{1}{4}\varepsilon_0 R_Q^2 \equiv \mu_0 \alpha^2 \quad (144)$$

This final expression is a remarkably powerful statement of unification. The dimensional consistency of Eq. 144 holds in both the SI system and the dimensionally-collapsed framework of this theory, further cementing its robustness.

## I. Dynamic Origin of Permittivity from Vacuum Power Principles

Having established the distinct roles of wave impedance ( $Z_0$ ) and dissipative resistance ( $R_Q$ ), we can now propose a dynamic origin for the vacuum's electric permittivity,  $\varepsilon_0$ . We move beyond the static picture of permittivity as a passive capacity to store fields and instead derive it from the vacuum's fundamental power dissipation and resistance properties.

First, we define a *Unitary Vacuum Power*,  $P_{unit}$ , as the power dissipated when the vacuum's intrinsic structural voltage ( $V \equiv \mu_0$ ) is applied across a unitary resistance ( $R = 1\Omega$ ). According to Joule's Law:

$$P_{unit} = \frac{V^2}{R} \equiv \frac{\mu_0^2}{1\Omega} \quad (145)$$

In this framework,  $P_{unit}$  represents the baseline rate of energy transfer or dissipation inherent to the vacuum's primary potential.

Next, we postulate that the vacuum's permittivity,  $\varepsilon_0$ , which represents its compliance or ability to "permit" an electric field, is an emergent property. It arises from this unitary power being modulated by the vacuum's own internal friction. The most natural choice for this friction is the **dissipative resistance**,  $R_Q$ , as it governs energy loss per oscillatory cycle. We therefore propose the following constitutive law:

$$\varepsilon_0 \equiv P_{unit} \cdot R_Q = \frac{\mu_0^2}{1\Omega} R_Q \equiv \mu_0^2 \cdot 2\sqrt{\frac{3}{5}4\pi} \quad (146)$$

where one can identify the second-order term and write the more exact expression

$$\varepsilon_0 \equiv \left( \mu_0^2 \cdot 2\sqrt{\frac{3}{5}4\pi} \right) (1 + 2\alpha + \dots)$$

### J. The Gravito-Entropic Power Equivalence

Building upon this last relationship, we can establish a direct link between the vacuum's primary inertial properties and its large-scale cosmological dissipation. Eq. 146 yields a direct equivalence:

$$\frac{1}{4}\mu_0^2 \equiv 2\pi\varepsilon_0 \cdot \alpha \equiv G_{glob} \cdot \zeta \quad (147)$$

This equation describes how the vacuum's capacity to store inertial or reactive energy in its transverse (electromagnetic) modes is intrinsically related to the vacuum's compliance to longitudinal deformation ( $G_{glob}$ ) and its damping factor ( $\zeta$ ). Ultimately, Eq. 147 signifies a profound equilibrium: *the vacuum's capacity to store inertial energy in its transverse mode is perfectly balanced by the power it dissipates in its longitudinal mode*. This relationship links the electromagnetic and gravitational sectors through a fundamental equilibrium.

### K. The Elementary Charge as a high-order stiffness of the Vacuum

An additional insight into the vacuum's fundamental properties arises when substituting the geometric definition of  $\alpha$  from Eq. 140 into the expression  $e \equiv \frac{2\alpha \cdot 1}{c^2} m$  (XVII A):

$$e \equiv \frac{2 \cdot 1}{c^2} m \left( \frac{1}{16\pi\sqrt{\frac{3}{5}4\pi}} \right) = \frac{1}{c^2} \frac{m}{8\pi\sqrt{\frac{3}{5}4\pi}}$$

Using the fundamental relation  $\mu_0\varepsilon_0 = 1/c^2$  to replace  $c^2$ , and substituting the permittivity  $\varepsilon_0$  from Eq. 146, yields:

$$\begin{aligned} e &\equiv \mu_0 \left( \mu_0^2 \cdot 2\sqrt{\frac{3}{5}4\pi} \right) \cdot \frac{1}{8\pi\sqrt{\frac{3}{5}4\pi}} \\ &= \frac{1}{8\pi\sqrt{\frac{3}{5}4\pi}} \cdot 2\mu_0^3\sqrt{\frac{3}{5}4\pi} = \frac{\mu_0^3 \cdot 1}{4\pi} m \rightarrow \boxed{e \equiv \frac{\mu_0^3 \cdot 1}{4\pi} m} \end{aligned} \quad (148)$$

where one can identify the second-order term and write the more exact expression

$$e \equiv \frac{\mu_0^3 \cdot 1}{4\pi} m \cdot (1 + 2\alpha + \dots)$$

This result shows how charge is a direct, emergent property of the vacuum itself, determined solely by its most fundamental characteristic—the transverse inertial stiffness  $\mu_0$ —projected through the geometry of three-dimensional space ( $\frac{1}{4\pi}$ ).

The cubic dependence ( $e \propto \mu_0^3$ ) signifies that charge is a non-linear excitation of the vacuum field. While the vacuum's primary elastic response is linear (as seen in wave propagation), the formation of a

stable, quantized charge represents a higher-order, self-interaction phenomenon. It is a measure of the vacuum's capacity to sustain a localized, persistent deformation against its own inertial resistance.

Importantly, this formulation elevates the magnetic permeability  $\mu_0$  to the status of the single, primary constant of the electromagnetic sector. The elementary charge  $e$  and the electric permittivity  $\varepsilon_0$  are both derived from it, establishing a clear hierarchy of fundamental constants.

Finally, equation 148 defines vacuum's capacity to manifest charge as a function of its intrinsic stiffness, contributing a big step into the geometrization of physics by demonstrating that not only forces, but also the sources of those forces, arise from the fabric of spacetime.

### L. The Quantum of Action as an Emergent Property of Vacuum Compliance

Following the derivation of the elementary charge from the vacuum's inertial stiffness ( $\mu_0$ ), we show how the quantum of action,  $\hbar$ , can also be derivable from the vacuum's complementary elastic property: its compliance, or permittivity ( $\varepsilon_0$ ). This demonstration serves to ground not only the sources of forces, but the very granularity of quantum mechanics, in the tangible, elastic properties of the vacuum.

Our starting point is the modal action for the electromagnetic field derived in Eq. 149, which established the dimensionality of action as a spacetime area:

$$\hbar \equiv \frac{1}{c^4} m^2 \quad (149)$$

To express  $\hbar$  in terms of permittivity, we use the fundamental relationship  $c^4 = 1/(\mu_0^2\varepsilon_0^2)$  into Eq. (149), which yields:

$$\hbar = (1 m^2) \cdot (\mu_0^2\varepsilon_0^2)$$

Replacing the inertial term  $\mu_0^2$  with its equivalent expression derived from the vacuum's dissipative properties, as given in Eq. 146, reveals  $\hbar$  as a pure function of vacuum compliance and geometry:

$$\begin{aligned} \hbar &= (1 m^2) \cdot \left( \frac{\varepsilon_0}{2\sqrt{\frac{3}{5}4\pi}} \right) \cdot \varepsilon_0^2 \\ &= \frac{\varepsilon_0^3 \cdot (1 m^2)}{2\sqrt{\frac{3}{5}4\pi}} \end{aligned}$$

This leads to a new fundamental expression for the reduced Planck constant:

$$\boxed{\hbar \equiv \frac{\varepsilon_0^3 \cdot (1 m^2)}{2\sqrt{\frac{3}{5}4\pi}}} \quad (150)$$

This result provides a compelling physical origin for the quantum of action, reinterpreting it not as an axiom, but as a direct consequence of the vacuum's elastic structure. The equation suggests that the fundamental discreteness of physical processes, quantified by  $\hbar$ , is an emergent phenomenon rooted in the vacuum's ability to permit electric fields ( $\varepsilon_0$ ), and thus a built-in consequence of the elastic and geometric properties of spacetime itself.

The cubic dependence on permittivity ( $\hbar \propto \varepsilon_0^3$ ) is highly significant. It indicates that quantum action arises from a complex, volumetric self-interaction of the vacuum's compliance. It can be physically pictured as the total "elastic energy potential" that can be stored in a unit of spacetime area, a potential that is non-linearly dependent on the medium's softness.

Importantly, this result forms a perfect symmetry with the derivation for the elementary charge ( $e \propto \mu_0^3$ ). Together, they paint a complete picture:

- The **fundamental quantum of charge** ( $e$ ) is a *non-linear function of the vacuum's inertial stiffness* ( $\mu_0$ ).
- The **fundamental quantum of action** ( $\hbar$ ) is a *non-linear function of the vacuum's elastic compliance* ( $\varepsilon_0$ ).

This duality between stiffness/charge and compliance/action represents a deep, foundational symmetry of the vacuum. It suggests that the laws of electromagnetism and the laws of quantum mechanics are two sides of the same coin, both emerging from the fundamental tension between inertia and compliance in the fabric of reality.

This derivation solidifies the framework's central claim: the constants that define our physical laws are not a random assortment of numbers, but are deeply interconnected parameters that describe the single, underlying, elastic quantum vacuum.

### M. Derivation of Vacuum Constants and the Speed of Light from a Single Parameter

The ultimate test of the framework's internal consistency lies in its ability to define the fundamental constants of the vacuum,  $\mu_0$  and  $\varepsilon_0$ , and subsequently the speed of light,  $c$ , from a single, dimensionless parameter. Using the equivalence between the static and dynamic origins of the gravitational constant  $G$  (from Eq. 126) and the expression for the dynamic origin of permittivity from vacuum power principles (from Eq. 146), we can substitute Eq. (146) into Eq. (126) to obtain that:

$$\begin{aligned} \left(\frac{3}{5}\right) 4\pi \left( \mu_0^2 \cdot 2\sqrt{\frac{3}{5}4\pi} \right) &\equiv \mu_0 \alpha^2 \rightarrow \\ \mu_0 &\equiv \alpha^2 \cdot \frac{1}{16\pi \cdot \sqrt{\frac{3}{5}4\pi}} \cdot \frac{10}{3} \rightarrow \\ \boxed{\mu_0 &\equiv \frac{10}{3} \alpha^3} \end{aligned} \quad (151)$$

where we have used  $\alpha \equiv \frac{1}{16\pi \cdot \sqrt{\frac{3}{5}4\pi}}$  (140) and the derived result is a first-order approximation. Using this result, we can express  $\varepsilon_0$  purely in terms of  $\alpha$ :

$$\begin{aligned} \left(\frac{3}{5}\right) 4\pi \varepsilon_0 &\equiv \frac{10}{3} \alpha^3 \alpha^2 \rightarrow \\ \boxed{\varepsilon_0 &\equiv \left(\frac{5}{3}\right)^2 \cdot \frac{1}{2\pi} \cdot \alpha^5} \end{aligned} \quad (152)$$

Finally, with both vacuum constants defined by  $\alpha$  and geometry, we derive the speed of light using the fun-

damental relation  $c^2 = 1/(\mu_0 \varepsilon_0)$ :

$$\begin{aligned} c^2 &\equiv \frac{1}{\left(\frac{10}{3}\alpha^3\right) \cdot \left(\frac{25}{18\pi}\alpha^5\right)} \rightarrow \\ c^2 &\equiv \frac{1}{\frac{125}{27\pi} \cdot \alpha^8} \rightarrow \boxed{c \equiv \frac{1}{\sqrt{\frac{125}{27\pi}} \cdot \alpha^4}} \end{aligned} \quad (153)$$

This result represents the culmination of the unified framework. The constants  $\mu_0$ ,  $\varepsilon_0$ , and  $c$  are no longer fundamental in their own right. They are revealed to be interdependent functions of a single, more primary parameter,  $\alpha$ . The entire framework of vacuum electrodynamics and spacetime kinematics is determined not just by a number, but by the *scalar norm* ( $\alpha$ ) of the rank-2 symmetric *structural damping tensor*,  $\zeta_{\mu\nu}$ , which encodes the vacuum's intrinsic dissipative properties.

Additionally, the relationship  $c \propto \alpha^{-4}$  confirms that the product  $c \cdot \alpha^4$  is indeed a constant composed of the geometric factors derived throughout this section. As anticipated in Section XII B, this is precisely the normalization constant required to reconcile the first-order, bare geometric expression for  $\hbar$  with its physical value, thus closing a crucial loop of internal coherence in the theory.

Moreover, the derived relationship,  $c \propto \alpha^{-4}$ , offers a profound insight into the nature of causality. The speed of light is not an arbitrary limit but an *emergent property* dictated by the vacuum's inherent "friction". A hypothetical, perfectly frictionless vacuum ( $\alpha \rightarrow 0$ ) would permit an infinite propagation speed, rendering causality instantaneous. It is therefore the small, non-zero damping of spacetime—the inherent viscosity of the quantum oscillator lattice—that establishes a finite causal speed limit, giving the universe its structure in time.

This synthesis elevates the fine-structure constant to the role of the primary architect of physical reality. It is the scalar measure of the vacuum's fundamental dissipative geometry. The value of  $\alpha$  dictates the vacuum's inertial resistance ( $\mu_0$ ) and elastic compliance ( $\varepsilon_0$ ), and the interplay between these two properties, governed by  $\alpha$ , sets the exact value of the cosmic speed limit,  $c$ . Knowing the norm of the vacuum's damping tensor is equivalent to knowing the fundamental operational rules of spacetime.

### N. Geometric Origin of $\alpha$ and $c$ as a Consistency Check

To close the synthesis of the framework, we provide a consistency check showing that both the value of the fine structure constant  $\alpha$  and the causal speed  $c$  emerge from purely geometric principles of minimal structure and dissipation.

*$\alpha$  as minimal geometric dissipation.*

As we have already pointed out, the fine structure constant derived in Eq. 140 can be rewritten as

$$\alpha = \frac{1}{16\pi \sqrt{\frac{3}{5}4\pi}} = \left(\frac{1}{4}\right) \cdot \left(\frac{1}{4\pi}\right) \cdot \left(\frac{1}{\sqrt{\frac{3}{5}4\pi}}\right). \quad (154)$$

This decomposition makes explicit three irreducible geometric factors:

- The *gauge factor*  $1/4$ , the canonical normalization for any covariant gauge field Lagrangian,
- The *Laplacian Green factor*  $1/(4\pi)$ , encoding the minimal isotropic flux dilution in three spatial dimensions,
- The *self-interaction factor*  $1/\sqrt{(3/5)4\pi}$ , representing the lowest-energy configuration of a uniform spherical distribution.<sup>6</sup>

The product of these three factors singles out  $\alpha$  as the minimal dissipation constant compatible with gauge consistency, spatial isotropy, and stable self-interaction. Any smaller value would be insufficient to stabilize these constraints simultaneously.

#### Variational Origin of the Fine-Structure Constant

The above can be based more deeply as the solution to a variational problem. Consider an effective scalar channel  $\Phi$  (trace mode) and a gauge sector  $A_\mu$  (torsional/EM) in a four-volume  $V$ . We write a conservative  $U(1)$ -invariant term plus a linear (Rayleigh-type) dissipation term:

$$S_{\text{eff}}[\Phi, A] = \underbrace{\alpha \int_V d^4x \mathcal{D}[\Phi, A]}_{\text{dissipation (to be minimized)}} + \underbrace{\frac{1}{4} \int_V d^4x F_{\mu\nu} F^{\mu\nu}}_{\text{fixed gauge term}} \quad (155)$$

where  $\mathcal{D} \geq 0$  encodes the positive-definite dissipative density and  $\alpha > 0$  measures the substrate damping. We impose three geometric normalization constraints that fix the relative scale of fields unambiguously:

(R1) *Gauge* Canonical gauge prefactor:  $\mathcal{N}_g = \frac{1}{4}$ .

(R2) *Laplacian/Green* On 3D slices, the Green function of  $-\nabla^2$  carries prefactor  $1/(4\pi)$ :

$$\int_{S^2_{\infty}} d\mathbf{S} \cdot \nabla G(\mathbf{x}) = -1 \implies G(\mathbf{x}) = \frac{1}{4\pi|\mathbf{x}|}, \quad (156)$$

hence  $\mathcal{N}_\Delta = \frac{1}{4\pi}$ .

(R3) *Self-interaction* The isotropic self-energy scale in a spherical cell is governed by  $\sqrt{(3/5)4\pi}$  (moment of inertia factor  $\frac{3}{5}$  and solid angle  $4\pi$ ), thus  $\mathcal{N}_{\text{self}} = \sqrt{(3/5)4\pi}$ .

Introduce Lagrange multipliers  $(\lambda_g, \lambda_\Delta, \lambda_{\text{self}})$  and define the constrained functional

$$\mathcal{J}(\alpha) = \alpha \underbrace{\int_V d^4x \mathcal{D}}_{\equiv \mathcal{Q} > 0} + \lambda_g \left( \frac{1}{4} - \mathcal{N}_g \right) + \lambda_\Delta \left( \frac{1}{4\pi} - \mathcal{N}_\Delta \right) + \lambda_{\text{self}} \left( \sqrt{(3/5)4\pi} - \mathcal{N}_{\text{self}} \right) \quad (157)$$

As (R1)–(R3) fix the relative normalizations of  $\Phi$  and  $A_\mu$ , the only remaining global freedom that can reduce the total dissipation is  $\alpha$ . A parsimonious closure is to require that the three geometric channels contribute *equally* to the dissipative budget. Equivalently, for a fixed product of geometric factors, the arithmetic mean is minimized when all contributions are balanced (AM–GM inequality). This uniquely selects the stationary (minimal-dissipation) value

$$\alpha_{\text{min}} = \frac{1}{16\pi \sqrt{\frac{3}{5} 4\pi}} \quad (158)$$

which coincides with Eq. 140. Thus,  $\alpha$  is not merely the product of independent geometric constants, but the *unique minimizer* of the global dissipation under the constraints (R1)–(R3).

#### Minimal action as minimal dissipation.

In this picture, the total dissipation along a trajectory is proportional to its geometric action:

$$D_{\text{path}} = \alpha S_{\text{path}}. \quad (159)$$

If the vacuum operates at the minimal dissipation value  $\alpha = \alpha_{\text{min}}$ , then the stability condition becomes

$$\delta D_{\text{path}} = \alpha_{\text{min}} \delta S_{\text{path}} = 0 \implies \delta S_{\text{path}} = 0. \quad (160)$$

Thus, the Principle of Least Action arises naturally as the principle of least dissipation: physical systems evolve along those histories that minimize deformation in a substrate already tuned to the lowest possible dissipative regime.

#### Causal velocity as dimensional scaling.

Propagation efficiency is inversely related to dissipation. In a four-dimensional spacetime, the most parsimonious scaling law links the causal speed  $c$  to the inverse dissipation with exponent equal to the dimensionality:

$$c \propto \left( \frac{1}{\alpha} \right)^D, \quad D = 4. \quad (161)$$

Therefore,

$$\boxed{c \propto \alpha^{-4}}. \quad (162)$$

The exponent four is not arbitrary: it is the signature of four-dimensional spacetime. In the limit  $\alpha \rightarrow 0$ , corresponding to a frictionless substrate, propagation would be instantaneous ( $c \rightarrow \infty$ ). The finite value of  $c$  is thus a direct consequence of the minimal dissipation condition.

#### Conclusion.

This consistency check confirms that the fine structure constant  $\alpha$  is fixed by the minimal geometric requirements of gauge normalization, isotropic propagation, and self-stability, and that the causal velocity  $c$  emerges as the maximal speed consistent with a four-dimensional dissipative substrate. This provides a geometric origin both for  $\alpha$  and  $c$ , and constitutes one of the jewels of QEG predictions.

<sup>6</sup> The appearance of the square root in Eq. 143 can be traced back to the fact that self-interaction energies are quadratic in field amplitudes, while impedances and resistances are linear. For a uniform spherical source the self-energy is  $U_{\text{self}} \sim \frac{3}{5} \frac{Q^2}{4\pi\epsilon_0 R}$ , so the relevant geometric factor enters as  $\frac{3}{5} 4\pi$  at the level of energy. When relating this to an effective resistance—a linear quantity—the square root must be taken, yielding the factor  $\sqrt{\frac{3}{5} 4\pi}$

## Part V: The thermo-entropic field: a fundamental prediction from quantum-elastic geometry

### XX. THE THERMO-ENTROPIC FIELD: A FUNDAMENTAL PREDICTION FROM QUANTUM-ELASTIC GEOMETRY

#### A. Motivation for a thermo-entropic field

The plausibility of a structured field theory uniting gravitational and entropic dynamics is supported by a range of independent theoretical and empirical findings:

- **Gravitational wave observations**, notably those by LIGO and Virgo, confirm that the gravitational field  $\vec{g}$  can vary with time [44]. This supports the existence of dynamical couplings with an auxiliary field  $\vec{T}$ , where temporal variations in the entropic sector may induce circulation-like components in  $\vec{g}$ .
- **Black hole thermodynamics** reveals deep links between gravitational phenomena and thermodynamic quantities such as entropy and temperature [45, 46], supporting the idea that the entropic field  $\vec{T}$  is not a derivative phenomenon, but rather a fundamental component of spacetime structure.
- **Experimental confirmations of gravitomagnetic effects**, such as those from Gravity Probe B [47], show that rotating masses generate a field component dependent on mass currents. This behavior is consistent with the idea that a circulating mass flow  $\vec{J}_m$  contributes to the generation of a complementary field  $\vec{T}$ , in analogy with magnetism.
- **Thermodynamic derivations of gravitational dynamics**, such as Jacobson's approach to Einstein's equations [10] and Verlinde's emergent gravity framework [11], suggest that gravity may arise from underlying entropic principles.
- **Thermoelectric relationships** further strengthen the proposal. In condensed matter physics, temperature gradients generate electric potentials (Seebeck effect), while electric currents produce or absorb heat (Peltier effect) [48]. These two-way couplings between energy and entropy mirror the kind of mutual interactions expected in a thermo-entropic field theory. Additionally, the Unruh effect shows how temperature can emerge from acceleration, reinforcing the connection between thermodynamics and spacetime structure.

These observations suggest the existence of a missing dynamic sector in fundamental physics. The Quantum Elastic Geometry (QEG) framework not only accommodates this possibility, *but inevitably predicts it*. While the compressive ( $\mathcal{G}_{00}$ ) and torsional ( $\mathcal{G}_{0i}$ ) modes of the unified field naturally yield General Relativity and Electromagnetism, the components of the spatial tensor,  $\mathcal{G}_{ij}$ , must correspond to a physical interaction. We term this the *thermo-entropic field*, which formalizes the interplay between gravity and entropy through

a field pair  $\{\vec{g}, \vec{T}\}$ . Here,  $\vec{g}$  represents the gravitational field, while  $\vec{T}$  denotes a circulating field analogous to magnetism, generated by mass currents. The name *thermo-entropic* reflects this intrinsic duality between radial, mass-induced effects (gravity) and an azimuthal, thermo-entropy-induced circulation. The following analysis will show how a single master dynamics for this field leads simultaneously to (i) Maxwell-like equations for entropic currents, (ii) an effective stress-energy tensor for Einstein's field equations, and (iii) diffusive equations characteristic of irreversible thermodynamics.

#### B. Field Content and Tensor Decomposition

The spatial part of the substrate deformation,  $\mathcal{G}_{ij}$ , is a symmetric rank-2 tensor in three dimensions. It decomposes uniquely as

$$\begin{aligned}\Theta(x) &\equiv C_\Theta \delta^{ij} \mathcal{G}_{ij}(x) \quad (\text{isotropic thermal scalar}) \\ \Sigma_{ij}(x) &\equiv C_\Sigma \left( \mathcal{G}_{ij}(x) - \frac{1}{3} \delta_{ij} \mathcal{G}_{kk}(x) \right) \\ &(\text{anisotropic shear tensor})\end{aligned}\quad (163)$$

where  $C_\Theta, C_\Sigma$  are normalization constants. We propose the physical identification of the scalar trace  $\Theta$  with the dynamics of local thermal energy, and the traceless tensor  $\Sigma_{ij}$  with anisotropic shear strains. The validity of this identification will be demonstrated by showing that the resulting dynamics correctly reproduce the known phenomenology of irreversible thermodynamics and lead to testable consequences. The sources for these fields are, respectively, the isotropic pressure/heat flux  $J_\Theta$  and the anisotropic stress tensor  $\sigma_{ij}$ .

This decomposition is mathematically unique: any symmetric rank-2 tensor in 3D can be uniquely decomposed into its trace and traceless part. The physical identification of the trace with a thermo-entropic scalar field and of the traceless part with anisotropies follows directly from the analogy with continuum mechanics (elasticity), where volumetric dilatation describes isotropic compression and shear describes volume-preserving deformations. This physical identification is also required by covariance: the scalar trace  $\Theta$  is the only component that can couple to scalar sources like isotropic pressure ( $p$ ), while the traceless tensor  $\Sigma_{ij}$  is required to couple to anisotropic tensor sources like shear stress ( $\sigma_{ij}$ ).

#### C. The Master Dynamics: Telegrapher's Equation

From the QEG action (Sec. 54), the equations of motion for  $\mathcal{G}_{ij}$  include both an elastic term governed by the universal stiffness  $\kappa$  and a viscous term governed by the universal damping invariant  $\zeta$ , both established as foundational properties of the substrate in Part I:

$$\kappa \square \mathcal{G}_{ij} - \zeta \partial_t \mathcal{G}_{ij} - \frac{\delta V}{\delta \mathcal{G}_{ij}} = J_{ij}. \quad (164)$$

In continuum mechanics, dissipation is introduced via a Rayleigh functional,  $\mathcal{R}$ , proportional to the square of the rate of strain. For a relativistic field theory, this functional must be a Lorentz scalar. We define the covariant dissipation functional as:

$$\mathcal{R} = \frac{1}{2} \zeta (u^\alpha \nabla_\alpha \mathcal{G}_{\mu\nu})(u^\beta \nabla_\beta \mathcal{G}^{\mu\nu}) \quad (165)$$

where  $u^\alpha$  is the 4-velocity of the substrate's local rest frame, and  $\zeta$  is the universal damping invariant. This construction ensures that dissipation is proportional to the rate of change of the deformation field as measured in the medium's own rest frame.

The total action for the system is  $S_{\text{total}} = \int (\mathcal{L}_{\text{QEG}} - \mathcal{R}) \sqrt{-g} d^4x$ . The principle of least action,  $\delta S_{\text{total}} = 0$ , leads to modified Euler-Lagrange equations. The variation of the dissipative term yields a force density term proportional to the covariant derivative of  $\partial \mathcal{R} / \partial (\nabla_\alpha \mathcal{G}_{\mu\nu})$ . In the substrate's rest frame ( $u^\alpha = (1, \vec{0})$ ), this adds the term  $-\zeta \partial_t \mathcal{G}_{ij}$  to the spatial components of the field equations.

As a result, applying the principle of dissipation in a covariant framework, linearizing around equilibrium and neglecting nonlinearities of  $V$ , we obtain the *Telegrapher's equation*, a master equation that governs the full dynamics of the thermo-entropic field:

$$\tau \partial_t^2 \mathcal{G}_{ij} + \partial_t \mathcal{G}_{ij} - D \nabla^2 \mathcal{G}_{ij} = \frac{1}{\zeta} J_{ij}, \quad (166)$$

with relaxation time  $\tau = \kappa/\zeta$  and diffusivity  $D$  defined by the propagation speed  $v_*$  such that  $D = v_*^2 \tau$ . As calibrated from the modal hierarchy (Sec. XVI), the characteristic propagation speed for this slow, dissipative mode is  $v_* = 1/c$ . This is the most general linear dynamics compatible with the QEG constitutive law and reciprocity principle.

The associated dispersion relation reads

$$\tau \omega^2 + i\omega - Dk^2 = 0. \quad (167)$$

Its structure guarantees that the field exhibits two complementary physical regimes.

#### D. Regime I: High-Frequency Elastic Response (Wave-like)

When  $\omega \gg 1/\tau$ , the inertial term dominates. Equation (166) reduces to a damped wave equation. The

This completes the derivation of a Maxwell-like system for the gravito-entropic sector. The structural analogy with electromagnetism, now established from first principles, is summarized below:

Quantity	Electromagnetism	Gravito-entropism
Source	Electric charge ( $q$ )	Mass ( $m$ )
Main field (circulatory)	$\vec{B} = \frac{\mu_0 \vec{I}}{2\pi r} \hat{\theta}$	$\vec{\mathcal{T}} = \frac{k_B \vec{I}_m}{2\pi r} \hat{\theta}$
Derived field (radial)	$\vec{E} = \frac{K_e Q}{r^2} \hat{r}$	$\vec{g} = \frac{GM}{r^2} \hat{r}$
Coupling constant	$\mu_0$	$k_B = \mu_0/c^2$
Gauss's law	$\nabla \cdot \mathbf{E} = 4\pi K_e \rho_q$	$\nabla \cdot \mathbf{g} = -4\pi G \rho_m$
No-monopole law	$\nabla \cdot \mathbf{B} = 0$	$\nabla \cdot \mathcal{T} = 0$
Faraday's law	$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$	$\nabla \times \mathbf{g} = -\frac{\partial \mathcal{T}}{\partial t}$
Ampère-Maxwell law	$\nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t}$	$\nabla \times \mathcal{T} = k_B \mathbf{J}_m + k_B \epsilon_T \frac{\partial \mathbf{g}}{\partial t}$

TABLE II. Comparison of Maxwell-like laws for electromagnetism and the proposed gravito-entropic sector.

dispersion relation becomes

$$\omega^2 \simeq \frac{D}{\tau} k^2 - i \frac{\omega}{\tau}, \quad (168)$$

with phase velocity  $v_* = \sqrt{D/\tau}$ . From the modal hierarchy (Sec. XVI), this is fixed to  $v_* = 1/c$ . Thus, at high frequency,  $\Theta$  and  $\Sigma_{ij}$  behave as propagating *entropic waves*.

*Maxwell-like formulation.* We define a thermo-entropic potential  $\Theta_\mu$  and its field strength

$$H_{\mu\nu} \equiv \partial_\mu \Theta_\nu - \partial_\nu \Theta_\mu. \quad (169)$$

This satisfies  $dH = 0$  identically, giving the homogeneous equations (source-free entropic analogues of Maxwell's). The inhomogeneous equations follow from the action

$$S_{\text{GE}} = \int d^4x \sqrt{-g} \left( -\frac{1}{4k_B} H_{\mu\nu} H^{\mu\nu} - \Theta_\mu J_S^\mu \right), \quad (170)$$

leading to

$$\nabla_\mu H^{\mu\nu} = k_B J_S^\nu, \quad (171)$$

where  $J_S^\mu$  is the entropy current. The coupling is fixed by QEG's constitutive law as  $k_B = \mu_0/c^2$  (Sec. XVI C). This reproduces a complete Maxwell-like system for the gravito-entropic sector.

*Notes:*

- The antisymmetric construction  $H = d\Theta$  is necessary to guarantee the Bianchi identity  $\nabla_{[\lambda} H_{\mu\nu]} = 0$ , which in turn implies the two homogeneous Maxwell-like equations. This is not a free choice, but the unique way to construct a gauge-invariant field from a 1-form potential while preserving covariance and antisymmetry.
- The normalization with  $1/4k_B$  is fixed, not chosen: in Sec. XVI C it was shown that  $k_B = \mu_0/c^2$  is the constitutive constant of the thermo-entropic modes. Therefore, it must appear as the coupling here, ensuring consistency with the universal constitutive law and the synthesis of constants checked in Sec. XIX.

Here,  $\vec{I}_m$  and  $\rho_m$  represent the effective mass current and density.

The constitutive constants for each sector— $(\mu_0, \varepsilon_0)$  for electromagnetism and  $(k_B, \varepsilon_T)$  for gravito-entropism—are not independent parameters but distinct projections of the substrate’s universal properties  $(\kappa, \zeta)$ , constrained by the relation  $k_B \varepsilon_T = \mu_0 \varepsilon_0 = 1/c^2$ , which is necessary to ensure a universal propagation speed for wave-like excitations (see also VIB).

A profound consequence of this framework is that the thermo-entropic compliance is the direct inverse of the baseline electromagnetic inertia,  $\varepsilon_T = \frac{1}{k_B \cdot c^2} = 1/\mu_0$ . This result reveals a fundamental symmetry between the substrate’s inertial and compliant properties. Physically,  $\mu_0$  represents the baseline inductive inertia of the vacuum (its resistance to changes in torsional flux), while  $\varepsilon_T$  represents its thermo-entropic capacitive compliance (its capacity to permit a radial deformation). The relation  $\varepsilon_T = 1/\mu_0$  therefore signifies that *the compliance of the thermo-entropic mode is the exact inverse of the baseline inertia of the electromagnetic mode*. This is not an isolated coincidence but a direct manifestation of the **Dynamic Principle of Modal Reciprocity** postulated at the foundation of this theory (VIB). It is the same principle that governs the relationship between the gravitational and electromagnetic sectors, where the vacuum’s compliance to longitudinal deformation (proportional to  $G$ ) is the reciprocal of its immense stiffness against transverse deformation (proportional to  $K_e$ ). Thus, the vacuum’s elastic properties are shown to be deeply interconnected, with the capacity to yield in one mode being intrinsically linked to its resistance in another.

This reveals a non-trivial prediction: while the primary coupling to sources scales differently between the modes ( $k_B$  vs.  $\mu_0$ ), the coefficient of the displacement current term ( $k_B \varepsilon_T$  and  $\mu_0 \varepsilon_0$ ) is a *modal invariant, equal to  $1/c^2$  in both sectors*. This implies that the efficiency with which the substrate converts a changing radial field ( $\vec{E}$  or  $\vec{g}$ ) into a circulatory field ( $\vec{B}$  or  $\vec{T}$ ) is a universal mechanism of the vacuum, independent of the specific interaction.

*Einstein-like formulation.* In this propagating regime, the entropic field also contributes to the effective stress-energy. The action quadratic in  $H_{\mu\nu}$  leads to a stress-energy tensor

$$T_{\mu\nu}^{(\text{GE})} = \frac{1}{k_B} \left( H_{\mu\alpha} H_{\nu}^{\alpha} - \frac{1}{4} g_{\mu\nu} H_{\alpha\beta} H^{\alpha\beta} \right). \quad (172)$$

Coupling this to gravity via the effective Einstein equations (Sec. VIII) gives

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G \left( T_{\mu\nu} + T_{\mu\nu}^{(\text{GE})} \right). \quad (173)$$

Thus, the thermo-entropic field enters Einstein’s equations on the same footing as electromagnetic fields, but with coupling fixed by  $k_B$ .

$\zeta$  *Note:* This form is inevitable: it follows from varying the action  $S_{\text{GE}}$  with respect to the metric  $g^{\mu\nu}$ . By the standard definition  $T_{\mu\nu} = -(2/\sqrt{-g}) \delta S / \delta g^{\mu\nu}$ , the structure of  $T_{\mu\nu}^{(\text{GE})}$  is uniquely determined. Thus, the contribution of the thermo-entropic field to Einstein’s equations is forced by the variational structure, not assumed.

## E. Regime II: Low-Frequency Overdamped Response (Diffusive)

When  $\omega \ll 1/\tau$ , the viscous term dominates and (167) reduces to

$$\omega \simeq -iDk^2, \quad (174)$$

the hallmark of diffusion. The tensorial fields obey:

### 1. Heat Equation (scalar mode):

$$(u^\mu \partial_\mu) \Theta - D \nabla_s^2 \Theta = \frac{1}{\zeta'} J_\Theta. \quad (175)$$

### 2. Viscous Stress Relaxation (tensorial mode):

$$(u^\mu \partial_\mu) \Sigma_{ij} - \nu \nabla_s^2 \Sigma_{ij} = \frac{1}{\zeta''} \sigma_{ij}. \quad (176)$$

These are covariant forms of relativistic heat and Navier-Stokes-like equations. Although these equations are parabolic and appear to break relativistic symmetry, they are in fact the low-frequency approximations of the full hyperbolic system (166). Causality is preserved because the master telegrapher equation propagates with finite velocity. This mirrors the passage from causal relativistic hydrodynamics (Israel-Stewart) to Fourier’s law in the quasistatic limit [49–52].

## F. The Geometrization of Thermodynamics using Hooke’s Law and Fourier’s Law

The diffusive dynamics for the scalar mode  $\Theta$  derived in Eq. (175) provide a direct microscopic origin for the phenomenological laws of thermodynamics. This equation is the covariant generalization of *Fourier’s Law of heat conduction*. This identification allows us to interpret the abstract quantities of our model in physical terms. By comparing  $-\nabla^2 \Theta \propto J_\Theta$  with Fourier’s Law ( $\vec{q} = -k \nabla T$ ), we can identify the scalar field  $\Theta$  with the local temperature and its gradient as the driver of heat flux. Furthermore, within the collapsed dimensional framework of QEG, the scalar potential whose gradient generates this flow can be identified with *entropy*,  $S$ . Thus, entropy is geometrized as the scalar potential of the thermo-entropic field.

We begin with the standard form of Fourier’s law in vector notation:

$$\vec{q}_A = -k \nabla T, \quad (177)$$

where  $\vec{q}_A$  is the heat flux per unit area [ $\text{W m}^{-2}$ ],  $k$  is the thermal conductivity [ $\text{W m}^{-1} \text{K}^{-1}$ ], and  $\nabla T$  is the temperature gradient [ $\text{K m}^{-1}$ ]. To match the dimensional structure of other fields in the unified elastic formalism, we multiply each side by a characteristic length  $L$ , thereby defining a *line-integrated heat flux field*:

$$\vec{q} \equiv \vec{q}_A \cdot L, \quad (178)$$

which now carries units of power per unit length [ $\text{W m}^{-1}$ ], consistent with the other vector fields in our framework. Then, Fourier’s law becomes:

$$\vec{q} = -k \nabla T \cdot L = -\nabla P. \quad (179)$$

where  $\nabla P$  is the power gradient [ $\text{W m}^{-1}$ ]. Here,  $P$  is interpreted as a scalar power potential whose gradient drives thermal energy flow, just as  $V$  and  $\Phi$  generate electric and gravitational fields, respectively. However, note that *in this framework, the scalar quantity  $P$  plays a role that is conceptually indistinguishable from entropy  $S$ : it quantifies the internal deformation of the vacuum associated with thermal processes.* Indeed, since both  $P$  and  $S$  are dimensionally equivalent in the collapsed vacuum-elastic formalism, and since they both act as sources of thermodynamic flow when modulated by temperature, *we can regard entropy as the natural scalar field driving thermo-entropic deformation.*

Thus, from the standpoint of vacuum elasticity, entropy becomes a geometrically grounded scalar field, whose gradients yield observable thermal forces and fluxes. The identification of  $S$  with the scalar potential of the thermo-entropic field completes its structural analogy with gravitation and electromagnetism.

In exact parallel with the voltage  $\mathcal{E} = \int \vec{E} \cdot d\vec{\ell}$  of electrostatics, the thermo-entropic field defines a *thermo-entropic electromotive force*

$$\mathcal{E}_T \equiv \int_{\ell} \vec{q} \cdot d\vec{\ell} \quad (180)$$

with units of power [W]. In the dimension-collapsed vacuum-elastic system adopted here, powers are dimensionless, so  $\mathcal{E}_T$  plays exactly the same algebraic role as an (dimensionless) electric EMF.

Given the dimensional equivalence between power and entropy in the vacuum-elastic unit system, the quantity  $\mathcal{E}_T$  can also be interpreted as a net entropy difference between two thermal regions. Thus, the thermo-motive force becomes not only a measure of energy flux, but also a geometric manifestation of entropic imbalance in the vacuum lattice—driving deformation analogously to how electric potential drives charge.

As a result, each classical field admits a corresponding integral expression that encodes its action over an extended region. The following table summarizes these correspondences:

Phenomenon	Field	Expression	Physical meaning
Heat	$\vec{q}$	$\int \vec{q} \cdot d\vec{A}$	Thermal flux
Electricity	$\vec{E}$	$\int \vec{E} \cdot d\vec{A}$	Electric flux
Magnetism	$\vec{B}$	$\int \vec{B} \cdot d\vec{A}$	Magnetic flux
Gravity	$\vec{g}$	$\int \vec{g} \cdot d\vec{A}$	Gravitational flux

If we denote by  $\Delta T$  the temperature differential between adjacent isothermal layers, we can express the thermo-entropic response in a strictly Hookean form:

$$S = \kappa_{\text{dyn}} \Delta T \quad (181)$$

Here,  $S$  represents the entropy associated with the deformation of the vacuum medium, and  $\kappa_{\text{dyn}}$  acts as a dynamic entropic stiffness with dimensions [ $L^{-1}$ ] in the vacuum-elastic unit system. This formulation reinforces the interpretation of entropy as a scalar elastic displacement field, and  $\Delta T$  as its driving cause.

To provide a more concrete physical picture, we can interpret this thermal deformation within the oscillator lattice model. While a coherent, compressional deformation corresponds to mass and a coherent, torsional one corresponds to charge, we can identify temperature ( $T_{\text{emp}}$ ) with the *incoherent, isotropic vibrational energy* of the lattice oscillators in a given region. In this view, temperature is a measure of the average squared amplitude of these random, uncoordinated oscillations. A temperature gradient ( $\nabla T$ ) is thus a gradient in the intensity of this background 'shimmering' of the vacuum, which naturally drives a net flow of energy (heat) from regions of high-amplitude vibration to regions of low-amplitude vibration, perfectly aligning with the phenomenological description of Fourier's Law.

Hookean elasticity	Thermo-entropic
$F$ (mechanical force)	$P$ (power) / $S$ (entropy)
$\Delta x$ (displacement)	$\Delta T$ (Temperature diff.)
$k$ (spring constant, [ $L^{-1}$ ])	$\kappa_{\text{dyn}}$ ([ $L^{-1}$ ] or $\text{W s}^{-1}$ )

As a result, the thermo-entropic field integrates consistently into our unified tensorial framework as a Hookean deformation mode, whose force-like response is power and whose generalized displacement is temperature difference.

### G. Stability and Causality

Equation (166) guarantees both causal propagation and Lyapunov stability:

$$\frac{d\mathcal{E}}{dt} = -\frac{\zeta}{2} \int d^3x |\partial_t \mathcal{G}_{ij}|^2 \leq 0, \quad (182)$$

with finite signal velocity  $v_* = 1/c$ . This aligns with the causal relativistic hydrodynamics of Israel–Stewart type. The monotonic decrease of the energy functional follows directly from the viscous term, which is strictly positive. This constitutes a Lyapunov stability proof and guarantees consistency with the Second Law of Thermodynamics: entropy production is non-negative.

### H. Conceptual Closure

The thermo-entropic field is not a speculative add-on but an inevitable consequence of the QEG framework:

- In the **elastic regime**, it manifests as a Maxwell-like system with propagating entropic waves, contributing to Einstein-like field equations via  $T_{\mu\nu}^{(\text{GE})}$ .
- In the **dissipative regime**, it obeys covariant heat and stress-relaxation equations, providing a microscopic origin for irreversibility.

This duality resolves deep puzzles:

- The **Arrow of Time**: emerges from the intrinsic diffusive regime of the field.
- The **Cosmic Isotropy**: anisotropies are naturally damped by spacetime viscosity.
- The **Dark Energy problem**: the isotropic  $\Theta$  field contributes a uniform thermal background energy density.

Thus, QEG not only recovers Einstein's and Maxwell's theories, but also extends them, predicting a new fundamental field of spacetime: the thermo-entropic field.

## I. Summary of structural analogies between elastic responses of the vacuum

We present below the structural correspondence between the linearized Fourier's law, Gauss's law in electrostatics, and Newton's law of gravitation:

Concept	Fourier's L.	Gauss's L.	Newton's L.
Source	$\nabla \cdot \vec{q}$	$\nabla \cdot \vec{E}$	$\nabla \cdot \vec{g}$
Potential	$P \equiv S$	$V$	$\Phi$
Field	$\vec{q} = -\nabla S$	$\vec{E} = -\nabla V$	$\vec{g} = -\nabla \Phi$
Poisson eq.	$\nabla^2 S = k_B \rho_{\text{temp}}$	$\nabla^2 V = -\frac{\rho_e}{\epsilon_0}$	$\nabla^2 \Phi = 4\pi G \rho_m$

where  $\rho_{\text{temp}}$  represents a localized temperature density or distribution acting as the source of entropic deformation. Each of these field laws describes how a scalar potential gives rise to a vector field through a gradient operation, and how the divergence of that field connects to a source density via a Poisson-type equation. In this structural analogy:

- The entropy  $S$  plays the role of a scalar thermal potential,
- The linearized heat flux  $\vec{q}$  is analogous to the electric field  $\vec{E}$  or the gravitational field  $\vec{g}$ ,
- And the Laplacian  $\nabla^2 S$  captures thermo-entropic curvature, in full parallel with electrostatic and gravitational curvature.

The thermo-entropic field, driven by entropy gradients and governed by thermal curvature, thereby integrates as a scalar deformation mode within the elastic manifold defined by the symmetric tensor  $\mathcal{G}_{\mu\nu}$ . This geometrization of heat completes the triad of scalar sources—mass, charge, and temperature—unifying their corresponding field interactions as elastic responses of the vacuum encoded in the symmetric tensor  $\mathcal{G}_{\mu\nu}$ .

## XXI. VALIDATION AND DEEPER IMPLICATIONS OF THE THERMO-ENTROPIC FIELD

### A. Fundamental parameters of the thermo-entropic field

Applying the universal scaling rule derived in XVII B, we can derive the fundamental parameters of the thermo-entropic field just dividing the parameters obtained for the electromagnetic field by  $c^2$ :

- We obtain the quantum of mass-energy for the thermo-entropic field  $m_{\text{entr}} = \frac{\hbar}{2\pi c \cdot 1 \text{ m}} \approx 5.6 \times 10^{-44} \text{ kg}$ .
- We obtain a quantum of mass density  $\rho_{\text{entr}} = \frac{\hbar}{2\pi c \cdot 1 \text{ m}^4} \text{ kg m}^{-3}$ .
- We can set the action as  $\frac{\hbar}{c^2}$ , which matches the result derived previously (XVIC).

### B. Boltzmann's Constant as the Fundamental Quantum of Thermo-Entropic Force

Building on the reinterpretation of  $k_B$  as a thermo-entropic force (see Sec. XVIC) and the fundamental

equivalence introduced earlier (Eq. 117), we now propose a novel formulation of the Boltzmann constant as an emergent relativistic force.

Assuming the dimensional equivalence  $1 \text{ K} \equiv 1 \text{ m} \equiv 1 \text{ s}$  within our elastic vacuum framework, we express  $k_B$  as:

$$k_B = \frac{\hbar c}{2 \cdot 1 \text{ m}} \cdot \frac{1}{2\alpha \cdot 1 \text{ s}} = \frac{E}{a} \quad (183)$$

In this expression,  $k_B$  acquires the form of a Newtonian-like force  $F = ma$ , with the following components:

- $\frac{\hbar c}{1 \text{ m}} \rightarrow m$ : The characteristic energy scale of the vacuum, associated with a fundamental quantum of mass or photon energy.
- $\frac{1}{2\alpha \cdot 1 \text{ s}} = \frac{\gamma}{1 \text{ s}} \rightarrow a$ : An effective proper acceleration, where the Lorentz-like factor  $\gamma = \frac{1}{2\alpha}$  encodes the vacuum's resistance to excitation.

Thus, the Boltzmann constant  $k_B$  emerges as a quantized force scale, representing the vacuum's intrinsic responsiveness to acceleration. In this interpretation, entropy and temperature arise from the inertial resistance of spacetime to deformation, with  $k_B$  capturing the proportionality between energetic input and induced entropic curvature.

This Unruh-inspired formulation reinforces the view that thermodynamic quantities—such as temperature, and heat—are fundamentally geometric in nature. Here,  $k_B$  bridges the gap between thermal response and relativistic motion, playing a role analogous to that of  $G$  or  $\mu_0$  in mediating the vacuum's reaction to mass or charge, respectively. In this sense,  $k_B$  can be viewed as the *thermo-entropic stiffness constant* of spacetime: a universal coupling between acceleration, information flow, and thermal excitation. This perspective helps unify quantum field theory, thermodynamics, and general relativity within a common elastic-dynamical substrate.

### C. Deriving Unruh effect from the thermo-entropic field parameters and Newton's law

There are several checks that we can perform to further justify the validity of the fundamental parameters derived for the thermo-entropic field. In this subsection, we will focus on showing that the Unruh effect can be directly derived from the application of the fundamental expression of mass-energy for the thermo-entropic field and Newton's Law.

The **Unruh effect** [53] states that an observer with constant proper acceleration  $a$  in vacuum perceives a thermal bath at a temperature

$$T_{\text{Unruh}} = \frac{\hbar a}{2\pi c k_B}. \quad (184)$$

Rearranging gives

$$k_B = \frac{\hbar a}{2\pi c T_{\text{Unruh}}}. \quad (185)$$

We have already shown how  $k_B$  can be treated dimensionally as a force (XVIC). Also, we have derived that the expression of mass-energy for the thermo-entropic field is  $m_{\text{entr}} = \frac{\hbar}{2\pi c \cdot \lambda}$  XXI A. By identifying the force

$F$  in Newton's second law with the Boltzmann constant  $k_B$  (as justified by the dimensional equivalence framework, see XVI C) and the characteristic length  $\lambda$  with the Unruh temperature  $T_{Unruh}$ , the Unruh effect emerges as the direct application of Newtonian dynamics to the derived properties of the field:

$$F = k_B = m_{entr} \cdot a = \frac{\hbar}{2\pi c T_{Unruh}} \cdot a \quad (186)$$

This rearranges to the exact Unruh formula.

#### D. The Common Origin of Thermal Radiation and Quantum Forces

A powerful consistency check of the unified framework arises from revealing the deep connection between two seemingly unrelated phenomena: blackbody thermal radiation, governed by the Stefan-Boltzmann constant ( $\sigma$ ), and quantum intermolecular forces, governed by the London dispersion coefficient ( $C_6$ ). We will demonstrate that their dimensional correspondence is not a coincidence, but a necessary consequence of their common origin in the fluctuations of the quantum vacuum.

##### *The Dimensional Signature of Vacuum-Mediated Interactions*

London dispersion forces are a direct manifestation of the quantum vacuum's activity. They arise from the interaction of transient dipoles induced by zero-point fluctuations of the electromagnetic field. For two neutral atoms, the interaction potential is:

$$U_{\text{London}}(r) = -\frac{C_6}{r^6}$$

where the  $C_6$  coefficient encapsulates the polarizability of the atoms and is fundamentally determined by the structure of vacuum fluctuations. In our unified dimensional framework, where energy has dimensions of length ( $[E] \equiv [L]$ ), the London coefficient carries dimensions:

$$[C_6] = [E] \cdot [L^6] \equiv [L^7].$$

Now, consider the Stefan-Boltzmann constant, whose classical expression is built from the fundamental constants of quantum mechanics, thermodynamics, and relativity:

$$\sigma = \frac{\pi^2 k_B^4}{60 \hbar^3 c^2}.$$

Within our dimensional system, where  $k_B$ ,  $\hbar$ , and  $c$  reduce to combinations of lengths, the Stefan-Boltzmann constant acquires:

$$[\sigma] = [\text{W m}^{-2} \text{K}^{-4}] \equiv [L^{-6}].$$

This remarkable outcome reveals that  $\sigma$  shares the inverse sixth-power length dependence characteristic of dipole-dipole vacuum interactions—a dimensional signature of shared physical origin. To formalize the connection, we define:

$$[\sigma \cdot C_6] = [L^{-6}] \times [L^7] = [L],$$

which is the dimension of energy in our framework. The product  $\sigma C_6$  thus defines an intrinsic energy scale of the vacuum, suggesting that  $\sigma$  and  $C_6$  are complementary macroscopic parameters arising from the same vacuum energy reservoir.

##### *Physical Interpretation: Thermal vs. Ground-State Excitations of the Vacuum*

This dimensional correspondence can be physically interpreted through the modal structure of quantum field theory. Both blackbody radiation and Casimir-London forces derive from the quantization of the electromagnetic field modes within boundary conditions:

- *London and Casimir forces* emerge from the *ground-state energy* (zero-point energy) of these modes, typically expressed as  $E_0 = \frac{1}{2} \hbar \omega$ . They represent mechanical stresses exerted by the vacuum in its lowest energy configuration.
- *Stefan-Boltzmann radiation*, in contrast, arises from the *thermal excitation* of the same set of vacuum oscillators. The Stefan-Boltzmann law captures the total energy flux when these modes are thermally populated according to Planck's distribution.

Thus, London forces and blackbody radiation are not fundamentally distinct; they are two manifestations of the same quantized vacuum structure—one probing the ground state, the other the thermal state. The constants  $C_6$  and  $\sigma$  parameterize these effects at the macroscopic level, but their shared dimensionality and combined energy scaling ( $\sigma C_6$ ) highlight their unified origin. This perspective reinforces the central thesis of this work: that all physical interactions emerge as distinct modal responses of a single, elastic quantum vacuum.

##### *Implications for the Unified Description of Vacuum Energy and Thermodynamics*

This unification between thermal radiation and quantum dispersion forces provides a profound bridge between macroscopic thermodynamics and microscopic quantum interactions. The vacuum behaves as an elastic medium whose modal excitations, whether thermal or mechanical, dictate both the radiation laws and intermolecular forces. Accordingly, the Stefan-Boltzmann constant encapsulates not just an empirical radiation law, but a thermodynamic response of the vacuum's elastic structure. The London coefficient reflects the mechanical manifestation of the same vacuum under ground-state conditions. This duality strengthens the proposal that space-time itself possesses elastic properties, with its interaction with vacuum fluctuations giving rise to all observed forces and thermodynamic behaviors.

## XXII. SYNTHESIS: GRAVITY AS AN EMERGENT PHENOMENON OF VACUUM THERMODYNAMICS

The internal consistency of the QEG framework culminates in a profound synthesis that quantitatively defines the relationship between gravity and thermodynamics. By combining two of the theory's core, independently derived results—the dynamic origin of the gravitational constant and the principle of causal universality—the nature of gravity is revealed as an emergent property of the vacuum's thermodynamic response.

The starting points are two foundational equations of this theory:

1. **The Dynamic Origin of G:** As established in Sec. VI, gravity emerges as a second-order, dissipative effect of the vacuum's primary stiffness, yielding the relation  $G = \mu_0 \alpha^2$ .
2. **The Dynamic Reciprocity Condition:** As a direct consequence of requiring a universal propagation speed  $c$ , we have derived the thermo-entropic compliance  $\varepsilon_T = 1/\mu_0$  (see Sec. XX). This can be rewritten as  $\mu_0 = 1/\varepsilon_T$ .

Substituting the second relation into the first yields a new, fundamental expression for the gravitational constant:

$$G = \frac{\alpha^2}{\varepsilon_T} \quad (187)$$

The implications of this equation are remarkable and provide a first-principles basis for the emergent gravity paradigm. It states that the gravitational constant  $G$  is not a fundamental parameter of nature, but is instead determined by the ratio of two more fundamental properties of the vacuum:

- It is directly proportional to the square of the vacuum's intrinsic **dissipation**, quantified by the fine-structure constant ( $\alpha^2$ ).
- It is inversely proportional to the vacuum's **thermo-entropic compliance** (its capacity to permit thermal/entropic deformations), quantified by  $\varepsilon_T$ .

This formulation makes two powerful, falsifiable predictions about the nature of gravity:

1. If the vacuum were a perfect, non-dissipative medium ( $\alpha \rightarrow 0$ ), then gravity as we know it would not exist ( $G \rightarrow 0$ ). The existence of gravity is therefore intrinsically linked to the irreversibility and inherent "friction" of spacetime.
2. If the vacuum were infinitely "soft" or compliant to entropic deformations ( $\varepsilon_T \rightarrow \infty$ ), gravity would be infinitely weak ( $G \rightarrow 0$ ). The strength of gravity is therefore a direct measure of the vacuum's "stiffness" against being deformed thermodynamically.

In conclusion, Eq. (187) positions gravity not as a primary interaction, but as a residual, second-order phenomenon emerging from the thermodynamics of the vacuum's quantum-elastic substrate. It provides a concrete, quantitative mechanism for the ideas proposed by Jacobson and Verlinde, explicitly linking Newton's constant to the thermodynamic and dissipative properties that govern the fabric of spacetime itself.

## Part VI: Validation and Cosmological predictions of the quantum elastic geometry

### XXIII. VALIDATION: THE UNIVERSAL FIELD STRUCTURE AND ITS FUNDAMENTAL SOURCES

A cornerstone of the QEG framework is the prediction, derived from the substrate's universal Laplacian response (Sec. II E), that all static fields must adhere to a common structure. Given that fields in QEG have

dimensions of inverse length, [ $L^{-1}$ ], this universal form can be expressed as:

$$\vec{\Phi}_X(r) = C_X \cdot \frac{1}{4\pi r} \cdot \hat{e}_X \quad (188)$$

where  $1/(4\pi r)$  is the universal elastic modulus of the vacuum, and  $C_X$  is a *dimensionless coefficient* that represents the strength (potential) of the source for the mode  $X$ . In this section, we validate this prediction by demonstrating that the standard laws of physics, when rewritten using QEG identities, collapse precisely into this form. Crucially, we will show that the source coefficients  $C_X$  are not arbitrary, but are directly related to the fundamental quanta of energy and action defined in the *Vacuum's Constitutive Equation* 117:

$$\mu_0 \cdot e \equiv k_B \cdot 2\alpha \equiv \frac{hc}{1 \text{ m}}$$

#### A. Justification of the Fundamental Source: The Primacy of the Quantum Harmonic Oscillator

A critical step in our validation is to establish, from first principles, the fundamental quantum source for each interaction. We will now demonstrate that the choice of these sources is not arbitrary, but is a necessary consequence of the central role of the Quantum Harmonic Oscillator (QHO) as the foundational building block of all stable, coherent excitations in nature.

*a. The QHO as a Consequence of First Principles.* The fundamental equation of non-relativistic quantum dynamics is the Schrödinger equation. For any system in a stable equilibrium, the potential energy  $V(x)$  can be Taylor-expanded around its minimum. The first non-trivial term is always quadratic,  $V(x) \approx \frac{1}{2}kx^2$ , the harmonic potential. This is not a choice, but a mathematical inevitability for small perturbations around a stable point. The QHO, as the solution to the Schrödinger equation with this potential, thus represents the *structural universal model for any minimal, stable quantum excitation*. It is the "atom" of quantum perturbation.

*b. The Irreducible Quantum of Energy.* The quantization of the QHO yields the famous discrete energy levels,  $E_n = (n + 1/2)\hbar\omega$ . The lowest possible energy state ( $n = 0$ ) is the non-zero ground state, or *Zero-Point Energy*:

$$E_0 = \frac{1}{2}\hbar\omega \quad (189)$$

This  $E_0$  is the *irreducible, non-zero quantum of coherent energy* that a stable vacuum fluctuation (the building block of an endogenous source) can possess. In our framework, we identify this as the fundamental source quantum for interactions mediated by stable, coherent particles. For a massless mode of characteristic wavelength  $\lambda$ , this energy is  $E_0 = \hbar c/(2\lambda)$ .

*As a result, the most fundamental unit of the unified field must be this irreducible quantum of coherent energy,  $E_0$ .*

*c. From Fundamental Energy to Primordial Source Flux.* In an elastic medium governed by a Laplacian response, the relationship between a source density  $\rho$  and the field it generates,  $\vec{\Phi}$ , is given by Poisson's equation,  $\nabla \cdot \vec{\Phi} = 4\pi G_0 \rho$ , where  $G_0$  is the fundamental,

dimensionless coupling constant of the undressed substrate. We set  $G_0 = 1$  for this baseline interaction. The total flux emerging from a source is found by integrating over a volume enclosing it, via Gauss's Law:

$$\text{Flux} = \oint_S \vec{\Phi} \cdot d\vec{A} = \int_V (\nabla \cdot \vec{\Phi}) dV = \int_V 4\pi\rho dV \quad (190)$$

For our fundamental point-like source, we identify its total content with the irreducible quantum of coherent energy,  $\int_V \rho dV = E_0$ . The primordial flux, which we identify as the fundamental source strength  $S_f$ , is therefore:

$$S_f = 4\pi \cdot E_0 = 4\pi \left( \frac{\hbar c}{2\lambda} \right) = \frac{\hbar c}{\lambda} \quad (191)$$

At the fiducial scale of our framework, we set  $\lambda = 1 \text{ m}$ . This yields the fundamental source flux for any mode:

$$S_f = \frac{\hbar c}{1 \text{ m}} \quad (192)$$

*d. A Powerful Consistency Check.* This result is profound. The primordial source flux for the unified field, derived here from the first principles of quantum mechanics (Schrödinger Eq.  $\rightarrow$  QHO  $\rightarrow$  Zero-Point Energy) and geometry (Poisson's Eq.), is *identically one of the three terms in the Vacuum's Constitutive Equation*:

$$\mu_0 \cdot e \equiv k_B \cdot 2\alpha \equiv \frac{\hbar c}{1 \text{ m}}$$

This demonstrates that our choice for each mode sources is not arbitrary. It is the *unique* choice that is both fundamentally derived from the nature of quantum excitations and perfectly consistent with the overarching structure of interconnected constants in the QEG framework.

## B. Verification of the Universal Structure

*a. 1. The Electric Field ( $\vec{E}$ ).* The standard expression is Coulomb's Law,  $\vec{E} = \frac{1}{4\pi\epsilon_0} \frac{e}{r^2} \hat{r}$ . This already matches the universal form proposed. Using the identity  $1/\epsilon_0 = \mu_0 c^2$ , we have:

$$\vec{E} = \left( \frac{e}{\epsilon_0 \cdot r} \right) \cdot \frac{1}{4\pi r} \hat{r} = \left( \frac{\mu_0 \cdot e \cdot c^2}{r} \right) \cdot \frac{1}{4\pi r} \hat{r} \quad (193)$$

Using the identity  $e \equiv 2\alpha \cdot \frac{1 \text{ m}}{c^2}$ , we can re-express it as:

$$\vec{E} = \left( \frac{2\alpha \cdot \mu_0 \cdot 1 \text{ m}}{r} \right) \cdot \frac{1}{4\pi r} \hat{r} \quad (194)$$

where  $\mu_0 \cdot 1 \text{ m}$  is the vacuum's base inductance and  $2\alpha$  is the dissipative term.

*b. 2. The Gravitational Field ( $\vec{g}$ ).* The standard expression is Newton's Law,  $\vec{g} = \frac{G_N m}{r^2} \hat{r}$ . Using the QEG identities for the local coupling,  $G_N \equiv G_{\text{loc}} = \mu_0 \alpha^2$ , and for the fundamental mass quantum,  $m = \hbar c / (2 \cdot 1 \text{ m}) = \frac{\hbar c}{4\pi \cdot 1 \text{ m}}$ , we derive:

$$\vec{g} = \frac{G_N m}{r^2} \hat{r} = \left( \frac{\mu_0 \alpha^2 \cdot \frac{\hbar c}{1 \text{ m}}}{r} \right) \cdot \frac{1}{4\pi r} \hat{r} \quad (195)$$

This expression expresses the microscopic sourcing of the gravitational field as a function of the vacuum's baseline potential ( $\mu_0$ ) quadratically suppressed by the universal damping factor ( $\alpha^2$ ), which rigorously explains its extreme weakness relative to other forces.

*c. 3. The Magnetic Field ( $\vec{B}$ ).* The fundamental expression from a point-like stationary current is  $\vec{B} = \frac{\mu_0 I}{4\pi r} \hat{\theta}$ <sup>7</sup>. This already matches the universal form. The source is the current  $I$ . For the electromagnetic mode, the characteristic current is the speed of light,  $I = c$ . Since  $c$  is dimensionless in QEG, we have

$$\vec{B} = \frac{\mu_0 I}{4\pi r} \hat{\theta} = \mu_0 c \cdot \frac{1}{4\pi r} \hat{\theta} = Z_0 \cdot \frac{1}{4\pi r} \hat{\theta} \equiv \frac{\vec{E} \cdot c}{2\alpha} \quad (196)$$

This expression reveals the magnetic field as a direct manifestation of the vacuum's torsional properties, whose strength is governed by a single dimensionless coefficient: the vacuum impedance,  $Z_0$ . Magnetism is thus interpreted as a *measure of the substrate's intrinsic "circulatory stiffness"*—its resistance to being twisted—driven by the characteristic velocity of the electromagnetic mode.

*d. 4. The Thermo-Entropic Field ( $\vec{T}$ ).* The source of the thermo-entropic field is the fundamental quantum of thermal energy exchange, which the Vacuum's Constitutive Equation identifies as  $k_B \cdot 2\alpha$ . The field is driven by this point-like stationary source and the characteristic baseline velocity,  $I = 1 \text{ m/s}$ :

$$\vec{T} = \frac{(k_B \cdot 2\alpha) I}{4\pi r} \hat{\theta} = \frac{k_B \cdot 2\alpha}{4\pi r} \hat{\theta} \quad (197)$$

Using the QEG identity  $k_B \equiv \mu_0 / c^2$ , we cast it into the universal form:

$$\vec{T} = \frac{(\mu_0 / c^2) \cdot 2\alpha}{4\pi r} \hat{\theta} = \left( \frac{2\alpha \cdot \mu_0}{c^2} \right) \cdot \frac{1}{4\pi r} \hat{\theta} \equiv \frac{\vec{E}}{c^2} \equiv \frac{\vec{g} \cdot c}{2\alpha} \quad (198)$$

Throughout this validation, the characteristic expressions for the fundamental fields have been constructed from the Vacuum's Constitutive Equation, which reflects the elementary contribution of the potential of a single discrete quantum oscillator.

## C. Synthesis: The Hierarchy of Fields and Electro-Gravitational Equivalence

The unified structure of the fundamental fields reveals a remarkable hierarchy. The framework suggests a paradigm shift where the circulatory, azimuthal modes ( $\vec{B}, \vec{T}$ ) are primary, and the radial force fields ( $\vec{E}, \vec{g}$ ) emerge as secondary phenomena when the circulatory fields are "amplified" by a universal conversion factor:

$$\begin{aligned} \vec{E} &= \vec{B} \cdot I_{\text{max}} \\ \vec{g} &= \vec{T} \cdot I_{\text{max}} \end{aligned} \quad (199)$$

where  $I_{\text{max}} \equiv G_{\text{Glob}} = 2\alpha / c$  is the maximum vacuum current derived previously 208. This relationship can be intuitively understood through a hydrodynamic analogy, where a circulatory flow (a vortex, like  $\vec{B}$ ) in the vacuum "fluid" generates a pressure gradient that results in a radial force (like  $\vec{E}$ ).

<sup>7</sup> From the Biot-Savart law applied to a localized oscillatory mode, yielding the expression  $\vec{B} = \mu_0 I / 4\pi r$ , which better reflects the point-like, modular nature of the vacuum excitations in this framework

Crucially, this universal conversion factor,  $I_{\max}$ , is not an arbitrary constant but is determined by the vacuum's most fundamental dissipative ( $\alpha$ ) and kinematic ( $c$ ) properties. This implies that the very existence of radial static forces is intrinsically linked to the dissipative nature of the substrate. In a hypothetical, frictionless vacuum where  $\alpha = 0$ , this conversion factor would vanish, and static forces would not be generated in this manner.

#### D. The Electro-Gravitational Equivalence: A Static-Dynamic Duality of the Vacuum

The unified structure of the fundamental fields, derived from the Vacuum's Constitutive Equation, reveals a profound and rigid hierarchy between the interactions. Beyond the relationship between radial and azimuthal modes, a more direct and powerful consequence is the inescapable link between the two radial force fields themselves. We find a direct relationship between the electric and gravitational fields at the fiducial scale:

$$\frac{\vec{E}}{c^2} \equiv \frac{\vec{g} \cdot c}{2\alpha} \rightarrow \vec{E} \cdot \frac{e}{1 m} \equiv \vec{g} \cdot c \quad (200)$$

*Physical Interpretation: The Static-Dynamic Equilibrium.*

This equation establishes a remarkable *Electro-Gravitational Equivalence*. It represents a fundamental equilibrium within the vacuum substrate. The left-hand side, the electric force per unit length, quantifies the *static elastic stress* of the vacuum in response to a fundamental charge. It is a measure of the substrate's capacity to exert a static, radial force. The right-hand side, the gravitational field multiplied by the characteristic EM current, quantifies the *dynamic gravito-inertial flux*. It is a measure of the substrate's capacity for dynamic response and flow. The equivalence reveals that these two aspects of the vacuum—its static tension and its dynamic flux potential—are not independent but are two perfectly balanced facets of a single underlying reality.

*Physical Manifestation: The Photoelectric Effect*

This abstract equilibrium finds a direct and stunning physical manifestation in one of the cornerstones of quantum mechanics: the photoelectric effect. The phenomenon, where a photon liberates a bound electron from a material, can be reinterpreted as a process governed by the Electro-Gravitational Equivalence.

- **The Bound Electron and the Static Stress:** An electron within a metal is bound by electrostatic forces. The energy required to overcome this binding and extract the electron is the material's **work function**,  $W$ . This work function is the integrated measure of the *static elastic stress* of the substrate (the LHS of Eq. 200) that confines the electron to its potential well.
- **The Incident Photon and the Dynamic Flux:** An incident photon is a localized, dynamic quantum of energy,  $E_\gamma = hf$ . In QEG, this energy acts as a source for a gravitational/inertial perturbation,  $g$ , and it propagates with the characteristic current of the EM

mode,  $c$ . The term on the RHS of our equivalence,  $gc$ , therefore represents the *dynamic impulsive flux* that this quantum of energy can deliver to the substrate.

- **The Threshold Condition as the Equilibrium Point:** The famous threshold condition for photoemission,  $hf_{\min} = W$ , is the precise physical realization of our equivalence principle. It marks the exact point where the dynamic flux delivered by the photon (the RHS) is perfectly balanced by the static stress binding the electron (the LHS).

Therefore, the photoelectric effect is revealed to be more than a simple energy exchange. It is a quantum manifestation of the fundamental static-dynamic equilibrium of the spacetime substrate, where a quantum of dynamic flux is absorbed to precisely overcome a quantum of static elastic stress. This connection provides powerful, independent evidence for the validity of the QEG framework, grounding its most abstract predictions in one of the most fundamental and well-verified phenomena of quantum physics.

## XXIV. GLOBAL GRAVITATIONAL CONSTANT AS THE FUNDAMENTAL NOETHER CURRENT OF THE VACUUM

In the QEG framework, the torsional sector (encoded in the mixed components  $\mathcal{G}_{0i}$ ) realises a  $U(1)$  gauge symmetry at low energies. In this section we (i) derive the associated Noether current directly from the QEG Lagrangian, (ii) evaluate its fundamental amplitude for the fiducial vacuum cell, and (iii) prove that this amplitude coincides with the global gravitational constant of the quasi-linear cosmological regime:

$$G_{\text{Glob}} \equiv I_{\max} \quad (201)$$

This identification turns the large-scale gravitational coupling into an unavoidable consequence of the vacuum's  $U(1)$  symmetry and its conserved current.

### A. Torsional sector of QEG and the $U(1)$ gauge symmetry

We model the torsional response of the substrate by a gauge 1-form  $A_\mu$  obtained as a linear projection of the mixed components of the deformation tensor,  $A_\mu \equiv C_A \Pi_\mu^{\nu i} \mathcal{G}_{0i}$ , with a fixed projector  $\Pi$ . The gauge-invariant field strength is  $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ . The universal torsional Lagrangian density (vacuum plus minimal matter) reads

$$\mathcal{L}_{\text{tor}} = -\frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} + (D_\mu \Psi)^* (D^\mu \Psi) - U(|\Psi|) \quad (202)$$

$$D_\mu = \partial_\mu + i e A_\mu,$$

where  $\Psi$  is a complex order parameter for the coherent torsional excitation (the minimal  $U(1)$  matter content)<sup>8</sup>,  $e$  is the charge quantum, and  $\mu_0$  is the vacuum's transverse stiffness previously identified in the constitutive relations.

<sup>8</sup> Physically,  $\Psi$  represents the effective, coarse-grained field describing the coherent phase and amplitude of the underlying substrate's torsional oscillators.

a. *Gauge structure and global  $U(1)$  symmetry.* The Lagrangian (202) is invariant under local  $U(1)$  transformations  $\Psi \rightarrow e^{ie\chi(x)}\Psi$ ,  $A_\mu \rightarrow A_\mu - \partial_\mu\chi$ . Noether's second theorem enforces identities for the gauge sector. For the purpose of a *conserved charge/current*, we restrict to the global subgroup  $\chi(x) = \theta = \text{const}$ . Then the variation  $\delta\Psi = ie\theta\Psi$  yields a strictly conserved Noether current:

$$j^\mu = \frac{\partial\mathcal{L}_{\text{tor}}}{\partial(\partial_\mu\Psi)}\delta\Psi + \text{c.c.} = i\left[\Psi^*(D^\mu\Psi) - (D^\mu\Psi)^*\Psi\right]$$

$$\partial_\mu j^\mu = 0. \quad (203)$$

This current is the unique, symmetry-mandated flow associated with the torsional  $U(1)$  sector of QEG. Its spatial integral gives the conserved Noether charge  $Q = \int d^3x j^0$ , which identifies the charge quantum  $e$  for a single coherent cell.

### B. Fundamental amplitude of the vacuum Noether current

Consider the uniform, single-mode, coherent solution  $\Psi(x) = \Psi_0 e^{-i\omega t}$ , with spatially homogeneous amplitude  $|\Psi_0| = \text{const}$  and vanishing background potential  $A_\mu = 0$  (Coulomb gauge; the evaluation at  $A_\mu = 0$  fixes the intrinsic, un-driven current scale). Then  $D_\mu\Psi = (\partial_\mu + ieA_\mu)\Psi \rightarrow \partial_\mu\Psi$ , so Eq. (203) gives

$$j^0 = 2e|\Psi_0|^2, \quad \mathbf{j} = 2e|\Psi_0|^2 \mathbf{v}_{\text{ph}}, \quad \mathbf{v}_{\text{ph}} \equiv \nabla(\omega t - \mathbf{k}\mathbf{x})/\omega. \quad (204)$$

In a fiducial QEG cell of linear size  $L_{\text{ref}} = 1 \text{ m}$  with periodic boundary conditions (the same cell used to define the constitutive identities), a purely temporal mode ( $\mathbf{k} = 0$ ) has  $\mathbf{v}_{\text{ph}} = \mathbf{0}$  and carries conserved charge  $Q = \int_{V_{\text{ref}}} j^0 d^3x = 2e|\Psi_0|^2 V_{\text{ref}}$ . Fixing the elementary quantum  $Q = e$  in the fiducial cell fixes the amplitude  $|\Psi_0|^2 = \frac{1}{2V_{\text{ref}}} = \frac{1}{2}$  since  $V_{\text{ref}} = 1 \text{ m}^3$ . A small, uniform gauge oscillation  $A_0 = 0$ ,  $A_i(t) = A_i^{(0)} \cos(\omega t)$  minimally couples as  $\delta\mathcal{L} \supset j^\mu A_\mu$  and induces a spatial current density  $\mathbf{j}(t) \propto \sin(\omega t)$  with amplitude proportional to  $\omega$ . Integrating over the cell cross-section (unit area in our fiducial construction) gives the *current amplitude*

$$I_{\text{max}} = e \cdot \omega_{\text{ref}}, \quad \omega_{\text{ref}} \equiv \frac{c}{L_{\text{ref}}} = \frac{c}{1 \text{ m}}. \quad (205)$$

Using the QEG constitutive identity for the elementary charge (Sec. XVII A),  $e \equiv \frac{2\alpha \cdot 1 \text{ m}}{c^2}$ , we obtain the *inevitable* magnitude of the vacuum's fundamental Noether current:

$$I_{\text{max}} = e \omega_{\text{ref}} = \left(\frac{2\alpha \cdot 1 \text{ m}}{c^2}\right) \left(\frac{c}{1 \text{ m}}\right) = \frac{2\alpha}{c}. \quad (206)$$

No further dynamical assumptions enter: Eq. (206) follows from (i) the  $U(1)$  symmetry of the torsional sector, (ii) Noether's theorem, and (iii) the QEG constitutive identities and fiducial scale.

### C. Independent derivation of the global gravitational constant

Independently, the global gravitational constant  $G_{\text{Glob}}$  that governs the quasi-linear, homogeneous

regime is fixed by the fine-structure constant,

$$\alpha = \frac{e^2}{4\pi\epsilon_0\hbar c} \implies G_{\text{Glob}} \equiv 2\pi\epsilon_0 = \frac{e^2}{2\alpha\hbar c}. \quad (207)$$

Substituting the QEG identities  $e \equiv \frac{2\alpha \cdot 1 \text{ m}}{c^2}$  and  $\hbar \equiv \frac{1 \text{ m}^2}{c^4}$  yields

$$G_{\text{Glob}} = \frac{\left(\frac{2\alpha \cdot 1 \text{ m}}{c^2}\right)^2}{2\alpha \left(\frac{1 \text{ m}^2}{c^4}\right) c} = \frac{2\alpha}{c}. \quad (208)$$

### D. Fundamental identity and hierarchy of fields

Equations (206) and (208) give the central identity

$$G_{\text{Glob}} \equiv I_{\text{max}}, \quad (209)$$

now seen as a corollary of (i) symmetry and conservation (Noether current of the torsional  $U(1)$  sector) and (ii) the constitutive synthesis of constants in QEG. This identity also clarifies the hierarchy relations established in Sec. XXIII:

$$\vec{E} = \vec{B} \cdot I_{\text{max}}, \quad \vec{g} = \vec{\tau} \cdot I_{\text{max}}, \quad (210)$$

so that the same invariant  $I_{\text{max}} = G_{\text{Glob}}$  functions as the universal conversion factor from primary azimuthal (circulatory) modes to secondary radial (potential) modes in both electromagnetic and gravito-entropic sectors.

### E. Validation: inductive origin of the vacuum energy density

As a quantitative check, the fiducial vacuum inductance is  $L_{\text{vac}} = \mu_0 \cdot 1 \text{ m}$ . Assuming equipartition between magnetic and electric energies in a fundamental coherent mode, the peak energy is twice the magnetic peak:

$$E_{\text{peak}} = 2L_{\text{vac}} I_{\text{max}}^2 = 2(\mu_0 \cdot 1 \text{ m}) \left(\frac{2\alpha}{c}\right)^2 \quad (211)$$

Interpreting  $E_{\text{peak}}$  as the mass-energy in the reference volume  $V_{\text{ref}} = 1 \text{ m}^3$  gives the vacuum mass density

$$\rho_{\text{vac}} = \frac{E_{\text{peak}}}{V_{\text{ref}}} = \frac{2(\mu_0 \cdot 1 \text{ m}) \left(\frac{2\alpha}{c}\right)^2}{(1 \text{ m})^3}$$

$$\approx 5.956 \times 10^{-27} \text{ kg m}^{-3} \quad (212)$$

in excellent agreement with cosmological observations [43]. This agreement validates the identification  $G_{\text{Glob}} \equiv I_{\text{max}}$  and the use of  $I_{\text{max}}$  as the fundamental, symmetry-fixed scale governing both the hierarchy of fields and the global gravitational coupling.

a. *Summary.* Starting from the QEG torsional Lagrangian and its  $U(1)$  symmetry, Noether's theorem uniquely fixes a conserved current whose fundamental amplitude in the fiducial cell is  $I_{\text{max}} = 2\alpha/c$ . An independent route from the definition of  $\alpha$  yields  $G_{\text{Glob}} = 2\alpha/c$ . Their equality is therefore not a numerical accident but a constitutive identity of the vacuum: the global gravitational constant is the fundamental Noether current amplitude of the torsional sector. This anchors the macroscopic gravitational coupling in the microscopic symmetry structure of the QEG substrate and explains, at once, the weakness of gravity and the universality of the conversion factor in the field hierarchy.

## XXV. CASIMIR CONSTANT AS THE FUNDAMENTAL QUANTUM OF VACUUM PRESSURE

The Casimir effect [54, 55] is a direct manifestation of the mechanical stress of the quantum vacuum. Within the QEG framework, we derive the fundamental quantum of this vacuum pressure from first principles, combining thermodynamic and geometric reasoning.

### A. From Zero-Point Energy to Elementary Force

The irreducible energy of a vacuum oscillator of characteristic wavelength  $L$  is its zero-point energy:

$$E_0(L) = \frac{\hbar c}{2L}. \quad (213)$$

As we have already seen in Section XXIII A, this is a universal result of the quantum harmonic oscillator and represents the minimal coherent excitation of the vacuum. The elementary force conjugate to the length scale  $L$  follows from the principle of virtual work:

$$F_0(L) = \left| -\frac{\partial E_0}{\partial L} \right| = \frac{\hbar c}{2L^2}. \quad (214)$$

This step uses only the variational identity  $F = -\partial E/\partial L$  and requires no additional assumptions.

### B. From Force to Isotropic Pressure: Laplacian Flux Conservation

To connect this force to a physical pressure, we appeal to the Laplacian structure of the substrate. In the static limit, the vacuum response is conservative and flux-preserving (Gauss's law). Thus, the total force flux  $F_0$  must distribute isotropically over the surface of a sphere of radius  $L$ , with area  $A = 4\pi L^2$ . The associated vacuum pressure quantum is therefore

$$P_0(L) = \frac{F_0(L)}{4\pi L^2} = \frac{\hbar c}{8\pi L^4}. \quad (215)$$

This is the universal unit of isotropic vacuum stress: its scaling  $\hbar c/L^4$  follows from dimensional analysis, while the prefactor  $1/(8\pi)$  follows uniquely from flux conservation in three dimensions, consistent with the Green kernel  $1/(4\pi r)$ .

### C. Thermodynamic Consistency Check

The same result can be obtained directly from thermodynamic first principles. Pressure is defined as the negative gradient of energy with respect to volume:

$$P = -\frac{\partial E}{\partial V}. \quad (216)$$

Assigning to a single isotropic mode of scale  $L$  the natural spherical volume  $V = \frac{4}{3}\pi L^3$ , one has  $dV = 4\pi L^2 dL$ . Applying the chain rule,

$$P_0(L) = -\frac{\partial E_0}{\partial L} \cdot \frac{dL}{dV} = \left( \frac{\hbar c}{2L^2} \right) \left( \frac{1}{4\pi L^2} \right) = \frac{\hbar c}{8\pi L^4}, \quad (217)$$

identical to the flux-conservation result in Eq. (215). This dual derivation—from both virtual work and thermodynamic definition—confirms that  $P_0(L)$  is the fundamental, geometry-independent quantum of vacuum pressure.

### D. From the Quantum to the Observable Casimir Force

The magnitude of the Casimir force per unit area  $A$  between two perfectly conducting plates separated by a distance  $d$  is classically given by:

$$\frac{F_C}{A} = -\frac{\pi^2 \hbar c}{240d^4} \approx \frac{1.3 \times 10^{-27} \text{ N} \cdot \text{m}^2}{d^4},$$

whereas using the quantum of vacuum pressure yields

$$\frac{F_C}{A} = \frac{\hbar c}{8\pi L^4} \approx \frac{1.26 \times 10^{-27} \text{ N} \cdot \text{m}^2}{L^4}$$

The agreement with both theoretical estimates and experimental measurements [56, 57] confirms the validity of this first-order approximation derived from QEG principles. The second-order approximations can be attributed to radiative corrections, and the topology and spectral properties of the boundary configuration.

## XXVI. THE GRAVITATIONAL CONSTANT AS AN EMERGENT SUM OVER QUANTUM VACUUM MODES

In this section, we present a consistent derivation of the Newtonian gravitational constant,  $G$ , from the first principles of the framework. We will demonstrate that  $G$  emerges from the collective effect of all quantum oscillatory modes of the vacuum. The derivation begins by establishing the correct on-shell action of spacetime geometry, which is then connected to the vacuum's intrinsic elastic force. The result reveals a profound link between gravity and the mathematical structure of quantum mechanics.

### A. The On-Shell Action of a Vacuum-Dominated Spacetime from the Einstein-Hilbert action

The most fundamental description of pure geometry is the Einstein-Hilbert action,  $S_{EH} = \frac{c^4}{16\pi G} \int R \sqrt{-g} d^4x$  [58] [59] [4]. To evaluate the action for the physical vacuum, we must evaluate it for the appropriate solution to the field equations. In a universe whose energy content is dominated by the vacuum energy density  $\rho_{\text{vac}}$  (and its associated cosmological constant  $\Lambda$ ), the correct solution is the de Sitter metric, for which the Ricci scalar is constant and given by  $R = 4\Lambda$ .

Crucially, our framework posits that the vacuum energy (and thus  $\Lambda$ ) is not a fundamental parameter to be added to the action, but an *emerging property* of the substrate's quantum fluctuations. Therefore, the physically consistent approach is to start with the action for pure geometry ( $\int R$ ) and to evaluate it for the physical vacuum solution generated by these fluctuations ( $R = 4\Lambda$ ). This is conceptually distinct from the standard on-shell action which evaluates the effective Lagrangian ( $R - 2\Lambda$ ), as our model derives  $\Lambda$

from the substrate's dynamics rather than postulating it. The action integral is therefore sourced by the full geometric response of the vacuum to its own emergent energy content:

$$S_{EH} = \frac{c^4}{16\pi G} \int 4\Lambda\sqrt{-g} d^4x \quad (218)$$

We can substitute the cosmological constant  $\Lambda$  via

$$\Lambda = 8\pi G \frac{\rho_{vac}}{c^2}, \quad (219)$$

to obtain that

$$\begin{aligned} S_{EH} &= \frac{c^4}{16\pi G} \int R\sqrt{-g} d^4x \\ &= \frac{c^4}{16\pi G} \int 4\Lambda\sqrt{-g} d^4x \\ &= \frac{c^4}{16\pi G} \int \frac{32\pi G}{c^2} \cdot \rho_{vac}\sqrt{-g} d^4x \\ &= \int 2 \cdot \rho_{vac} \cdot c^2 \sqrt{-g} d^4x \end{aligned} \quad (220)$$

Substituting, we have that

$$2 \cdot \rho_{vac} \cdot c^2 = 2 \frac{\hbar c}{1 m^4} c^2 = 2 \frac{\hbar c^3}{1 m^4} \quad (221)$$

In an almost flat universe, spacetime is only slightly curved, and the metric tensor  $g_{\mu\nu}$  deviates minimally from the flat Minkowski metric  $\eta_{\mu\nu}$ . Therefore, the determinant of the metric tensor  $g$  can be expressed as:

$$\sqrt{-g} \equiv 1 + \frac{1}{2}\delta g. \quad (222)$$

For practical purposes in an almost flat universe,  $\delta g$  is so small that  $\sqrt{-g} \approx 1$  is a valid approximation. This is fully justified at the modal-cell level, where deviations from flatness are negligible compared to the scale of coarse-graining.

To evaluate this action, we integrate over the baseline 4-volume of the substrate,  $d^4x_{base} = (1 \text{ m})^4$ , which represents the fundamental cell of spacetime in our framework. This yields the total geometric action generated by the vacuum's quantum fluctuations within a reference unit:

$$S_{EH} = \left( 2 \frac{\hbar c^3}{1 m^4} \right) \cdot (1 m^4) = 2\hbar c^3 \quad (223)$$

Using the framework's fundamental identity  $\hbar \equiv 1 \text{ m}^2/c^4$  (derived in Sec. XII B), this action has a clear geometric interpretation:

$$S_{EH} = 2 \left( \frac{1 \text{ m}^2}{c^4} \right) c^3 = \frac{2 \text{ m}^2}{c} \quad (224)$$

This quantity, with dimensions of action, represents the total geometric action generated by the full energy content of the vacuum's quantum fluctuations.

## B. The Fundamental Elastic Force of the Substrate

Our framework is built upon the principle that the vacuum behaves as an elastic medium. The relationship between an applied action, the resulting deformation, and the emergent restoring force can be

described by a consistent set of Hookean relations. As established, the force-like response ( $F$ ) and the deformation-like displacement ( $x$ ) are related by the substrate's stiffness ( $k$ ):

$$F = -kx \quad (225)$$

Furthermore, we have established a dimensionally consistent constitutive law unique to this framework, which relates the deformation  $x$  to the applied action  $S$  via the same stiffness parameter:

$$x = kS \quad (226)$$

Combining these two expressions yields a direct relationship between the action applied to the substrate and the resulting microscopic restoring force it exerts:

$$F = -k^2S \quad (227)$$

To evaluate this fundamental force, we must identify its two components from the principles already derived:

- **The Geometric Stiffness ( $k$ ):** The stiffness  $k$  represents the vacuum's intrinsic resistance to deformation. For a three-dimensional isotropic medium, this response is governed by the Green's function of the Laplacian operator. Its geometric signature is the factor  $1/(4\pi r)$ . The stiffness, with dimensions of inverse length, is therefore the modulus of this geometric response evaluated at the reference scale ( $r = 1 \text{ m}$ ):

$$k \equiv \frac{1}{4\pi \cdot 1 \text{ m}} \quad (228)$$

- **The Vacuum Action ( $S$ ):** The action  $S$  corresponds to the dynamics of the vacuum in a baseline spacetime cell. As derived in Section XXVI A, the most fundamental description of pure geometry, Einstein-Hilbert action, evaluates to the quantum-relativistic constant  $S_{EH} \equiv \frac{2 \text{ m}^2}{c}$ .

## C. Derivation of the Gravitational Constant

By substituting the geometric stiffness (228) and the vacuum action (XII B) into the fundamental force law (227), we can calculate the characteristic Hookean force of the vacuum substrate:

$$\begin{aligned} F &= k^2S = \left( \frac{1}{4\pi \cdot 1 \text{ m}} \right)^2 \cdot \left( 2 \frac{(1 \text{ m})^2}{c} \right) \\ &= \left( \frac{1}{16\pi^2 \cdot (1 \text{ m})^2} \right) \cdot \left( 2 \frac{(1 \text{ m})^2}{c} \right) \\ &= \frac{1}{16\pi c} \cdot \left( \frac{2}{\pi} \right) \end{aligned} \quad (229)$$

This expression represents the microscopic, fundamental elastic force intrinsic to the vacuum. To connect this to the macroscopic world, we recall the expression for the gravitational constant  $G$  derived previously in Section XIX E:

$$G \equiv \frac{1}{16\pi c} \quad (230)$$

Comparing the fundamental force  $F$  (229) with the gravitational constant  $G$  (230), we uncover a direct and strikingly simple relationship:

$$\boxed{G = F \cdot \left( \frac{\pi}{2} \right)} \quad (231)$$

#### D. Physical Interpretation: Gravity as the Wallis Product of Vacuum Modes

The identity in Eq. (231) reveals the deepest nature of gravity within this framework. The numerical factor  $\pi/2$  is not an arbitrary constant, but a well-known mathematical result with profound connections to quantum physics: the *Wallis Product*.

$$\frac{\pi}{2} = \prod_{n=1}^{\infty} \frac{4n^2}{4n^2 - 1} = \left(\frac{2 \cdot 2}{1 \cdot 3}\right) \cdot \left(\frac{4 \cdot 4}{3 \cdot 5}\right) \cdot \left(\frac{6 \cdot 6}{5 \cdot 7}\right) \cdot \dots \quad (232)$$

*Geometric factor as a modal product: the inevitability of  $\pi/2$*

1. *The Physical Basis: A Network of Quantum Oscillators* Assume that the vacuum substrate is a network of quantum harmonic oscillators (QHO), each corresponding to a mode  $n$  with characteristic frequency  $\omega_n = n\omega_0$ . The global geometric factor that rescales the gravitational constant  $G$  must therefore emerge from the combined contribution of all these modes.

We assume that each mode  $n$  contributes to the total geometric factor through a dimensionless term  $g_n$ . If the modal contributions are independent or multiplicative (as in partition functions or Gaussian determinants), the total factor is given by the infinite product:

$$\Gamma = \prod_{n=1}^{\infty} g_n.$$

2. *The Geometric Contribution of a Single Mode* The form of  $g_n$  must reflect the physics of the oscillator. In a dynamical system, equilibrium constants typically arise from a balance between *excitation* (self-interaction of the mode) and *restoration* (anchoring of the mode to the network). For the  $2n$ -th harmonic, the intensity of self-interaction scales as  $(2n)^2$  (two creation/annihilation operators in a quadratic transition), while the most immediate restoration comes from its neighbors  $2n \pm 1$ . Thus,

$$g_n \equiv \frac{\text{Excitation}}{\text{Restoration}} = \frac{(2n)^2}{(2n-1)(2n+1)} = \frac{4n^2}{4n^2 - 1}.$$

This is not ad hoc: it is the natural ratio between the “weight” of the mode itself and the minimum “anchoring” imposed by its nearest neighbors in the modal ladder.

3. *The Collective Result: the Wallis Product* The collective geometric factor is then

$$\Gamma = \prod_{n=1}^{\infty} \frac{4n^2}{4n^2 - 1}.$$

This infinite product is exactly the *Wallis product*, and it converges to

$$\Gamma = \frac{\pi}{2}.$$

Therefore, the geometric scaling that links the fundamental elastic force of the substrate to the gravitational constant  $G$  is uniquely fixed, without additional freedom, by the completeness of the modal summation.

*Formal justification (mathematical and quantum, in two steps)*

(i) *Mathematical—Euler’s sine product*. The classical Euler identity

$$\sin x = x \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{\pi^2 n^2}\right)$$

implies, upon evaluation at  $x = \pi/2$ , that

$$\frac{2}{\pi} = \prod_{n=1}^{\infty} \left(1 - \frac{1}{4n^2}\right) \implies \prod_{n=1}^{\infty} \frac{4n^2}{4n^2 - 1} = \frac{\pi}{2}.$$

Thus, closing the product over the entire modal ladder  $n = 1, 2, \dots$  inevitably enforces the  $\pi/2$  factor.

(ii) *Quantum—Modal determinant in the QHO*. In the path integral formalism, the propagator of the one-dimensional harmonic oscillator reads

$$K(x_f, T; x_i, 0) = \sqrt{\frac{m\omega}{2\pi i \hbar \sin(\omega T)}} \exp\left[\frac{i}{\hbar} S_{\text{cl}}(x_i, x_f; T)\right],$$

where the prefactor  $[\sin(\omega T)]^{-1/2}$  arises from the *infinite product* over Fourier modes (Gaussian determinant) and is evaluated using Euler’s sine product. For a natural “quadrature shift” of the modal cell,  $\omega T = \pi/2$ , one obtains

$$\prod_{n=1}^{\infty} \left(1 - \frac{1}{4n^2}\right) = \frac{2}{\pi} \iff \prod_{n=1}^{\infty} \frac{4n^2}{4n^2 - 1} = \frac{\pi}{2}.$$

That is, upon performing the complete modal decomposition in the standard formalism, the Wallis factor appears inevitably.

*Conclusion* The factor  $\pi/2$  is not a numerical coincidence nor an empirical fit: it is the *statistical signature* of coherently summing the entire ladder of oscillatory modes of the quantum-elastic substrate. Consequently, *the constant of gravitation  $G$  is the macroscopic measure of the vacuum’s fundamental elastic force, scaled by the normalized, collective contribution of the infinite ladder of quantum oscillatory modes of the substrate.*

This final result has two transformative implications:

1. **Gravity as Quantum Vacuum Pressure:** The connection solidifies the interpretation of gravity as a phenomenon emergent from the quantum vacuum, analogous to the Casimir effect. The value of  $G$  is a direct measure of the “pressure” exerted by the vacuum’s zero-point fluctuations. Its weakness is a consequence of the specific geometric and elastic properties of the substrate.
2. **Gravity as an Entropic/Informational Constant:** Because the Wallis product is a sum over all possible modes ( $n = 1, \dots, \infty$ ), the value of  $G$  is determined by the total number of degrees of freedom of the spacetime substrate. This aligns perfectly with the paradigm of entropic gravity, where gravity is not a force but a statistical manifestation of information. The constant  $G$  ceases to be a simple coupling constant and becomes a measure of the information capacity of the vacuum itself.

In conclusion, this framework establishes that the gravitational constant is the macroscopic manifestation of the vacuum's collective oscillatory structure. The Wallis factor ensures the unique consistency of the modal summation, thereby unifying the geometric, elastic, and quantum aspects of the substrate in a single, inevitable identity.

## XXVII. THE COSMOLOGICAL CONSTANT AS AN EMERGENT PROPERTY OF THE QUANTUM VACUUM

In this section, we derive the origin and magnitude of the cosmological constant,  $\Lambda$ , from the first principles of Quantum-Elastic Geometry. We demonstrate that  $\Lambda$  is not a fundamental parameter to be added to the action, but is instead an inevitable consequence of the granular, oscillatory structure of the spacetime substrate. By modeling the vacuum as a network of quantum harmonic oscillators, we show that the collective zero-point energy of these modes, regularized by the substrate's own fundamental scale, gives rise to a vacuum energy density that, in the macroscopic limit, is identified with the observed cosmological constant. This result elevates  $\Lambda$  from a persistent puzzle to a core prediction of the theory.

### A. The Substrate as a Quantum Oscillator Network and its Zero-Point Energy

The foundational postulate of QEG is that spacetime is a physical, elastic medium quantized as a lattice of oscillators. Each normal mode of this lattice, indexed by its wavevector  $\vec{k}$ , behaves as an independent quantum harmonic oscillator (QHO). As established by the universal principles of quantum mechanics, any such oscillator possesses an irreducible ground-state energy, or zero-point energy, given by:

$$E_{0,k} = \frac{1}{2} \hbar \omega_k \quad (233)$$

As already derived in Section XV, the measured energy density ( $\rho_{\text{vac}}$ ) is a macroscopic, cosmological observable. It reflects the net effect of an immense number of uncoordinated vacuum oscillators, with their phases and spatial orientations being statistically random. At the macroscopic level, statistical isotropy enforces a coarse-grained average over random phases, entirely analogous to the emergence of  $g_{\mu\nu} = \langle G_{\mu\nu} \rangle$ . The observable energy density is therefore

$$\rho_{\text{eff}} \equiv \frac{1}{2\pi} \int_0^{2\pi} \mathcal{L}_{\text{modal}} d\theta = \frac{\mathcal{L}_{\text{modal}}}{2\pi} = \frac{\hbar c}{2\pi \cdot 1 \text{ m}^4} XV$$

### B. From Microscopic Energy to Macroscopic Curvature: The Emergence of $\Lambda$

The coarse-graining of the high-frequency quantum fluctuations of the substrate field  $\mathcal{G}_{\mu\nu}$  leads to an effective low-energy theory for the macroscopic metric  $g_{\mu\nu}$ , which takes the form of the Einstein-Hilbert action. In this emergent framework, the microscopic vacuum energy density  $\rho_{\text{vac}}$  acts as a persistent, isotropic source term for the macroscopic geometry. The standard relationship from General Relativity,

$$\Lambda = \frac{8\pi G}{c^2} \rho_{\text{vac}} \quad (234)$$

is reinterpreted in QEG not as a definition, but as a *consistency condition* that connects the microscopic physics of the vacuum to the large-scale curvature it generates. By substituting our derived expression for  $\rho_{\text{vac}}$  (Eq. XV) into this condition, we obtain a first-principles prediction for the cosmological constant:

$$\Lambda = \frac{8\pi G}{c^2} \left( \frac{\hbar c}{2\pi \cdot 1 \text{ m}^4} \right) = \frac{4 \cdot G \hbar}{c \cdot 1 \text{ m}^4} \quad (235)$$

### C. Synthesis with the Planck Scale and Physical Interpretation

This result reveals a spectacular connection to the fundamental scale of quantum gravity. By using the definition of the Planck length,  $l_p^2 = \frac{G \hbar}{c^3}$ , we can rewrite Eq. 235 as:

$$\Lambda = \frac{4G\hbar}{c \cdot 1 \text{ m}^4} \implies \boxed{\Lambda = 4 \frac{(l_p \cdot c)^2}{1 \text{ m}^4}} \quad (236)$$

*Mechanical Interpretation:  $\Lambda$  as Vacuum Kinetic Pressure and Torsional Inertia*

The expression for the cosmological constant in Eq. (236) allows for a profound physical interpretation in mechanical terms, further grounding the properties of spacetime in the tangible dynamics of the QEG substrate. This interpretation can be viewed from two complementary perspectives, linear and rotational.

*a. Kinetic Momentum Interpretation.* Within the QEG framework, where velocity is dimensionless, the quantity  $p_{pl} = l_p \cdot c$  can be identified as the fundamental *Planck momentum*. It represents the momentum of the most primordial quantum fluctuation: a Planck-scale excitation propagating at the maximum causal speed. With this identification, Eq. (236) becomes:

$$\Lambda = 4 \frac{p_{pl}^2}{(1 \text{ m})^4} \quad (237)$$

Given that the square of momentum ( $p^2$ ) is directly related to kinetic energy, this expression reveals that the cosmological constant can be understood as the *density of kinetic pressure* exerted by the zero-point fluctuations of the vacuum. The accelerated expansion of the universe is thus reinterpreted not as the effect of a mysterious "dark energy," but as the macroscopic manifestation of the incessant kinetic impulse of the spacetime substrate's own fundamental quantum vibrations.

*b. Torsional Inertia Interpretation.* Alternatively, the term  $(l_p \cdot c)^2$  has dimensions of  $[L^2]$ , which is dimensionally equivalent to the *moment of inertia per unit mass* ( $I/m \sim r^2$ ). We can therefore identify this term with the *fundamental rotational inertia* of a primordial cell of the vacuum substrate. The formula for  $\Lambda$  then acquires a new meaning:

$$\Lambda = 4 \frac{I_{\text{fund}}/m_{\text{fund}}}{(1 \text{ m})^4} \quad (238)$$

From this perspective, the cosmological constant measures the *density of the vacuum's torsional inertia*. It reflects spacetime's intrinsic resistance to rotational or shear-like deformations. This view powerfully connects the cosmic expansion to the torsional and

shear modes of the spatial substrate itself (encoded within  $\mathcal{G}_{ij}$ ), suggesting that the universe's expansion could be the result of the substrate "unwinding" a primordial torsional stress, governed by its intrinsic inertia.

These two interpretations—linear momentum and rotational inertia—are complementary facets of the same underlying mechanical reality. They solidify the view of spacetime as a dynamic, physical medium, whose fundamental properties dictate the large-scale evolution of the cosmos.

#### D. Consistency with Geometric and Thermo-Entropic Structures

The QEG prediction for  $\Lambda$ , derived from quantum principles, must be consistent with the well-established geometric and thermodynamic roles it plays in physics. This subsection demonstrates this perfect correspondence.

*From Planck Momentum to Spacetime Curvature:  
The Microscopic Origin of Vacuum Energy in  
General Relativity*

The standard framework of General Relativity (GR) accommodates the observed cosmic acceleration by introducing the cosmological constant,  $\Lambda$ , which is interpreted as the energy density of the vacuum itself. In this subsection, we demonstrate that the phenomenological description of vacuum energy in GR is a direct macroscopic consequence of the fundamental principles of QEG, originating from the primordial momentum of the substrate's quantum fluctuations.

In GR, the energy content of the vacuum is described by an effective energy-momentum tensor of a perfect fluid:

$$T_{\mu\nu}^{\text{vac}} = (\rho_{\text{vac}} + p_{\text{vac}})u_{\mu}u_{\nu} + p_{\text{vac}}g_{\mu\nu} \quad (239)$$

For this tensor to be Lorentz invariant, as the vacuum must be, it requires that the pressure be exactly  $p_{\text{vac}} = -\rho_{\text{vac}}$ . This leads to the well-known form  $T_{\mu\nu}^{\text{vac}} = \rho_{\text{vac}}g_{\mu\nu}$ , and the consistency condition  $\rho_{\text{vac}} = \Lambda c^2 / (8\pi G)$ . While GR requires this negative pressure, it does not explain its physical origin.

QEG provides this missing physical foundation. We have derived the cosmological constant from the first principles of the quantum-elastic substrate (Eq. 236):

$$\Lambda = \frac{4(l_p \cdot c)^2}{(1 \text{ m})^4} = \frac{4p_{pl}^2}{(1 \text{ m})^4} \quad (240)$$

where we have identified  $p_{pl} \equiv l_p \cdot c$  as the fundamental *Planck momentum*. By substituting this result into the GR consistency condition, we obtain a first-principles expression for the vacuum energy density:

$$\rho_{\text{vac}} = \frac{\Lambda c^2}{8\pi G} = \frac{c^2}{8\pi G} \left( \frac{4p_{pl}^2}{(1 \text{ m})^4} \right) = \frac{p_{pl}^2 c^2}{2\pi G \cdot (1 \text{ m})^4} \quad (241)$$

This equation is a powerful bridge between the micro and macro worlds. It dictates that the vacuum energy density is determined by the kinetic pressure of the substrate's most fundamental quantum fluctuations.

This perspective provides a direct physical origin for the mysterious *negative pressure*. The kinetic pressure exerted by the zero-point fluctuations is inherently isotropic—it pushes outwards in all directions. In an expanding spacetime, a pressure that pushes outwards does negative work on its surroundings ( $dW = -pdV$ ), which is the defining characteristic of negative pressure. Therefore, the enigmatic negative pressure of dark energy is reinterpreted in QEG as the natural and inevitable consequence of the kinetic pressure of the vacuum's quantum vibrations.

This result provides the concrete microscopic mechanism for the emergent gravity paradigm, as hinted at by Jacobson and Verlinde [10, 11]. Their frameworks posit that gravity emerges from underlying thermodynamic and entropic principles. QEG provides the physical substrate for these ideas: the vacuum is a network of quantum oscillators whose collective kinetic and inertial properties manifest simultaneously as the *thermodynamics of the vacuum* (entropy and temperature) and as its *gravitational effect* (spacetime curvature via  $\Lambda$ ).

*Thermo-Entropic Action Density and the Geometric Structure of the Cosmological Constant*

Substituting  $\hbar \equiv \frac{1}{c^4} m^2$  from (XII B) and  $G \equiv \frac{1}{16\pi c}$  from (134) in Eq. 235, we arrive at:

$$\Lambda \equiv \frac{1}{4\pi c^6 \cdot 1 \text{ m}^2} \quad (242)$$

Recalling that the thermo-entropic modal action was shown to be  $S_{\text{th}} = \frac{\hbar}{c^2}$ , we see that

$$\Lambda = \frac{S_{\text{th}}}{4\pi \cdot 1 \text{ m}^4}$$

This reveals the cosmological constant as the manifestation of a universal *modal action density per 4D volume*. It acts as a tension-like Lagrangian density of the vacuum, coupling entropy and expansion across a fundamental volume cell.

This relationship can be further illuminated by rewriting it to explicitly reveal its constitutive components:

$$\Lambda = \left( \frac{1}{4\pi \cdot 1 \text{ m}} \right) \cdot \left( \frac{S_{\text{th}}}{1 \text{ m}^3} \right) \quad (243)$$

This formulation is particularly powerful as it provides a direct mechanical origin for the cosmological constant. The first term is precisely the *Geometric Stiffness Modulus* of the substrate, derived from the 3D Laplacian Green's function, which quantifies the vacuum's intrinsic elastic resistance to deformation. The second term represents the *Thermo-Entropic Action Density* per unit volume, the measure of the substrate's minimal, dissipative quantum action. Thus, the cosmological constant is revealed to be the product of the vacuum's fundamental stiffness and its baseline entropic action density. This recasts  $\Lambda$  not as an abstract energy value, but as the *intrinsic elastic tension* of the spacetime substrate, generated in response to its own fundamental thermodynamic state. This perfectly unifies the geometric, thermodynamic, and quantum aspects of the vacuum within a single, coherent mechanical framework.

### Interpretation as Geometric Surface Tension

Indeed, note that setting  $r = c^3 \cdot 1 \text{ m}$  situates  $\Lambda$  as an effective curvature density, with  $4\pi r^2 = 4\pi c^6 \cdot 1 \text{ m}^2$  representing the "surface" of an expanding spherical volume. Thus, the cosmological constant acquires a direct geometrical interpretation as an *inverse areal curvature density*, analogous to curvature or density of a spherical boundary in expanding space, projecting a constant action flux over the expanding boundary of the universe. From this viewpoint,  $\Lambda$  is not a bulk energy density but a quantized surface effect—a geometric relic of the thermo-entropic elasticity of the vacuum.

This form provides a physical interpretation in which the large-scale expansion of the universe is driven by a steady energy flow that distributes itself over the expanding boundary, dynamically adjusting the effective curvature density as the volume of the universe grows. This interpretation not only aligns with the curvature requirements of an accelerating universe but also positions  $\Lambda$  as a fundamental invariant describing how the vacuum tension distributes minimal action quanta across areal elements, reinforcing the idea that the cosmological constant is, in essence, the *vacuum's curvature Lagrangian*.

### The Inevitable Gauge-like Structure of the Thermo-Entropic Field

The principles of QEG demand that every mode of the unified substrate  $\mathcal{G}_{\mu\nu}$  be described by a self-consistent, covariant field theory. Having identified the scalar ( $\mathcal{G}_{00}$ ) and vector ( $\mathcal{G}_{0i}$ ) modes with gravity and electromagnetism, the remaining degrees of freedom within the spatial tensor  $\mathcal{G}_{ij}$ —which we have identified as the thermo-entropic field—must also adhere to these principles.

For a massless or very light field emerging in the low-energy limit, the principles of Lorentz covariance and locality uniquely constrain the form of its kinetic action. The most general, non-trivial Lagrangian at the two-derivative level must be quadratic in a field-strength tensor derived from an underlying potential. *This is not an analogy, but a structural necessity for any fundamental interaction.* Therefore, the dynamics of the thermo-entropic sector *must* be describable by an effective field strength tensor,  $\mathcal{F}_{\mu\nu}^{(GE)}$ , constructed from the underlying modes of the substrate:

$$\mathcal{F}_{\mu\nu}^{(GE)} := \partial_\mu \mathcal{G}_\nu^{(T)} - \partial_\nu \mathcal{G}_\mu^{(T)} \quad (244)$$

where  $\mathcal{G}_\mu^{(T)}$  represents the effective potential of the thermo-entropic modes. The corresponding Lagrangian density for this sector is thus structurally fixed to the canonical form:

$$\mathcal{L}_{GE} = -\frac{1}{4k_{GE}} \mathcal{F}_{\mu\nu}^{(GE)} \mathcal{F}^{(GE)\mu\nu} \quad (245)$$

where  $k_{GE}$  is the modal stiffness constant for the thermo-entropic sector.

The total energy density of the vacuum,  $\rho_{vac}$ , is the sum of the zero-point energies of all substrate modes. The cosmological constant, as the macroscopic manifestation of this energy, is therefore the vacuum expectation value of the sum of the Lagrangians of all field modes,  $\Lambda = \langle \sum_i \mathcal{L}_i \rangle$ . In a vacuum dominated by the thermo-entropic fluctuations, this simplifies to  $\Lambda \approx \langle \mathcal{L}_{GE} \rangle$ .

This provides a profound meaning to our derived result reflected in Eq. 235 and Eq. 236:

$$\Lambda = 4 \cdot \frac{\hbar G}{c \cdot 1 \text{ m}^4} = 4 \frac{(l_p \cdot c)^2}{1 \text{ m}^4} \quad (246)$$

The equations reflect precisely the canonical form of Eq. XXVII D, and validates them as a fundamental result derived from the vacuum's zero-point energy, which—when formalized into a consistent field theory—necessarily adopts the canonical normalization of a gauge-like structure. The correspondence is a powerful confirmation of the theory's internal consistency.

### Gravity as an Emergent Thermodynamic Action

The reduction of the Einstein-Hilbert action in a vacuum-dominated universe to the form (220)

$$S_{EH} = \frac{c^4}{16\pi G} \int 4\Lambda \sqrt{-g} d^4x = \frac{c^4}{4\pi G} \int \Lambda \sqrt{-g} d^4x \quad (247)$$

reveals a profound identity when combined with our identification of the cosmological constant  $\Lambda$  as the effective Lagrangian density of the thermo-entropic field,  $\Lambda = \langle \mathcal{L}_{GE} \rangle$ . Substituting this identification back into the Einstein-Hilbert action, we obtain:

$$S_{EH} = \frac{c^4}{4\pi G} \int \langle \mathcal{L}_{GE} \rangle \sqrt{-g} d^4x \quad (248)$$

This result has fundamental implications for the nature of gravity. It demonstrates that the *Einstein-Hilbert action is, up to a prefactor, identical to the effective action of the thermo-entropic field.* The dynamics of spacetime geometry are thus shown to be a direct macroscopic manifestation of the underlying thermodynamics of the vacuum. This provides a concrete physical basis for the emergent gravity paradigm, realizing the vision hinted at by Jacobson and Verlinde [10] [11]. The principle of least action for geometry ( $\delta S_{EH} = 0$ ) is reinterpreted as a principle of *extremal entropic action*, and the prefactor  $\frac{c^4}{4\pi G}$  becomes the fundamental conversion factor between the two descriptions.

### E. Derivation of the Planck Length from the First Principles of QEG

The network of constitutive identities developed within the QEG framework not only predicts cosmological constants but also allows for the derivation of nature's most fundamental scales from its primordial principles. In this subsection, we demonstrate how the Planck length,  $l_p$ —the minimal granularity of spacetime—emerges as a direct consequence of the substrate's quantum-elastic properties.

We begin with the standard definition of the Planck length:

$$l_p^2 = \frac{G\hbar}{c^3} \quad (249)$$

Within QEG,  $G$  and  $\hbar$  are not independent, fundamental constants but are instead emergent properties of the substrate. Using the identities derived in previous sections:

- The gravitational constant, as a manifestation of the vacuum's elasticity (Sec. XIX E):  $G \equiv \frac{1}{16\pi c}$ .
- The quantum of action, as the minimal deformation area of the electromagnetic mode (Sec. 149):  $\hbar \equiv \frac{1}{c^4} m^2$ .

We substitute these QEG identities into the definition of the Planck length:

$$l_p^2 = \frac{\left(\frac{1}{16\pi c}\right) \left(\frac{1}{c^4} m^2\right)}{c^3} = \frac{1}{16\pi c^8} m^2 \quad (250)$$

Taking the square root, we obtain the Planck length expressed purely in terms of the speed of light and the fiducial meter X D scale:

$$l_p = \frac{1}{4\sqrt{\pi}} \frac{m}{c^4} \quad (251)$$

We can rewrite this expression to reveal its profound structure by using the identity  $\hbar \equiv (1 m)^2/c^4$ :

$$l_p = \frac{\hbar}{4\sqrt{\pi} \cdot 1 m} \quad \Rightarrow \quad \boxed{l_p = \left(\frac{1}{4\pi \cdot 1 m}\right) \cdot \hbar \cdot \sqrt{\pi}} \quad (252)$$

*Physical Interpretation: The Granularity of Spacetime as a Quantum Deformation*

Equation (252) is one of the most powerful validations of the QEG framework. It reveals that the Planck length is not an axiom but a derived theorem. Its structure can be decomposed into three elements, each with a clear physical meaning:

1. **The Geometric Stiffness**  $\left(\frac{1}{4\pi \cdot 1 m}\right)$ : This term represents the intrinsic elastic resistance of the substrate to being deformed. The  $1/(4\pi)$  factor is the unmistakable signature of the Laplacian operator in three dimensions, which governs all static responses of the medium.
2. **The Quantum of Action**  $(\hbar)$ : This is the primordial source of the scale, representing the minimal deformation or geometric "excitation" that the substrate can support.
3. **The Quantum Normalization Factor**  $(\sqrt{\pi})$ : This is not an arbitrary factor but the mathematical fingerprint of the substrate's nature as a network of quantum harmonic oscillators (QHOs). Its origin lies in the *Gaussian integral*  $(\int e^{-x^2} dx = \sqrt{\pi})$ , which is the foundation for normalizing quantum vacuum states and for calculating the path integrals that describe their dynamics. It also emerges from the fundamental value of the *Gamma function* at  $\Gamma(1/2) = \sqrt{\pi}$ , which quantifies the geometric factor of a primordial quantum excitation.

Therefore, Equation (252) is interpreted as follows: *the smallest possible length in nature ( $l_p$ ) is the deformation generated by the primordial quantum of action ( $\hbar$ ), scaled by the elastic stiffness of spacetime and normalized by the intrinsic geometric factor of its own quantum fluctuations.*

*Consistency with the Foundational Hookean Structure.* This result is in perfect agreement with the foundational Hookean structure of the theory, which posits that any fundamental deformation ( $x$ ) arises from the product of the substrate's stiffness ( $k$ ) and the driving source action ( $S_{\text{source}}$ ). In this context, the Planck length ( $l_p$ ) represents the minimal possible deformation of the substrate. Our derived expression (Eq. 252) precisely matches this constitutive law:

$$l_p = \underbrace{\left(\frac{1}{4\pi \cdot 1 m}\right)}_{\text{Stiffness } (k)} \cdot \underbrace{\hbar}_{\text{Effective Source } S} \cdot \underbrace{\sqrt{\pi}}_{\text{Geometric factor}} \quad (253)$$

Here, the stiffness of the substrate is its geometric modulus ( $k$ ), while the driving term is the fundamental quantum of action ( $\hbar$ ), modulated by a dimensionless geometric factor ( $\sqrt{\pi}$ ) arising from the vacuum's quantum nature. Thus, the Planck length emerges not only from kinematic identities but also as the direct result of the substrate's fundamental elastic response, solidifying the deep mechanical consistency of the QEG framework.

## XXVIII. THE SCALE-DEPENDENT GRAVITATIONAL COUPLING AND THE RESOLUTION OF COSMOLOGICAL TENSIONS

The QEG framework predicts that gravity is not characterized by a single universal constant, but exhibits distinct responses depending on the geometric distribution of the source energy and the self-interactions involved. This behavior is an inevitable consequence of the duality of self-energy (Sec. IX). As we show in this section, this duality provides a first-principles, parameter-free explanation for the Hubble tension, and naturally extends to the dark sector, structure growth, and gravitational lensing anomalies.

### A. The Two Asymptotic States of Gravity

As a direct consequence of the duality of self-energy (Sec. IX), the QEG framework inevitably predicts that the effective gravitational coupling is not a universal constant, but depends on the geometric distribution of the source energy. We can derive the two expected asymptotic values for this coupling by examining the universal geometric factors that arise from integrating over two distinct idealized source geometries:

1. **A Volume-Dominant, Self-Interacting Geometry:** This regime corresponds to localized, clumpy structures where energy density fills a volume. The standard calculation for the total self-energy of such a spherical distribution universally yields a geometric prefactor of  $3/5$ .
2. **A Surface-Dominant, Quasi-Linear Geometry:** This regime corresponds to the large-scale, homogeneous universe where strong nonlinearities are averaged out. The energy is effectively treated as residing on a boundary, analogous to the charge on a spherical conductor. This calculation universally yields a geometric prefactor of  $1/2$ .

These purely geometric factors allow us to define two distinct gravitational couplings as a direct application of Sec. IX A:

- **Local Coupling ( $G_{\text{loc}}$ ):** The volume-dominant case is identified with the local, strongly self-interacting compressive channel ( $\mathcal{G}_{00}$ ), where the total work of formation ( $U_{\text{glob}}$ ) is the relevant energy. It is identified with the standard Newtonian constant  $G_N$ :

$$G_{\text{loc}} = \frac{3}{5} 4\pi\epsilon_0 \quad (254)$$

- **Global Coupling ( $G_{\text{glob}}$ ):** The surface-dominant case is identified with the global, quasi-linear volumetric channel ( $\text{Tr}(\mathcal{G}_{ij})$ ). This fits with the homogeneous regime of the cosmos, where non-linearities are averaged out and only the external field energy ( $U_{\text{loc}}$ ) is relevant. Its value is:

$$G_{\text{glob}} = \frac{1}{2} 4\pi\epsilon_0 = 2\pi\epsilon_0 \quad (255)$$

Crucially, these two values are not free parameters adjusted to fit data. They are **parameter-free predictions** derived uniquely from the universal geometry of the substrate's elastic response. The fact that these values, when inserted into the Friedmann equations, precisely resolve the Hubble tension (as shown in the next subsection) provides powerful, non-trivial evidence for the theory's central claim: that *gravity is fundamentally geometric in nature*.

## B. A Parameter-Free Resolution of the Hubble Tension

The first Friedmann equation [60] [61] [62] is given by:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3} \rho_{\text{vac}} - \frac{kc^2}{a^2} + \frac{\Lambda c^2}{3}, \quad (256)$$

This equation relates the rate of expansion (the Hubble parameter,  $H = \dot{a}/a$ ) to the energy density of the universe. Assuming a nearly flat universe ( $k \approx 0$ ), the Hubble parameter can be calculated as

$$H^2 = \frac{8\pi G}{3} \rho_{\text{vac}} + \frac{\Lambda c^2}{3},$$

Substituting with the classical expression for  $\Lambda$  (234), we have that

$$\begin{aligned} H^2 &= \frac{8\pi G}{3} \rho_{\text{vac}} + \frac{\Lambda c^2}{3} = \\ &= \frac{8\pi G}{3} \rho_{\text{vac}} + \frac{8\pi G \rho_{\text{vac}} c^2}{3} = \\ &= 2 \left( \frac{8\pi G}{3} \rho_{\text{vac}} \right) = \frac{16\pi G}{3} \rho_{\text{vac}} \end{aligned} \quad (257)$$

Substituting  $G_{\text{Loc}}$  and  $G_{\text{Glob}}$  into the Friedmann equation yields two distinct predictions for the Hubble constant:

- **Local Hubble constant ( $H_{0,\text{Loc}}$ ):**

$$H_{0,\text{Loc}} = \sqrt{\frac{16\pi G_{\text{Loc}}}{3} \rho_{\text{vac}}} \approx 73.17 \text{ km/s/Mpc}, \quad (258)$$

in excellent agreement with the SH0ES value of  $73.0 \pm 1.0 \text{ km/s/Mpc}$  [63].

- **Global Hubble constant ( $H_{0,\text{Glob}}$ ):**

$$H_{0,\text{Glob}} = \sqrt{\frac{16\pi G_{\text{Glob}}}{3} \rho_{\text{vac}}} \approx 66.81 \text{ km/s/Mpc}, \quad (259)$$

in excellent agreement with the Planck 2018 result of  $67.4 \pm 0.5 \text{ km/s/Mpc}$  [64].

The Hubble tension is thus reinterpreted not as a conflict, but as observational evidence of a dual gravitational response. The early Universe probes the global, weakly-coupled channel, while late-time observations probe the local, strongly self-interacting one.

## C. The Dynamical Transition Mechanism: From Bare to Dressed Gravity

The existence of two asymptotic values for the gravitational coupling suggests a dynamical interpolation between the two regimes. Within QEG, this transition is not a postulate but a direct consequence of the "dressing" of the bare gravitational interaction by the dynamics of the *thermo-entropic field*.

As established in Sec. IIE, the bare, static response of the geometro-elastic substrate is governed by the Laplacian operator, corresponding to a propagator of the form  $1/k^2$  in momentum space. This describes the fundamental, unscreened interaction. However, the substrate is not merely elastic but also dissipative, with its dynamics governed by the thermo-entropic field (Sec. XX).

In the quasi-static cosmological limit, this field's dynamics become diffusive, introducing a characteristic physical scale,  $\lambda$ , related to the substrate's stiffness  $\kappa$  and viscosity  $\zeta$ . This dynamic screening mechanism modifies, or "dresses", the bare gravitational propagator. The resulting effective gravitational coupling,  $G_{\text{eff}}(k)$ , must therefore interpolate between the two asymptotic states. The general mathematical form for such a screened interaction, which transitions from a global, unscreened value at large scales ( $k \rightarrow 0$ ) to a local, bare value at short scales ( $k \rightarrow \infty$ ), is given by:

$$G_{\text{eff}}(k) = G_{\text{Glob}} + (G_{\text{Loc}} - G_{\text{Glob}}) \frac{(k\lambda)^2}{1 + (k\lambda)^2} \quad (260)$$

which can be rearranged to

$$G_{\text{eff}}(k) = G_{\text{Loc}} \left[ 1 + \frac{G_{\text{Glob}} - G_{\text{Loc}}}{G_{\text{Loc}}} \frac{1}{1 + (k\lambda)^2} \right], \quad (261)$$

where  $\lambda$  is the transition scale set by the substrate's stiffness  $\kappa$  and viscosity  $\zeta$ . This form ensures

$$\lim_{k \rightarrow \infty} G_{\text{eff}}(k) = G_{\text{Loc}}, \quad \lim_{k \rightarrow 0} G_{\text{eff}}(k) = G_{\text{Glob}}.$$

Eq. 261 is not a phenomenological ansatz, but the expected functional form for a dressed coupling constant in a theory with a diffusive screening scale. The running of  $G$  with scale is thus a dynamically inevitable consequence of the interplay between the substrate's fundamental elastic (Laplacian) response and its dissipative (thermo-entropic) dynamics.

*a. Phenomenological Model for Observational Tests.* To connect this theoretical prediction with cosmological observables, we need a phenomenological model for the evolution of  $G_{\text{eff}}$  as a function of

redshift,  $z$ . A logistic function provides a simple and flexible parameterization for such a smooth transition:

$$G_{\text{eff}}(z) = G_{\text{Glob}} + \left( \frac{G_{\text{Loc}} - G_{\text{Glob}}}{1 + e^{a(z-z_c)}} \right) \quad (262)$$

While the parameters  $z_c$  (transition redshift) and  $a$  (transition rate) are treated as free parameters when fitting to cosmological data, our theory predicts that they are not fundamental. They are effective parameters that, in principle, can be derived from the substrate's properties ( $\kappa, \zeta$ ). Specifically, the transition redshift  $z_c$  would correspond to the epoch where the mean density of the universe dropped below a critical threshold related to the substrate's intrinsic stiffness, while the rate  $a$  would be governed by its viscosity  $\zeta$ .

This model provides a concrete framework for testing the theory against a suite of cosmological data (SN Ia, BAO, RSD). A successful, consistent fit for  $z_c$  and  $a$  across multiple probes would provide compelling evidence for the dynamical transition predicted by QEG, offering a clear path to either validate or falsify the framework.

#### D. Observational Consequences

This framework makes several testable predictions:

- **Redshift evolution:**  $H(z)$  must interpolate smoothly between  $H_{\text{Glob}} \approx 67$  km/s/Mpc at high  $z$  and  $H_{\text{Loc}} \approx 73$  km/s/Mpc at low  $z$ .
- **Structure formation:** Growth rates differ depending on whether  $G_{\text{Glob}}$  or  $G_{\text{Loc}}$  dominates, offering an explanation for the  $S_8$  tension.
- **Gravitational lensing:** Weak lensing (Mpc scales) should systematically underestimate masses if analyzed with  $G_{\text{Loc}}$ , by a factor  $\sim G_{\text{Glob}}/G_{\text{Loc}} \approx 0.83$ .
- **Gravitational waves:** Propagation is governed by the tensorial channel ( $\Sigma_{ij}$ ), which remains consistent with LIGO/Virgo observations since the radiative sector is not altered by the scalar/tensorial decomposition of static gravity.

#### E. Consistency with Gravitational Wave Observations

A potential concern with scale-dependent gravity is consistency with gravitational-wave constraints. In the present framework, this consistency emerges as a natural consequence of the modal decomposition of the substrate. Gravitational waves are sourced by the traceless shear modes of  $\mathcal{G}_{ij}$  and propagate as high-frequency excitations on a background set by  $G_{\text{eff}}$ . Since the scale-dependent interpolation is governed by the *isotropic* thermo-entropic (trace) channel, it renormalizes the background coupling  $G$  without altering the hyperbolic, radiative tensor sector. Therefore, the speed and dispersion of gravitational waves remain unchanged to leading order, in agreement with LIGO/Virgo observations, while cosmological probes remain sensitive to the running of  $G_{\text{eff}}$ .

#### F. Theoretical Context and Consistency

This proposal does not modify Einstein's equations, but the effective coupling between geometry and mat-

ter. In the local, non-linear regime, General Relativity with  $G_{\text{Loc}}$  is recovered. In the global, quasi-linear regime, averaging over large volumes renormalizes the coupling to  $G_{\text{Glob}}$ . Friedmann equations thus remain valid, but with a scale-dependent  $G_{\text{eff}}(z)$ .

#### G. Implications for the Dark Sector

The scale-dependent nature of gravity reframes the interpretation of the dark sector. The fundamental quantity driving cosmic acceleration is the geometric invariant  $\Lambda$ . The effective vacuum energy density that would be inferred by an observer assuming a constant  $G$  is given by:

$$\rho_{\text{inferred}} = \frac{\Lambda c^2}{8\pi G}$$

while the real effective value would be

$$\rho_{\text{vac}} = \frac{\Lambda c^2}{8\pi G_{\text{eff}}(z)}$$

This has two profound consequences:

- **Dark energy:** The accelerated expansion is driven by the geometric invariant  $\Lambda$ , not by an ad hoc fluid. The "dark energy" of  $\Lambda$ CDM is an artifact of assuming a constant  $G$ .
- **Dark matter:** The discrepancy arises from interpreting *local dynamics* within the *global cosmological context* provided by the  $\Lambda$ CDM model. The  $\Lambda$ CDM framework is calibrated on the largest scales (e.g., the CMB), implicitly embedding the weaker global coupling,  $G_{\text{glob}}$ , into its predictions for cosmic structure and mass content. However, the actual gravitational dynamics within a local system (e.g., a galaxy) are governed by the stronger local coupling,  $G_{\text{loc}}$ . An observer comparing the observed strong local gravity to the mass content predicted by the  $G_{\text{glob}}$ -based model will find a significant gravitational deficit, which is then attributed to "missing mass" (dark matter). Therefore, a substantial fraction of the inferred dark matter may be an artifact of applying a global gravitational framework to a local regime where gravity is intrinsically stronger.

#### H. A Unified Framework for Other Cosmological Tensions

Finally, the dual couplings  $G_{\text{Loc}}$  and  $G_{\text{Glob}}$  explain other major tensions:

- **Structure growth ( $S_8$  tension):** Since  $G_{\text{Glob}}/G_{\text{Loc}} \approx 0.83$ , the true growth rate is suppressed relative to  $\Lambda$ CDM predictions, explaining why LSS surveys infer lower  $S_8$  values than the CMB.
- **Gravitational lensing anomalies:** Strong lensing (local regime) agrees with  $G_{\text{Loc}}$ , while weak lensing (large-scale regime) reflects  $G_{\text{Glob}}$ . This dichotomy predicts a  $\sim 17\%$  systematic mass underestimate in weak lensing analyses, matching observed discrepancies.

In summary, the scale-dependent gravitational coupling derived from QEG provides a single, parameter-free mechanism that resolves the Hubble tension, the  $S_8$  tension, and anomalies in lensing, while simultaneously reinterpreting the dark sector.

## I. Derivation of the MOND Acceleration from First Principles

The phenomenological success of Modified Newtonian Dynamics (MOND) is centered on a fundamental acceleration scale,  $a_0 \approx 1.2 \times 10^{-10} \text{ m/s}^2$ . Within the QEG framework, we demonstrate that this is not an ad-hoc parameter, but an emergent property of the dissipative, elastic vacuum.

We identify  $a_0$  as the critical threshold where the vacuum's response to deformation transitions from being purely *coherent and elastic* (the regime of Newtonian gravity) to being dominated by *incoherent, dissipative dynamics* (the regime of the thermo-entropic field). At this precise transition point, we invoke a principle of *equipartition*: the substrate's capacity to respond to the deformation is equally partitioned between the elastic and dissipative channels.

Consequently, the effective stiffness,  $k_{\text{eff}}$ , available to sustain the *coherent* deformation is exactly half of the total geometric stiffness derived in Sec. II E. At the fiducial scale ( $r = 1 \text{ m}$ ), this effective stiffness is:

$$k_{\text{eff}} = \frac{1}{2} k_{\text{fiducial}} = \frac{1}{2} \left( \frac{1}{4\pi \cdot 1 \text{ m}} \right) = \frac{1}{8\pi \cdot 1 \text{ m}} \quad (263)$$

The fundamental acceleration scale is the rate at which the substrate can dynamically sustain this effective elastic tension, given by the product of this effective stiffness and the characteristic velocity of the dissipative mode,  $v_{\text{th}} = 1/c$  (Sec. XVI):

$$a_0 \equiv v_{\text{th}} \cdot k_{\text{eff}} = \frac{1}{c} \cdot \frac{1}{8\pi \cdot 1 \text{ m}} \quad (264)$$

Numerically, this yields a value of  $a_0 \approx 1.33 \times 10^{-10} \text{ m/s}^2$ , which is in remarkable agreement with the empirically determined MOND parameter.

*a. Physical Interpretation.* This result provides a first-principles origin for MOND. It suggests that MOND-like behavior is a manifestation of the vacuum's thermo-entropic dynamics becoming co-dominant with the elastic response. The constant  $a_0$  is the fundamental threshold of equipartition between coherence and dissipation in the spacetime substrate.

## J. Synthesis of Gravitational Modifications: A Two-Level Framework

The QEG framework predicts two distinct modifications to standard gravity, which operate in different regimes and resolve different observational puzzles. Rather than being contradictory, they form a complementary, two-level description of gravitational phenomena.

*a. Level 1: Scale-Dependent Coupling ( $G_{\text{loc}}$  vs.  $G_{\text{glob}}$ ).* As derived from the duality of self-energy (Sec. IX), the effective gravitational coupling constant itself depends on the scale and geometry of the source distribution. This principle governs the overall strength of gravity, explaining discrepancies between the local universe (governed by the stronger  $G_{\text{loc}}$ ) and the global cosmological background (governed by the weaker  $G_{\text{glob}}$ ). Its primary success is the parameter-free resolution of large-scale cosmological tensions, such as the Hubble and  $S_8$  tensions. This framework sets the "background" value of  $G$  for a

given system.

*b. Level 2: Acceleration-Dependent Dynamics ( $a_0$  and MOND).* Within a local regime where the coupling is already set to  $G_{\text{loc}}$ , a second modification emerges. As derived in Sec. XXVIII, when the local acceleration falls below the critical threshold  $a_0$ —a fundamental constant of the dissipative vacuum—the dynamical response of the substrate transitions from a purely elastic to a dissipative regime. This explains the anomalous rotation curves of galaxies without invoking dark matter.

*c. Conclusion: A Unified Picture.* There is no conflict between these two effects. The MOND phenomenology, governed by  $a_0$ , describes a change in the *dynamical law* at low accelerations *inside* a system (like a galaxy) which is itself governed by the stronger *coupling constant*  $G_{\text{loc}} \equiv G_N$ . Together, they provide a complete, self-consistent picture of gravity from galactic peripheries to the cosmic horizon, all emerging from the same underlying physics of a quantum-elastic, dissipative spacetime.

## XXIX. FINAL VALIDATION: FROM COVARIANT FORMALISM TO A MINIMAL GENERATIVE MODEL

The culmination of the QEG framework is the synthesis of all its derived relationships into a single, predictive mathematical object. This final validation proceeds in two stages. First, we present the complete, covariant **Unified Response Operator** in its 10-component (4x4) form, demonstrating that this rigorous formalism perfectly reproduces all the physical laws and constants derived in this paper. Second, we reveal that this complex structure is not fundamental, but emerges from a remarkably simple **Minimal Substrate Model** (a 3x3 matrix) based on first principles of symmetry. This not only validates the theory's internal consistency but also showcases its profound adherence to the principle of parsimony.

### A. Validation via Operator Synthesis: The Unified Response Tensor

The culmination of the QEG framework is the synthesis of all its derived relationships into a single, predictive mathematical object. Following a rigorous operator formalism, we can construct the **Unified Response Tensor**,  $K_{\mu\nu}$ , which dictates the substrate's complete elastic and dissipative response. This provides the ultimate validation of the theory, demonstrating that the entire structure of physical law emerges from a minimal set of generative principles.

The total response is the linear superposition of an elastic part and a dissipative part,  $K_{\mu\nu} = K_{\text{el}} + K_{\text{dis}}$ . Each part is obtained by factorizing the problem into its minimal components: a set of modal projectors and a diagonal matrix of physical weights. This is nothing but the modal restatement of the constitutive law

$$Q_i = k_i S_i, \quad (265)$$

already established in Eq. (37).

### 1. The Modal Projectors

The foundation of the structure is a complete and orthogonal set of projectors  $\{\Pi_A\}$  that decompose the space of symmetric tensors into the physically relevant subspaces. For QEG, these are:

- $\Pi_{00}$ : Projects onto the scalar, time-like compressive mode (static gravity).
- $\Pi_{0i}$ : Projects onto the vector, time-space torsional modes (electromagnetism).
- $\Pi_{\text{tr}}$ : Projects onto the scalar, isotropic spatial mode (thermo-entropics, trace of  $\mathcal{G}_{ij}$ ).
- $\Pi_{\text{shear}}$ : Projects onto the tensor, anisotropic spatial mode (gravitational waves, traceless part of  $\mathcal{G}_{ij}$ ).

These projectors satisfy  $\sum_A \Pi_A = \mathbb{I}_{\text{sym}}$  and  $\Pi_A \Pi_B = \delta_{AB} \Pi_A$ .

### 2. The Unified Response Tensor

The effective stiffness for each mode is encoded by applying a diagonal operator of “weights” in the basis of these projectors.

*a. The Elastic Response ( $K_{el}$ ).* The elastic part is governed by the baseline inertial stiffness  $\mu_0$  and the causal modulator  $c$ . The operator is constructed by assigning the derived elastic stiffness to each modal projector:

$$K_{el} = \underbrace{(\mu_0 c^2 \cdot \Pi_{0i})}_{\text{Electromagnetism (Stiff)}} + \underbrace{\left(\frac{\mu_0}{c^2} \cdot \Pi_{\text{tr}}\right)}_{\text{Thermo-entropics (Compliant)}}. \quad (266)$$

This operator correctly assigns the maximum stiffness ( $\propto c^2$ ) to the transverse EM mode and the maximum compliance ( $\propto c^{-2}$ ) to the longitudinal thermo-entropic mode.

*b. The Dissipative Response ( $K_{dis}$ ).* The dissipative part is governed by the baseline stiffness  $\mu_0$  and the universal damping coefficient  $\alpha$ . As derived, gravity is a second-order dissipative phenomenon. Therefore, its stiffness is assigned to the gravitational projectors:

$$K_{dis} = \underbrace{(\mu_0 \alpha^2 \cdot \Pi_{00})}_{\text{Static Gravity}} + \underbrace{(\mu_0 \alpha^2 \cdot \Pi_{\text{shear}})}_{\text{Gravitational Waves}}. \quad (267)$$

Strictly speaking, the dissipative operator is of telegraph type (hyperbolic–parabolic), but  $\mu_0 \alpha^2$  captures its static limit consistently.

*c. The Total Response Operator.* The complete Unified Response Tensor is the sum of the elastic and dissipative parts:

$$K_{\mu\nu} = \left( \mu_0 c^2 \Pi_{0i} + \frac{\mu_0}{c^2} \Pi_{\text{tr}} \right) + \left( \mu_0 \alpha^2 \Pi_{00} + \mu_0 \alpha^2 \Pi_{\text{shear}} \right) \quad (268)$$

### 3. Reproduction of Physical Equations and New Predictions

This operator formalism reproduces all constitutive relations derived in this paper. For any deformation

state, represented by a vector  $|\psi\rangle$  in the space of symmetric tensors, the resulting physical interaction is governed by the eigenvalue of the response operator:

$$K_{\mu\nu} |\psi\rangle = \lambda |\psi\rangle. \quad (269)$$

- If  $|\psi\rangle$  lies in the subspace of  $\Pi_{0i}$ , the operator returns the electromagnetic stiffness:  $K_{\mu\nu} |\psi\rangle = \mu_0 c^2 |\psi\rangle$ .
- If  $|\psi\rangle$  lies in the subspace of  $\Pi_{00}$ , the operator returns the gravitational stiffness:  $K_{\mu\nu} |\psi\rangle = \mu_0 \alpha^2 |\psi\rangle$ .
- The same holds for thermo-entropic and shear-gravitational modes, reproducing all the derived values of  $G$ ,  $K_e$ , and  $k_B$ .

*a. New Predictions.* This formalism does not yield new numerical constants, but it makes a profound prediction: the **structural uniqueness and rigidity of physical law**. It implies that there can be only four fundamental types of long-range interaction, corresponding to the four orthogonal projectors. Their strengths are not independent but are rigidly locked by the operator structure. This opens the door to systematically calculating higher-order corrections (mixing between projectors) and analyzing the running of the constants under the renormalization group.

### B. The Underlying Simplicity: A Minimal Generative Model

Having established the validity of the complete 10-component operator, we now demonstrate that this entire structure is not fundamental, but can be generated from a far simpler model based on first principles of symmetry and parsimony. To analyze the substrate’s response, we distinguish between the full 10-component unified tensor  $\mathcal{G}_{\mu\nu}$  and its purely spatial deformation components. We define a *Symmetric State Tensor*,  $S_{ij}$ , as the 3x3 matrix representation of the spatial block  $\mathcal{G}_{ij}$ . This object is not an abstract source, but represents the physical “shape” of the spatial deformation at a given point. Any such symmetric tensor can be uniquely and irreducibly decomposed into two fundamental modes:

- **Isotropic (Radial) Modes:** These phenomena represent spherically symmetric, compressive/decompressive deformations of the substrate, like a “breathing” motion. They have no intrinsic directionality other than the radial line from the source. This family includes **Electrostatics**, **Newtonian Gravity**, and the diffusive flow of the **Thermo-Entropic Field**.
- **Anisotropic (Torsional/Circulatory) Modes:** These phenomena represent deformations with an intrinsic directionality or orientation, such as shear or rotation. This family includes **Magnetism**, **Gravitational Waves**, and other torsional effects like frame-dragging.

This physical dichotomy is directly mapped to the mathematical structure of the unified tensor, providing a rigorous, non-arbitrary classification of all physical law. The response modulator matrices (which we denote  $M_e$  and  $M_a$ ) act upon these intrinsic modes of the state tensor  $S$  to produce the observed hierarchy of forces.

*Derivation of the Minimal Modulator Matrices from First Principles*

We seek the simplest law for the response coefficients consistent with the substrate's fundamental symmetries: homogeneity, isotropy, index-shift equivariance, and left/right separability ( $K_{ij} = a_i b_j$ ). As a mathematical theorem, the only non-trivial form satisfying these conditions is an exponential scaling law,  $K_{ij} \propto \chi^{j-i}$ <sup>9</sup>. This rule is not a postulate but a deduction from symmetry, yielding two **Dimensionless Modulator Matrices**:

$$M_c = \begin{pmatrix} 1 & c & c^2 \\ c^{-1} & 1 & c \\ c^{-2} & c^{-1} & 1 \end{pmatrix}, \quad M_\alpha = \begin{pmatrix} 1 & \alpha & \alpha^2 \\ \alpha^{-1} & 1 & \alpha \\ \alpha^{-2} & \alpha^{-1} & 1 \end{pmatrix}. \quad (270)$$

*a. Dimensional consistency.* The scales  $c$  and  $\alpha$  within these matrices are treated as pure numbers. All physical dimensions are carried by the baseline prefactors.

*The Constitutive Law: Superposition of Responses*

The total response of the substrate to a symmetric deformation state  $S$  is a linear superposition of the elastic and dissipative channels, analogous to the Kelvin-Voigt model. The observable field,  $\mathcal{Q}$ , is generated by applying the Green's operator,  $\mathcal{G}$ , to the sum of the internal source responses:

$$\mathcal{Q} = \mathcal{G} \left[ \mu_0 \text{Sym}(M_c \odot S) + \eta_0 \text{Sym}(M_\alpha \odot S) \right] \quad (271)$$

where  $\mu_0$  is the dimensionless baseline elastic strength,  $\eta_0$  is the baseline dissipative strength,  $\odot$  is the Hadamard product, and  $\text{Sym}[X] := \frac{1}{2}(X + X^\top)$  ensures a symmetric response. The operator  $\mathcal{G}$  represents the appropriate Green's function for the d'Alembertian, which in the static 3D limit becomes the Laplacian kernel  $\mathcal{G}(r) = 1/(4\pi r)$ .

*Additive Law vs. Multiplicative Couplings.*

The constitutive law (Eq. 271) is fundamentally *additive*. However, when projected onto a single modal subspace  $A$ , the effective scalar coupling  $\kappa_A$  can be factorized into a multiplicative form:

$$\kappa_A(\omega) = \mu_0 c^{\Delta_A} \left[ 1 + \Lambda_A(\omega) \right], \quad \Lambda_A(\omega) = g(\omega) \frac{\eta_0}{\mu_0} \left( \frac{\alpha}{c} \right)^{\Delta_A}$$

This rigorously shows why the measured effective constants (like  $G \propto \eta_0 \alpha^2$ ) appear as products, even though the underlying operator law is a sum.

<sup>9</sup> *Lemma (Toeplitz-separable  $\Rightarrow$  exponential weights).* Under index-shift equivariance and left/right separability,  $K_{ij} = a_i b_j$  with  $K_{i+r, j+r} = K_{ij}$  implies  $K_{ij} = \text{const} \cdot \chi^{j-i}$  for a constant  $\chi$  independent of  $i, j$ . Hence the coefficient masks are uniquely (up to normalization) of the form  $\chi^{j-i}$ .

*Recovery of Physical Constants and Predictions*

This minimal formalism reproduces all static scales as channel-dominated limits of the law. The even-powered exponents ( $\Delta = 0, \pm 2$ ) correspond to isotropic, radial, stable, long-range forces (EM, Thermo-entropics, Gravity), while the odd-powered exponents ( $\Delta = \pm 1$ ) are correctly identified as anisotropic, torsional, dynamical, short-range couplings, such as the vacuum impedance  $Z_0 \propto \mu_0 c$ :

- **Electrostatics** ( $K_e \propto \mu_0 c^2$ ): The strongest static force, arising from the stiff pole ( $\Delta = 2$ ) of the elastic channel acting on an isotropic state.
- **Thermo-Entropics** ( $k_B \propto \mu_0 c^{-2}$ ): The weakest elastic force, from the compliant pole ( $\Delta = -2$ ) of the elastic channel on an isotropic state.
- **Gravity** ( $G \propto \eta_0 \alpha^2$ ): A static force arising from the stable, second-order dissipative response ( $\Delta = 2$ ) on an isotropic state. We unify the framework by identifying  $\eta_0 \equiv \mu_0 \propto \alpha^3$  151.
- **Magnetism** ( $Z_0 = \mu_0 c$ ): A transient, dynamic coupling arising from the first-order elastic response ( $\Delta = 1$ ) on an anisotropic state.

It is crucial to distinguish between the physical state of the substrate and the laws that govern it. The deformation itself is described by a *tensor field*, such as the symmetric state  $S_{ij}$ , which has a value at each point in spacetime. The response matrices,  $M_c$  and  $M_\alpha$ , are not tensors in this sense. They are **operators** that represent the substrate's intrinsic constitutive law. Their indices do not span spacetime coordinates, but rather an abstract "modal space" that captures the different types of response (e.g., maximally stiff, baseline, maximally compliant). The Hadamard product ( $\odot$ ) is the mathematical tool that describes how this abstract rulebook (the matrix) acts upon the concrete physical state (the tensor) to determine the outcome of an interaction. This formalism allows us to separate the universal rules of the substrate from its specific, local deformation.

**C. Discussion: Two Formalisms, One Physics**

This two-tiered validation opens a profound discussion. The 4x4 operator formalism provides the complete, covariant description necessary to map interactions to the geometry of spacetime. However, the 3x3 minimal model demonstrates that the underlying "genetic code" of physical law is far simpler and is rigidly constrained by symmetry.

This suggests that the 4x4 formalism is the necessary "phenotypic" expression of the underlying "genotypic" simplicity of the 3x3 structure. The question of which representation is more "fundamental" is a deep one. QEG suggests that the truest answer lies in their synthesis: the universe operates on the simplest possible principles (the 3x3 model), which necessarily manifest in the complete covariant structure (the 4x4 operator) required by the geometry of spacetime. This dual perspective provides a powerful, self-consistent, and remarkably simple foundation for a unified theory of physics.

As a final note, the Causal-Dissipative Equivalence  $c \propto \alpha^{-4}$  XIX M reveals that the speed of light is an emergent property determined by the vacuum's quantum friction. A universe with less friction ( $\alpha \rightarrow 0$ ) would have a faster speed of light

( $c \rightarrow \infty$ ). The existence of a finite cosmic speed limit is a direct consequence of the fact that spacetime is a dissipative medium. The value of  $\alpha$  sets the value of  $c$ , unifying causality and quantum dissipation into a single, profound principle. Therefore, the minimum generative model could be simplified even more, just to the simplest geometric tool (the Laplacian operator  $\nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ ) and the simplest geometric dissipation dictated by symmetry, isotropy and auto-interactions ( $\alpha \equiv \frac{1}{16\pi\sqrt{\frac{3}{5}}4\pi} 140$ ).

### XXX. FINAL CONCLUSIONS

#### A. A Coherent and Predictive Framework from First Principles

This work has introduced Quantum-Elastic Geometry (QEG), a unified framework built upon a small set of foundational axioms: the universe consists of a single, unified substrate—spacetime itself—which is described as a quantum, elastic, and dissipative medium. From these physically-motivated first principles, we have demonstrated that a remarkably coherent and internally consistent picture of fundamental physics emerges.

The theory's primary strength lies not in predicting constants *ex nihilo*, but in revealing the inevitable network of relationships that constrains them. By postulating a single, unified field  $\mathcal{G}_{\mu\nu}$ , we have shown that:

1. **The Laws of Physics are Emergent:** General Relativity and Electromagnetism are not independent theories, but the macroscopic manifestations of the compressive and torsional deformation modes of the substrate, respectively. The theory further predicts a third, thermodynamic field from the substrate's spatial modes, providing a first-principles origin for irreversible thermodynamics.
2. **Fundamental Constants are Interdependent:** Constants such as  $G$ ,  $\hbar$ , and  $\alpha$  are not arbitrary inputs, but are uniquely determined by the substrate's properties (stiffness  $\kappa$ , dissipation  $\alpha$ ) and by the stringent requirements of internal consistency. The successful parameter-free prediction of  $\alpha$  and the resolution of cosmological tensions like the Hubble crisis serve as powerful, non-trivial validations of the framework.
3. **Sources are Endogenous:** Mass and charge are not external entities that deform spacetime, but are themselves specific, stable configurations of deformed spacetime.

In summary, QEG presents a paradigm in which the universe is not a collection of disparate laws and constants, but a deeply interconnected system whose entire structure is rigorously determined by the properties of a single underlying entity.

#### B. Correspondence with Formal Gauge-Theoretic Frameworks

The physical principles of QEG find a profound conceptual correspondence in the formal gauge theory of Unified Gravity (UG) proposed by Partanen and Tulkki [23]. While QEG proceeds from physical and

geometric principles, UG arrives at a similar vision from the mathematical rigor of gauge theory, successfully deriving a dimensionless gravitational coupling and demonstrating one-loop renormalizability.

The correspondence is striking: QEG's "modal projections" of the  $\mathcal{G}_{\mu\nu}$  tensor can be seen as the physical manifestation of UG's distinct  $U(1)$  gauge symmetries. Furthermore, UG provides a formal, first-principles derivation of the stress-energy-momentum tensor as the source for gravity, justifying the physical coupling at the heart of QEG. This suggests that the UG formalism may provide the rigorous, renormalizable quantum field theory for which the QEG framework lays the physical and conceptual foundation, lending significant theoretical support to the idea that the principles and derivations of QEG reflect a deep, symmetric structure of our universe.

#### C. Broader Implications: The Torsional Substrate and the Cosmic Spin Anomaly

Beyond its ability to unify fields and constants, the QEG framework offers a novel perspective on outstanding observational puzzles that challenge the standard cosmological model. One such puzzle is the recently reported evidence for a large-scale spin asymmetry in the universe, suggesting a preferred axis or "cosmic spin" [65, 66]. Such an observation, if confirmed, would represent a profound violation of the cosmological principle of isotropy and would be difficult to reconcile with the  $\Lambda$ CDM model.

The QEG framework, however, not only accommodates this possibility but provides a natural physical mechanism for it. The connection arises from three of the theory's core tenets:

1. **Primordial Torsional Modes:** The unified field  $\mathcal{G}_{\mu\nu}$  possesses torsional modes ( $\mathcal{G}_{ij}$ ) as a fundamental component of its structure. It is therefore entirely plausible that the initial state of the universe contained a net primordial angular momentum, a coherent torsion embedded within the substrate itself. The observed spin asymmetry in galaxies could be a "fossil" of this initial condition, a faint remnant of the spacetime fabric's own intrinsic rotation.
2. **Inertia of the Vacuum:** As derived in Sec. XXVII, the cosmological constant itself can be interpreted as the density of the vacuum's rotational inertia. A non-zero primordial spin is a natural consequence of a substrate possessing such an intrinsic property. The observed cosmic axis would then correspond to the axis of this primordial inertia.
3. **Intrinsic Dissipation and the Fading Anisotropy:** The most elegant aspect of this explanation lies in its consistency with the extreme isotropy of the Cosmic Microwave Background (CMB). The QEG substrate is not only elastic but also inherently *dissipative*, possessing an intrinsic "viscosity" (Sec. VII). This property ensures that any primordial anisotropy, such as a large-scale rotation, would be naturally and inexorably damped over cosmic time.

This leads to a compelling narrative: the universe may have been born with a significant spin, but the dissipative nature of spacetime has smoothed out this anisotropy over 13.8 billion years. What we would observe today is not a violently rotating cosmos, but a nearly isotropic one, retaining only a ghostly, statistically subtle residue of its primordial spin—precisely

what the observational anomaly suggests. Thus, the potential discovery of a cosmic axis, far from being a crisis for QEG, would stand as one of the most powerful pieces of observational evidence for the torsional, dissipative, and elastic nature of the spacetime substrate itself.

#### D. A Final Word on the Structure of Physical Law: The Pond and the Rulebook

At its core, the synthesis presented in this paper can be understood through a simple yet powerful physical analogy: the universe is a vast, dynamic medium—the spacetime substrate—akin to the surface of a pond. The laws of physics are not arbitrary rules imposed upon this medium, but are the very description of its intrinsic properties and how waves form and propagate within it.

##### *The Tensor as the State of the Pond.*

The physical state of the universe at any point is described by the deformation tensor,  $\mathcal{G}_{\mu\nu}$  or its simplified form, the state tensor  $S_{ij}$ . This tensor is the **pond itself**. Its components describe the precise shape of the water's surface at every location: where there are peaks, where there are troughs, how steep the slopes are. It is a dynamic, living object—a geometric description of "*what is happening*" in the substrate.

##### *The Matrices as the Rulebook of Water.*

The minimal response matrices,  $M_c$  and  $M_\alpha$ , are not part of the pond's surface. They are the **rulebook** that describes the fundamental properties of the "*water*". They are not a tensor in spacetime, but an abstract operator that answers questions like: How much does this water resist being compressed? (Stiffness). How quickly do waves die out? (Viscosity). This rulebook is universal and the same everywhere. Its structure is not arbitrary but is a direct consequence of fundamental symmetries, as we have shown.

- The **Elastic Matrix** ( $M_c$ ) is the chapter on the water's "tension and density". It catalogues the different responses based on the causal speed,  $c$ : the immense resistance to creating a static peak ( $\propto c^2$ ), the impedance to a traveling wave ( $\propto c$ ), and the profound ease of a slow, global tide ( $\propto c^{-2}$ ).
- The **Dissipative Matrix** ( $M_\alpha$ ) is the chapter on the water's "viscosity". It catalogues the different ways the pond dissipates energy, governed by the damping coefficient,  $\alpha$ . It describes the weak, second-order friction that governs the most heavily suppressed phenomena, like gravity ( $\propto \alpha^2$ ).

##### *The Laplacian as Universal Syntax, the Tensor as Physical Semantics.*

The emergence of all physical law from this foundation is the interplay of three elements:

1. **The Source** ( $J_{\mu\nu}$ ): The "stone" thrown into the pond.

2. **The Universal Rule of Propagation (The Laplacian,  $\nabla^2$ )**: This is the universal "syntax" of the universe, a rule of geometry dictated by isotropy. It governs the  $1/r$  shape of all long-range potentials, just as geometry dictates that waves in the pond will be circular.

3. **The Character of the Interaction (The Unified Response,  $K_{\mu\nu}$ )**: This is the "semantics" of the universe, constructed from the rulebook matrices. It does not change the geometric shape of the interaction, but it determines its **amplitude** (the elastic part) and its **duration** (the dissipative part).

In this view, the rich tapestry of physical law is not ad-hoc. The immense strength of electromagnetism, the subtle nature of thermodynamics, and the weak, persistent presence of gravity are the direct and inevitable consequences of the interplay between the universal syntax of geometry and the specific semantic properties of our universe's elastic and dissipative fabric.

#### E. A Final Word on the Nature of Quantum-Elastic Geometry

At its core, Quantum-Elastic Geometry is the logical conclusion of viewing the universe as a single, unified entity. If spacetime is the only true substance, then all phenomena must be manifestations of its properties. QEG formalizes this by modeling spacetime as a quantum elastic medium, a postulate justified by the fundamentally oscillatory, wave-like nature of everything that exists. Its capacity for different types of vibration gives rise to the distinct interaction modes we call gravity (compressive), electromagnetism (torsional), and thermodynamics (diffusive), which differ not in substance, but only in topology and strength.

Perhaps the most profound shift is the reinterpretation of physical sources. In QEG, entities like mass and charge are not external agents that deform spacetime; they are localized, stable deformations of spacetime. This leads to the ultimate insight of the Hookean framework: because the classical "sources" are now understood as the "deformations", the true driving term in the universe's constitutive law becomes the *action* associated with stabilizing that deformation. In this view, all of physics above the Planck scale is unified under a single, supreme elastic law, where the geometry of spacetime itself acts, reacts, and resonates in response to the flow of quantum action.

In summary, the results of Quantum-elastic geometry challenge our notions of what is fundamental in the universe. If gravity, electromagnetism, and quantum phenomena all arise from the same oscillatory vacuum, then the distinction between these forces are more illusory than real. They are just expressions of the same underlying reality, a vibrating cosmos that resonates through every level of existence—from the quantum realm to the largest cosmic structures.

This realization suggests that the universe is not a fragmented collection of forces and constants, but a deeply interconnected whole, where every phenomenon is an expression of the same underlying dynamics, and divisions between forces and fields are merely artifacts of our limited understanding, and where every aspect of reality is a manifestation of the same fundamental processes.

Quantum-elastic geometry also resonates with

the philosophical principle of simplicity, or "Occam's Razor", which suggests that the simplest explanation that accounts for all phenomena is likely to be correct. The notion that the universe's complexity—spanning from quantum mechanics to general relativity—can be fundamentally explained through the dynamics of vacuum oscillations provides a powerful example of how simplicity can reveal profound truths.

The metaphysical vision offered as a byproduct by this model invites us to reconsider the nature of the universe as a whole. It suggests a cosmos that is not a static structure governed by immutable laws but a dynamic, evolving system where everything is

interconnected. This invites a more holistic view of the cosmos, where complexity and diversity arise from simple, fundamental vibrations at the heart of reality itself.

#### Declaration of generative AI and AI-assisted technologies in the writing process

*During the preparation of this work the author used Generative AI to improve the quality of the narrative of the Paper, and peer-review and check the internal consistency of the theoretical derivations. After using this tool/service, the author reviewed and edited the content as needed, and takes full responsibility for the content of the publication.*

- 
- [1] Steven Weinberg. *Dreams of a Final Theory: The Scientist's Search for the Ultimate Laws of Nature*. Knopf Doubleday Publishing Group, 2011.
- [2] John F. Donoghue, Eugene Golowich, and Barry R. Holstein. *Dynamics of the Standard Model*. Cambridge University Press, 1994.
- [3] Albert Einstein. The foundation of the general theory of relativity. *Annalen der Physik*, 354(7):769–822, 1916.
- [4] Robert M. Wald. *General Relativity*. University of Chicago Press, Chicago, 1984.
- [5] Sean M. Carroll. *Spacetime and Geometry: An Introduction to General Relativity*. Addison-Wesley, 2004.
- [6] Steven Weinberg. The cosmological constant problem. *Reviews of Modern Physics*, 61(1):1–23, 1989.
- [7] Sean Carroll. *Dark Matter, Dark Energy: The Dark Side of the Universe*. The Teaching Company, 2007.
- [8] V. Trimble. Existence and nature of dark matter in the universe. *Annual Review of Astronomy and Astrophysics*, 25:425–472, 1987.
- [9] Carlo Rovelli and Lee Smolin. Loop space representation of quantum general relativity. *Nuclear Physics B*, 331(1):80–152, 1988.
- [10] Ted Jacobson. Thermodynamics of spacetime: The einstein equation of state. *Physical Review Letters*, 75(7):1260–1263, 1995.
- [11] Erik P. Verlinde. On the origin of gravity and the laws of newton. *Journal of High Energy Physics*, 2011(4):29, 2011.
- [12] S.W. Hawking. Particle creation by black holes. *Communications in Mathematical Physics*, 43(3):199–220, 1975.
- [13] Shinsei Ryu and Tadashi Takayanagi. Holographic Derivation of Entanglement Entropy from AdS/CFT. *Phys. Rev. Lett.*, 96:181602, 2006.
- [14] Mark Van Raamsdonk. Building up spacetime with quantum entanglement. *Gen. Rel. Grav.*, 42:2323–2329, 2010.
- [15] A. Belfiglio, O. Luongo, S. Mancini, and S. Tomasi. Entanglement entropy evolution during gravitational collapse. 2025.
- [16] Carlos Barceló, Stefano Liberati, and Matt Visser. Analogue Gravity. *Living Rev. Rel.*, 14:3, 2011.
- [17] W. G. Unruh. Experimental black-hole evaporation? *Phys. Rev. Lett.*, 46:1351–1353, 1981.
- [18] Michael E. Peskin and Daniel V. Schroeder. *An Introduction to Quantum Field Theory*. Addison-Wesley, Reading, Mass., 1995.
- [19] Carlo Rovelli. *Quantum Gravity*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 2004.
- [20] Fay Dowker. Introduction to causal sets and their phenomenology. *General Relativity and Gravitation*, 45:1651–1667, 2013.
- [21] Ted Jacobson and David Mattingly. Gravity with a dynamical preferred frame. *Phys. Rev. D*, 64:024028, 2001.
- [22] V. Alan Kostelecký and Neil Russell. Data Tables for Lorentz and CPT Violation. *Rev. Mod. Phys.*, 83:11–31, 2011.
- [23] Mikko Partanen and Jukka Tulkki. Gravity generated by four one-dimensional unitary gauge symmetries and the standard model. *Reports on Progress in Physics*, 88(5):057802, 2025.
- [24] Jean-Philippe Uzan. Varying constants, gravitation and cosmology. *Living Rev. Rel.*, 14:2, 2011.
- [25] R. R. Cuzinatto, R. F. L. Holanda, and S. H. Pereira. Observational constraints on varying fundamental constants in a minimal CPC model. *Eur. Phys. J. C*, 82(6):516, 2022.
- [26] S. Deser. Self-interaction and gauge invariance. *Annals of Physics*, 59:248–253, 1970.
- [27] D. Lovelock. The einstein tensor and its generalizations. *Journal of Mathematical Physics*, 12:498–501, 1971.
- [28] P. J. Mohr, B. N. Taylor, and D. B. Newell. Fine-structure constant. *National Institute of Standards and Technology (NIST)*, 2019.
- [29] Steve Long. Resonant circuits – resonators and q. Course notes for ECE145B / ECE218B, Electrical & Computer Engineering, U.C. Santa Barbara, 2004.
- [30] B. J. van Wees, H. van Houten, C. W. J. Beenakker, J. G. Williamson, L. P. Kouwenhoven, D. van der Marel, and C. T. Foxon. Quantized conductance of point contacts in a two-dimensional electron gas. *Physical Review Letters*, 60(9):848–850, 1988.
- [31] D. A. Wharam, T. J. Thornton, R. Newbury, M. Pepper, H. Ahmed, J. E. F. Frost, D. G. Hasko, D. C. Peacock, D. A. Ritchie, and G. A. C. Jones. One-dimensional transport and the quantisation of the ballistic resistance. *Journal of Physics C: Solid State Physics*, 21(8):L209–L214, 1988.

- [32] Grant R. Fowles and George L. Cassiday. *Analytical Mechanics*. Thomson Brooks/Cole, 7th edition, 2005.
- [33] David J. Griffiths and Darrell F. Schroeter. *Introduction to Quantum Mechanics*. Cambridge University Press, Cambridge, UK, 3rd edition, 2018.
- [34] John David Jackson. *Classical Electrodynamics*. John Wiley & Sons, 3rd edition, 1999.
- [35] Jan R. Westra, Chris J. M. Verhoeven, and Arthur H. M. van Roermund. *Oscillators and Oscillator Systems: Classification, Analysis and Synthesis*. Springer Science & Business Media, 2007.
- [36] David M. Pozar. *Microwave Engineering*. Wiley, 4th edition, 2011.
- [37] W. K. H. Panofsky and M. Phillips. *Classical Electricity and Magnetism*. Addison-Wesley, 1962.
- [38] Edwin F. Taylor and John Archibald Wheeler. *Spacetime Physics: Introduction to Special Relativity*. W. H. Freeman, 2nd edition, 1992.
- [39] G. E. Volovik. *The Universe in a Helium Droplet*, volume 117 of *International Series of Monographs on Physics*. Oxford University Press, 2003.
- [40] T. Padmanabhan. Thermodynamical aspects of gravity: New insights. *Reports on Progress in Physics*, 73(4):046901, 2010.
- [41] Erik P. Verlinde. On the origin of gravity and the laws of newton. *Journal of High Energy Physics*, 2011(4):29, 2011.
- [42] Richard P. Feynman, Robert B. Leighton, and Matthew Sands. The feynman lectures on physics, volume ii, chapter 8: Electrostatic energy, 1964.
- [43] Planck Collaboration. Planck 2015 results. xiii. cosmological parameters. *Astronomy & Astrophysics*, 594:A13, 2016.
- [44] J. Abadie et al. Search for gravitational waves from low mass compact binary coalescence in ligo's sixth science run and virgo's science runs 2 and 3. *Physical Review D*, 85(8):082002, 2012.
- [45] P. C. W. Davies. Thermodynamics of black holes. *Reports on Progress in Physics*, 41(8):1313–1355, 1978.
- [46] S. W. Hawking and R. Penrose. The singularities of gravitational collapse and cosmology. *Proceedings of the Royal Society A*, 314(1519):529–548, January 1970.
- [47] C. W. F. Everitt and D. B. et al. DeBra. Gravity probe b: Final results of a space experiment to test general relativity. *Physical Review Letters*, 106(22):221101, 2011.
- [48] H. Julian Goldsmid. *Introduction to Thermoelectricity*, volume 121 of *Springer Series in Materials Science*. Springer, 2010.
- [49] W. Israel. Nonstationary irreversible thermodynamics: A causal relativistic theory. *Annals of Physics*, 100:310–331, 1976.
- [50] W. Israel and J. M. Stewart. Transient relativistic thermodynamics and kinetic theory. *Annals of Physics*, 118:341–372, 1979.
- [51] C. Cattaneo. Sulla conduzione del calore. *Atti del Seminario Matematico e Fisico dell'Università di Modena*, 3:83–101, 1948.
- [52] P. Vernotte. Some possible complications in the phenomenological theory of heat conduction. *Comptes Rendus*, 246:3154–3155, 1958.
- [53] William G. Unruh. Notes on black-hole evaporation. *Physical Review D*, 14(4):870–892, 1976.
- [54] H. B. G. Casimir. On the attraction between two perfectly conducting plates. *Proc. K. Ned. Akad. Wet.*, 51:793–795, 1948.
- [55] M. Bordag, U. Mohideen, and V. M. Mostepanenko. New developments in the casimir effect. *Physics Reports*, 353(1–3):1–205, 2001.
- [56] S. K. Lamoreaux. Demonstration of the casimir force in the 0.6 to 6  $\mu\text{m}$  range. *Physical Review Letters*, 78(1):5–8, 1997.
- [57] G. Bressi, G. Carugno, R. Onofrio, and G. Ruoso. Measurement of the casimir force between parallel metallic surfaces. *Physical Review Letters*, 88(4):041804, 2002.
- [58] Albert Einstein. The field equations of gravitation. *Sitzungsberichte der Preussischen Akademie der Wissenschaften zu Berlin*, pages 844–847, 1915.
- [59] David Hilbert. Die Grundlagen der Physik. *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse*, pages 395–407, 1915.
- [60] Alexander Friedmann. On the curvature of space. *Zeitschrift für Physik*, 10:377–386, 1922.
- [61] P.J.E. Peebles. *Principles of Physical Cosmology*. Princeton University Press, 1993.
- [62] LibreTexts Physics Team. 7.5: The friedmann equation. *Physics LibreTexts*, 2020.
- [63] Lucas M. Macri et al. Adam G. Riess. A comprehensive measurement of the local value of the hubble constant with 1 km/s/mpc uncertainty from the hubble space telescope and the sh0es team. *The Astrophysical Journal Letters*, 934(1):L7, 2022.
- [64] Planck Collaboration. Planck 2018 results. vi. cosmological parameters. *Astronomy & Astrophysics*, 641:A6, 2020.
- [65] L. Shamir. Large-scale patterns of galaxy spin directions: review and data analysis. *Universe*, 8(8):397, 2022.
- [66] L. Shamir. Galaxy spin directions show a dipole axis in the sky. *Astronomische Nachrichten*, 343(6):e20220010, 2022.