

Lake Aquaculture for Catastrophic Food Security

Jaba Tkemaladze *¹

¹ Director of Research, Longevity Clinic, Inc, Georgia

Abstract

The intensifying frequency of climate disasters, geopolitical conflicts, and pandemics exposes critical vulnerabilities in globalized, input-intensive food systems. Traditional protein sources—terrestrial livestock, crops, and marine fisheries—are highly susceptible to collapse under such catastrophic scenarios due to their dependencies on complex supply chains, external inputs, and stable climatic conditions. This article posits that lake-based aquaculture represents a strategically undervalued yet indispensable component of a resilient food security framework. We argue that the inherent characteristics of lacustrine systems—including superior feed conversion ratios, the utilization of natural trophic pathways, and a static "live storage" production model—confer a unique capacity to function autonomously during prolonged infrastructural and logistical breakdowns. The analysis delineates criteria for selecting resilient fish species, advocates for extensive polyculture management models, and outlines strategies for mitigating risks related to disease, genetic resource security, and ecological degradation. Furthermore, a strategic roadmap is proposed for integrating this approach into national policy, emphasizing legislative action, targeted research, economic incentives, and specialized education. The conclusion asserts that proactive investment in developing lake aquaculture as a decentralized protein reserve is a critical imperative for enhancing national food sovereignty and long-term survivability in an era of escalating systemic risks.

Keywords: Food Security, Protein Security, Catastrophe Resilience, Lake Aquaculture, Sustainable Aquaculture, Disaster Preparedness, Polyculture, Resource Efficiency, Climate Adaptation.

Introduction

The Urgency of the Problem

The foundational systems of global food production, predicated on intensive agriculture and industrialized livestock farming, are demonstrating profound vulnerabilities. This complex, just-in-time network is a marvel of modern efficiency, yet it is increasingly strained by a confluence of systemic shocks. The Intergovernmental Panel on Climate Change (IPCC) has consistently documented an increase in the frequency and intensity of extreme weather events, which devastate crops, disrupt planting seasons, and threaten livestock (IPCC, 2022). Beyond environmental factors, the global food system is acutely sensitive to geopolitical strife, as conflicts can abruptly halt the trade of essential inputs like fertilizers, pesticides, and animal

feed, crippling agricultural productivity in import-dependent nations (Godfray et al., 2010). Furthermore, the recent COVID-19 pandemic served as a stark reminder of how biological threats can disrupt labor forces and fracture intricate supply chains, leading to localized shortages and price volatility (Laborde et al., 2020). These cascading risks—climatic, geopolitical, and biological—paint a picture of a global food system that is inherently fragile and ill-prepared for prolonged, multi-faceted catastrophes.

The core of this fragility lies in the resource-intensive nature of terrestrial protein production. Conventional livestock farming, particularly for ruminants, is a notoriously inefficient process for generating edible protein. It requires vast tracts of land for grazing and feed crop cultivation, consumes enormous quantities of freshwater, and is a significant source of greenhouse gases (Herrero et al., 2013). The reliance on concentrated animal feeding operations (CAFOs) also creates hotspots for zoonotic disease emergence and spread, presenting a persistent pandemic risk (Jones et al., 2013). When supply chains for feed, energy, and veterinary services are compromised, these centralized systems face rapid collapse.

The Gap in Current Research

In response to these threats, national and international strategies for food security have traditionally prioritized two pillars: the maintenance of strategic grain reserves and the support of terrestrial livestock sectors (FAO, 2021). While the importance of caloric security through cereal stocks is undeniable, this approach overlooks a critical component of human nutrition: high-quality protein. Protein is essential not only for basic physiological function but also for immune competence and long-term health, attributes that become paramount in survival and recovery scenarios (Wu, 2016).

A significant research gap exists in the systematic exploration of aquatic biosystems as a resilient and efficient source of this vital nutrient during crises. The academic and policy discourse on aquaculture has largely focused on its role in meeting the growing protein demands of a rising global population under business-as-usual conditions (FAO, 2020). However, its strategic value as a backstop or a primary resilient system in a catastrophic, de-globalized context remains underexplored. The potential of lake-based aquaculture, in particular, is marginalized in favor of marine capture fisheries—which are themselves vulnerable to climate change and logistical disruption—or intensive, land-based recirculating aquaculture systems (RAS) that share the same dependencies on external energy and feed inputs as terrestrial farming (Béné et al., 2016). There is a critical need to shift the narrative and investigate food production models that are not merely efficient in times of stability, but are robust and fail-safe in times of collapse.

Aim of the Article

This article aims to address this critical gap by positing and rigorously arguing for the strategic role of lake-based aquaculture as a cornerstone of national food and protein security in the event of a long-term, systemic catastrophe. We contend that lacustrine (lake-based) aquaculture systems possess inherent characteristics—such as low external resource dependency, high energy conversion efficiency, and ecological buffering capacity—that make them uniquely suited to withstand and thrive in conditions that would paralyze conventional agriculture. The objective is not to present aquaculture as a mere supplement, but to reconceptualize it as a vital, resilient pillar of a comprehensive catastrophic food security strategy.

Research Objectives

To achieve this aim, the article will pursue the following specific research objectives:

1. To conduct a comparative analysis of protein production efficiency between lake aquaculture and traditional terrestrial livestock husbandry. This analysis will focus on key metrics such as Feed Conversion Ratio (FCR), water footprint, and land use, drawing on established life-cycle assessment principles to demonstrate the superior resource efficiency of aquatic systems (Pahlow et al., 2015).
2. To identify and profile fish species best suited for sustainable cultivation under crisis conditions. Selection criteria will prioritize physiological robustness, trophic level (favoring herbivorous and omnivorous species), and ability to utilize natural lake productivity, thus minimizing dependence on externally sourced formulated feeds (Yuan et al., 2020).
3. To develop a conceptual management model for lake aquaculture enterprises operating under a regime of minimal imported inputs. This model will integrate principles of polyculture, integrated multi-trophic aquaculture (IMTA), and nutrient recycling from agricultural by-products to create a semi-closed, self-sustaining system (Granada et al., 2016).
4. To propose concrete policy and practical recommendations for the integration of resilient lake aquaculture into national and regional food security frameworks. This includes guidance on site selection, germplasm conservation, and the development of decentralized, low-energy processing and preservation techniques to ensure the viability of the protein supply from water to table in a post-catastrophe landscape.

By fulfilling these objectives, this article will provide a foundational framework for policymakers, disaster planners, and agricultural scientists to leverage the untapped potential of inland water

bodies as strategic food reserves, thereby enhancing societal resilience in an increasingly uncertain world.

Vulnerabilities of Traditional Protein Production Systems in the Face of Catastrophe

A comprehensive assessment of food security strategies for catastrophic scenarios must begin with a critical examination of the incumbent systems. The global food supply's heavy reliance on terrestrial livestock, intensive crop agriculture, and marine capture fisheries creates a network of critical points of failure. These systems, optimized for efficiency under stable conditions, possess inherent vulnerabilities that become existential threats when global stability is disrupted (Hertel et al., 2010).

Livestock Production: A House of Cards

Industrial livestock production is characterized by a high degree of specialization and concentration, making it exceptionally fragile. Its dependence on external inputs is perhaps its greatest weakness. The sector is a major consumer of concentrated feed, primarily soy and maize, the production of which is often geographically disconnected from the livestock operations themselves (Godfray et al., 2010). A catastrophe disrupting global trade would sever this supply chain, leading to immediate and catastrophic herd collapses. As noted by Gilchrist et al. (2007), the modern livestock industry is "precariously dependent on a continuous supply of feed ingredients," with even minor disruptions causing significant economic and production losses.

Furthermore, the practice of Concentrated Animal Feeding Operations (CAFOs) creates perfect epidemiological storm conditions. The high density of genetically similar hosts facilitates the rapid transmission and mutation of pathogens. Events like the avian influenza (H5N1) and African swine fever outbreaks demonstrate the devastating speed with which diseases can wipe out populations in such settings (Capua & Alexander, 2004; Sánchez-Vizcaíno et al., 2013). In a catastrophe scenario, the parallel collapse of veterinary supply chains—including vaccines, antibiotics, and other pharmaceuticals—would remove the primary tools for disease control, turning CAFOs from protein production centers into mass graveyards.

The resource intensity of livestock farming further compounds its vulnerability. It is the largest anthropogenic user of freshwater, with a single kilogram of beef requiring between 15,000 to 20,000 liters of water, primarily for feed irrigation (Mekonnen & Hoekstra, 2012). This makes the sector highly susceptible to droughts and water scarcity, which are likely to be exacerbated by climate change. The energy footprint is equally staggering, encompassing inputs for feed production, transport, and operation of housing facilities (Pelling et al., 2015). In an energy-constrained post-catastrophe world, maintaining such energy-intensive operations would be untenable.

Crop Production: The Precarious Harvest

While crop production is the foundation of the global food system, providing calories for direct human consumption and feed for animals, it is acutely vulnerable to environmental and economic shocks. Its dependence on predictable weather patterns is a fundamental weakness. Climate change is already increasing the frequency and severity of extreme weather events, such as droughts and floods, which can lead to catastrophic crop failures (Lesk et al., 2016). Drought stress, in particular, has been shown to cause non-linear yield losses in major cereals like maize and wheat, threatening global staple supplies (Lobell et al., 2014).

Modern high-yield agriculture is also heavily reliant on synthetic fertilizers and pesticides, the production of which is energy-intensive and often dependent on global supply chains for raw materials like phosphate rock and natural gas. A systemic collapse would halt the production and distribution of these agrochemicals, leading to a rapid decline in yields. Studies have shown that removing synthetic nitrogen fertilization can lead to yield reductions of 20-50% in major cereal systems (Ten Berge et al., 2019). This "nitrogen shock" would be immediate and severe.

Moreover, the stability of monoculture systems is threatened by pests and diseases. The reduced genetic diversity in modern crop varieties creates a homogeneous landscape susceptible to emerging pathogens and pests. The disruption of supply chains for effective pesticides would leave crops defenseless, while the loss of international phytosanitary monitoring and response capabilities would allow new threats to spread unchecked (Sharma et al., 2017). The combined pressure of abiotic stresses (drought, flood) and biotic stresses (pests, diseases) in a disrupted world could lead to simultaneous breadbasket failures, a scenario the current system is ill-equipped to handle.

Marine Capture Fisheries: A Faltering Pillar

Marine fisheries, often perceived as a "wild" and resilient source of protein, are in fact highly susceptible to the cascading effects of a catastrophe. The first vulnerability lies in the direct impact of climate change on fish stocks. Ocean warming, acidification, and deoxygenation are altering marine ecosystems, causing shifts in species distribution, reducing maximum catch potential, and disrupting the productivity of entire food webs (Free et al., 2019). A single climate event, such as a marine heatwave, can have devastating and long-lasting impacts on local fisheries (Smale et al., 2019). In a catastrophe scenario, the baseline productivity of the oceans would already be under severe stress.

The logistical complexity of marine fishing presents a second critical vulnerability. The industry is dependent on functional infrastructure: fuel for vessels, functioning ports for landing catch, and reliable refrigeration for storage and transport. A breakdown in any one of these links—for instance, a fuel shortage—would instantly immobilize the vast majority of the global fishing fleet (Allison & Horemans, 2006). Unlike terrestrial food production, which can sometimes regress to lower-tech methods, open-ocean fishing is inherently technology- and energy-dependent.

Finally, the pre-catastrophe state of many fish stocks is already one of overexploitation. According to the FAO, the fraction of fish stocks within biologically sustainable levels has been steadily declining for decades (FAO, 2020). In a scenario where governance and international cooperation break down, the risk of a "race to fish" and a complete collapse of already stressed fisheries would be extremely high (Pinsky et al., 2018). Marine fisheries cannot be relied upon as a resilient protein source when they are already operating at their sustainable limits and are entirely dependent on a functioning global logistic and energy system.

In conclusion, the three pillars of conventional protein production—livestock, crops, and marine fisheries—are all built upon assumptions of stability, globalization, and resource abundance that would be invalidated by a major, prolonged catastrophe. Their interconnected vulnerabilities highlight the urgent need to develop and integrate more localized, resource-light, and resilient alternatives, such as lake-based aquaculture.

The Superiority of Lake Aquaculture as a Resilient Model

In contrast to the fragile pillars of conventional protein production, lake-based aquaculture presents a paradigm of remarkable resilience and efficiency. Its inherent biophysical and logistical characteristics align closely with the requirements for sustaining food security in a post-catastrophe world, offering a decentralized, resource-light, and ecologically buffered system for generating high-quality protein.

Energy and Resource Efficiency

The most compelling advantage of aquaculture lies in its superior biological efficiency, most clearly quantified by the Feed Conversion Ratio (FCR). The FCR measures the mass of feed required to produce a unit mass of animal protein. Due to their poikilothermic (cold-blooded) nature and buoyancy in water, fish expend far less metabolic energy on thermoregulation and structural support compared to terrestrial livestock (Boyd et al., 2020). This physiological efficiency translates into dramatically lower FCRs. While beef cattle require 6.0–10.0 kg of feed to produce 1.0 kg of live weight, and pigs require 2.7–5.0 kg, herbivorous and omnivorous fish like tilapia (*Oreochromis niloticus*) and carp (*Cyprinus carpio*) achieve FCRs as low as 1.2–1.5 under optimal conditions (Hua & Bureau, 2012). This stark differential means that for the same input of feed, lake aquaculture can produce several times more edible protein than the most common forms of livestock husbandry, a critical advantage when feed resources are scarce.

This fundamental efficiency is further amplified by the ability of lake aquaculture to utilize natural trophic pathways, a feature absent in terrestrial systems. Unlike land-based animal production that relies almost exclusively on provided feed, lake aquaculture, particularly in extensive and semi-intensive models, can leverage the natural productivity of the water body. Phytoplankton and zooplankton, fueled by solar energy and biogeochemical cycling of nutrients, form a natural, in-situ food source that can sustain a significant portion of the fish biomass (Yuan et al., 2020). This drastically reduces dependence on externally sourced, formulated feeds. The practice of

polyculture, which will be discussed in detail later, optimizes this by stocking complementary species that exploit different ecological niches within the lake. For instance, filter-feeding silver carp (*Hypophthalmichthys molitrix*) consume phytoplankton, while bottom-feeding common carp (*Cyprinus carpio*) utilize detritus and benthic organisms (Milstein, 2019). This creates a more complete and efficient utilization of the ecosystem's inherent productivity, moving towards a semi-closed system.

Resilience to Logistic Disruption

The static and localized nature of lake aquaculture is a critical strategic advantage in a scenario characterized by fragmented supply chains and energy scarcity. A lake is a fixed-production asset that does not require constant fuel input to remain productive. This stands in stark contrast to marine fisheries, which are entirely dependent on diesel for vessel operations, and terrestrial supply chains that require constant fuel for the transport of feed, animals, and products (Béné et al., 2016). Once a lake is stocked, the production process is geographically contained and largely self-sustaining from an energy perspective, relying on natural primary production.




Figure 1: Feed Conversion Ratio (FCR) Comparison: Livestock vs. Aquaculture Species.

This localization also dramatically simplifies post-harvest logistics and reduces vulnerability. The requirement for complex, energy-intensive cold chains is significantly reduced. Without reliable

refrigeration or freezing, marine catches can spoil within hours. In contrast, fish in a lake represent a "live storage" system—a dynamic protein bank that can be harvested on demand, as needed, directly by local communities (Brugère et al., 2019). This eliminates the need for the extensive, energy-dependent infrastructure for processing, freezing, and long-distance transport that characterizes modern food systems. The lake itself acts as a natural, zero-energy bioreactor, holding tank, and refrigeration system, ensuring protein availability without external energy inputs.

Ecological Stability and Buffering Capacity

The aquatic environment of a lake provides a significant natural buffer against environmental fluctuations, offering a more stable habitat for cultivation than terrestrial settings. Water has a high specific heat capacity, meaning it heats up and cools down much more slowly than air. This thermal inertia dampens daily and seasonal temperature swings, protecting the cultivated biomass from sudden thermal stress that can devastate crops and terrestrial animals (Ficke et al., 2007). While climate change does impact lakes (e.g., through warming and stratification), the rate of temperature change within the water column is generally slower and more predictable than atmospheric changes, allowing for more gradual biological adaptation and management interventions.

This buffering capacity extends to other abiotic factors. The water body provides physical protection from direct atmospheric phenomena like wind, hail, and frost, which are major causes of catastrophic crop loss (Lesk et al., 2016). Furthermore, the substantial volume of water in a lake dilutes metabolic wastes and provides a stable osmotic environment, reducing the physiological stress on the animals compared to the often crowded and managed environments of terrestrial farms (Boyd & Tucker, 2014). This inherent stability makes lake-based systems less prone to a total collapse from a single acute weather event, a common vulnerability in modern agriculture.

Minimal Competition for Land Resources

In a catastrophe scenario where every hectare of arable land becomes critical for direct caloric production from staple crops, the ability to produce protein without competing for this precious resource is a decisive benefit. Lake aquaculture operates in a distinct ecological domain—the lentic water body—that is generally unsuitable for conventional agriculture (Bogard et al., 2017). By utilizing these existing water bodies for food production, societies can effectively expand their productive footprint without the need to convert forests or grasslands into farmland, a process that is both ecologically damaging and irreversible in the short term.

This represents a form of sustainable intensification that avoids the high environmental costs of land conversion. Moreover, lake aquaculture can be integrated synergistically with land-based agriculture in a circular economy model. Agricultural by-products and effluents, rich in organic matter and nutrients, can be responsibly managed to fertilize lakes, thereby enhancing natural productivity (a practice that must be carefully managed to prevent eutrophication) (Nhan et al., 2019). This creates a synergistic relationship where "waste" from one system becomes a

valuable resource for another, enhancing overall system resilience, reducing pollution, and minimizing the need for external synthetic inputs.

In summary, lake aquaculture's combination of superior biological efficiency, logistical simplicity, environmental buffering, and non-competitive land use positions it as a uniquely robust model for protein production. It is a system designed for persistence and autonomy rather than peak efficiency under globalized, ideal conditions, making it an indispensable component of a strategic framework aimed at weathering prolonged systemic disruptions.

Species Selection and Management Models for a Post-Catastrophe Scenario

The theoretical resilience of lake aquaculture must be operationalized through pragmatic, crisis-aware strategies for species selection and system management. In a post-catastrophe scenario, the overarching objective shifts from economic maximization to robust survivability and continuous protein production under conditions of severe resource scarcity and infrastructural collapse. This necessitates a fundamental re-evaluation of both the biological agents and the operational frameworks of aquaculture.

The Principle of "Hardiness and Autonomy" in Species Selection

The selection of fish species is the primary determinant of system resilience. The ideal species for a catastrophic food security strategy are those possessing a suite of traits that confer environmental tolerance, dietary flexibility, and disease resistance, thereby minimizing their dependence on external inputs and technological interventions (Béné et al., 2016). Key selection criteria must include:

- **Environmental Tolerance:** High resilience to fluctuations in temperature, dissolved oxygen, pH, and salinity. This is crucial as monitoring and water quality management capabilities may be severely degraded.
- **Disease Resistance:** Innate immunological robustness to common pathogens, reducing catastrophic losses in the absence of consistent access to vaccines, antibiotics, and other veterinary pharmaceuticals.
- **Trophic Level and Dietary Flexibility:** A strong preference for herbivorous and omnivorous species that can utilize a wide range of natural food sources (phytoplankton, zooplankton, detritus, aquatic vegetation) and alternative, low-quality feeds (agricultural by-products, food waste).
- **Reproductive Capacity:** Species that can be reliably spawned in simple, low-tech hatchery conditions or that exhibit natural recruitment within lake environments, ensuring a self-sustaining population without reliance on complex breeding facilities.

Based on these stringent criteria, a select group of species emerges as particularly suitable for post-catastrophe aquaculture:

- Common Carp (*Cyprinus carpio*): A quintessential resilient species, the common carp is a bottom-feeding omnivore (benthivore) renowned for its exceptional tolerance of poor water quality, including low oxygen levels (Rakus et al., 2017). It efficiently recycles nutrients by consuming detritus, benthic invertebrates, and other organic matter, playing a key role in ecosystem functioning within the lake (Woynarovich et al., 2010).
- Silver Carp (*Hypophthalmichthys molitrix*) and Bighead Carp (*Hypophthalmichthys nobilis*): These filter-feeding species are unparalleled in their ability to directly harvest primary and secondary production. Silver carp primarily consume phytoplankton, while bighead carp target zooplankton, allowing them to convert solar energy, captured in the base of the aquatic food web, directly into fish biomass without the need for any processed feed (Cremen et al., 2007; Liu et al., 2018).
- Grass Carp (*Ctenopharyngodon idella*): As a dedicated herbivore, the grass carp is highly effective at controlling aquatic macrophytes and can be sustained on terrestrial grasses and agricultural waste products, providing a direct link between land-based plant biomass and aquatic protein (Shireman & Smith, 2018).
- Nile Tilapia (*Oreochromis niloticus*): This tropical species is a cornerstone of global aquaculture due to its fast growth rate, high fecundity, and robust disease resistance. As an omnivore with a strong preference for phytoplankton and periphyton, it efficiently utilizes natural productivity (El-Sayed, 2019). Its primary limitation is a sensitivity to cold water, confining its use to temperate regions with warm summers or to tropical and subtropical climates.
- African Catfish (*Clarias gariepinus*) and Channel Catfish (*Ictalurus punctatus*): These species are exceptionally tolerant of hypoxic conditions and high stocking densities. Their carnivorous-to-omnivorous feeding habits allow them to consume a wide range of agricultural by-products, food waste, and even offal, making them excellent recyclers of organic waste streams in a collapsed economy (Hossain et al., 2021).

Management Models: Extensivity, Semi-Intensivity, and Polyculture

The management philosophy for catastrophic resilience must pivot decisively from high-input intensity to low-input optimization. The goal is to maximize output per unit of external input, rather than pursuing absolute maximum yield.

Schematic of a Polyculture Lake Ecosystem




Figure 2: Schematic of a Polyculture Lake Ecosystem.

- Extensive and Semi-Intensive Models: These models form the operational backbone of a resilient system. Extensive aquaculture relies almost entirely on the natural productivity of the lake to support fish growth, with stocking densities carefully calibrated to the ecosystem's inherent carrying capacity. Semi-intensive systems provide a pragmatic middle ground, supplementing natural food sources with minimal, locally available inputs such as cereal grains (wheat, maize), agricultural by-products (rice bran, oilcakes), and household food waste (Edwards, 2015). This approach drastically reduces or eliminates dependence on manufactured, protein-rich pelleted feeds, the industrial production of which would likely cease in a major catastrophe. The focus is on leveraging and enhancing the lake's natural trophic status, a fundamental shift from intensive systems that rely on constant technological and chemical intervention to overcome ecological limits.
- Polyculture: This is the practice of cultivating multiple, ecologically complementary fish species in the same water body, and it is a powerful strategy for optimizing the use of all available trophic niches. By stocking species that occupy different feeding guilds, polyculture mimics natural ecosystem structure and leads to a more complete utilization of the aquatic food web, thereby increasing total productivity without increasing external inputs (Milstein, 2019). A classic and highly effective polyculture combination for temperate and subtropical regions includes:

- Silver Carp: Harvests the phytoplankton niche.
- Bighead Carp: Harvests the zooplankton niche.
- Grass Carp: Controls aquatic weeds and consumes provided vegetation.
- Common Carp: Recycles nutrients from the bottom sediments.
This synergistic combination results in a more stable and productive ecosystem than any single species could achieve alone, as it minimizes the waste of natural food resources and creates a more balanced aquatic environment, reducing the risk of algal blooms and oxygen depletion (Liu et al., 2018).

Creating a Closed-Loop System: Integration with Agriculture

To achieve a high degree of autonomy and resilience, lake aquaculture must be integrated into a broader agro-ecological system. This involves creating deliberate nutrient loops that connect fish farming with terrestrial crop and livestock production, effectively mimicking and harnessing natural biogeochemical cycles.

The most direct form of this integration is the controlled use of organic fertilizers, such as livestock manure (from poultry, pigs, or ruminants), to fertilize the lake. When applied in judicious quantities, the nutrients in manure (primarily nitrogen and phosphorus) stimulate the growth of phytoplankton, which forms the base of the food web for filter-feeding and omnivorous fish (Nhan et al., 2019). This practice, a cornerstone of Integrated Agriculture-Aquaculture (IAA), effectively transforms animal waste into high-quality fish protein.

This creates a virtuous, closed-loop cycle: agricultural by-products feed the livestock, livestock manure fertilizes the lake, the lake produces fish for human consumption, and the nutrient-rich sediments that accumulate on the lake bottom can later be excavated and used as a potent fertilizer for crops, completing the loop (Prein, 2020). Such integrated systems are inherently more resilient because they diversify food outputs (grain, vegetables, meat, fish) and reduce dependency on external inputs to near zero. They create self-sustaining, localized food production nodes that can persist independently in a post-catastrophe landscape. The key to success lies in careful, knowledge-based management to prevent over-fertilization and cultural eutrophication, a challenge that requires skill and observation rather than expensive technology.

Overcoming Potential Risks and Challenges

While lake aquaculture presents a compelling model for resilience, its strategic implementation requires the proactive mitigation of inherent risks. A post-catastrophe scenario exacerbates vulnerabilities related to disease, genetic resource availability, ecological degradation, and food preservation. Addressing these challenges through pre-emptive planning and robust, low-tech protocols is essential for ensuring the long-term viability of these aquatic food systems.

Epidemiological Control: Biosecurity and Prevention

In a landscape where access to veterinary pharmaceuticals, vaccines, and professional diagnostics is severely limited or nonexistent, the paradigm for disease management must shift from treatment to prevention. The cornerstone of this approach is the implementation of stringent biosecurity protocols designed to minimize pathogen introduction and spread (Stuart et al., 2023). Key measures include the quarantine of new stock before introduction into lakes, restrictions on the movement of equipment and personnel between different water bodies, and the control of potential wildlife disease vectors where feasible (Murray & Peeler, 2015).

Beyond biosecurity, a fundamental strategy is the cultivation of genetic resistance. Selective breeding programs focused on enhancing disease resilience, rather than solely on growth performance, are critical for long-term sustainability. For example, breeding for resistance to specific pathogens like Koi Herpesvirus in common carp (*Cyprinus carpio*) has shown promising results and should be a priority for genetic stock development (Rakus et al., 2017). Furthermore, maintaining lower stocking densities, a core tenet of semi-intensive polyculture, inherently reduces stress and the probability of explosive disease outbreaks, creating a more stable and resilient epidemiological environment (Buller, 2004).

Germplasm Security and Broodstock Networks

The continuity of lake aquaculture is entirely dependent on a reliable supply of juvenile fish (fingerlings). The globalization of aquaculture has led to a situation where many regions depend on international trade for their seed stock, a supply chain that would be instantly severed by a catastrophe (Brugère et al., 2019). To mitigate this, it is imperative to establish decentralized, national or regional networks of broodstock banks and hatcheries.

These facilities would maintain genetically diverse breeding populations (broodstock) of the key resilient species discussed in Section 4. The network should be geographically distributed to safeguard against localized disasters and managed with a focus on maintaining genetic diversity to prevent inbreeding depression and preserve adaptive potential (López et al., 2021). These hatcheries would operate on a low-tech, resilient model, capable of producing fingerlings using simple hormone induction techniques for spawning and relying on natural plankton as initial feed for larvae (Mylonas et al., 2010). Securing this most fundamental input—the seed—is a non-negotiable prerequisite for a self-sufficient food security strategy.

Monitoring and Preventing Ecological Degradation

Aquaculture, if mismanaged, can become a source of environmental degradation, primarily through eutrophication caused by the accumulation of nutrient-rich wastes. In a post-catastrophe world, without the energy for mechanical aeration or the chemicals for water treatment, prevention through ecological management is the only viable strategy. The primary lever for control is the strict regulation of stocking density and nutrient loading to remain within the lake's assimilative capacity (Boyd & Tucker, 2014).

The polyculture models previously described are a key mitigation tool, as the diverse species assemblage facilitates more efficient nutrient recycling within the ecosystem (Liu et al., 2018). Additionally, the integration with agriculture must be carefully calibrated; while manure application can stimulate food production, over-application will lead to algal blooms, oxygen depletion, and system collapse. Low-tech monitoring methods, such as Secchi disks for water transparency and observation of indicator species, can provide adequate environmental assessment without sophisticated laboratories (Wang et al., 2020). The application of permaculture principles, such as creating wetland filters at inflow/outflow points to process nutrients, can further enhance the ecological stability and sustainability of the system.

Low-Energy Processing and Preservation

The reliance on energy-intensive freezing and refrigeration for food preservation is a critical vulnerability in a catastrophic scenario. To ensure the nutritional benefits of the fish harvest are not lost, it is essential to revitalize and optimize traditional, low-energy preservation techniques. Salting, drying, and smoking are highly effective methods that have sustained human populations for millennia (Ghaly et al., 2010).

Solar drying, in particular, represents a perfect synergy of resilience and practicality, using an abundant energy source to remove moisture and inhibit microbial growth (Doe, 2022). Salting (both dry-salting and brining) and fermentation (as used in traditional fish sauces and pastes) are also highly effective and require minimal infrastructure (Sampels, 2015). Investing in the research and development of these techniques to improve their efficiency, safety, and nutritional retention is a crucial component of a comprehensive food security strategy. By decentralizing preservation to the household or community level using these methods, the system becomes robust against the loss of large-scale, centralized cold chains.

A strategic focus on overcoming these four critical challenges—through biosecurity, genetic autonomy, ecological balance, and low-tech preservation—will transform lake aquaculture from a theoretically resilient concept into a practical, fail-safe pillar of long-term catastrophic food security.

A Strategic Roadmap for Integrating Lake Aquaculture into National Security Frameworks

The compelling case for lake aquaculture as a catastrophic food security solution necessitates a concrete and multi-faceted implementation strategy. Translating this theoretical resilience into practical readiness requires coordinated action across legislative, scientific, economic, and educational domains. This section outlines a strategic roadmap to embed resilient lake aquaculture within the core of national food security policy.

Legislative and Policy Foundation: Mainstreaming Resilience

The foundational step is the formal recognition of "Resilience Aquaculture" within national strategic documents. Specifically, governments should amend National Food Security Doctrines and Disaster Preparedness Plans to explicitly include lake-based aquaculture as a critical component of protein security during long-term emergencies (Brugère et al., 2019). This legislative action would mandate the identification, mapping, and zoning of suitable lakes and reservoirs for this purpose within regional spatial planning frameworks (Beveridge et al., 2018). Such zoning would prioritize water bodies based on ecological carrying capacity, proximity to population centers, and low vulnerability to pollution or extreme climatic events. This process would also involve establishing clear legal frameworks for water use rights and access protocols during a state of emergency, preventing conflict and ensuring that these aquatic resources can be swiftly mobilized for food production when needed (Bennett et al., 2021).




Figure 3: Strategic Roadmap for Integrating Resilient Lake Aquaculture into National Security.

Scientific and Technical Research: Building Autonomy

Strategic government investment in Research & Development (R&D) is crucial for optimizing system resilience and overcoming the technical challenges outlined previously. Key research priorities should include:

- **Genetic Breeding Programs:** Funding long-term selective breeding programs focused not on maximal growth under ideal conditions, but on enhancing traits critical for survival in low-input systems. This includes genetic selection for disease resistance (e.g., to

pathogens like *Aeromonas hydrophila* and Koi Herpesvirus), tolerance to fluctuating temperature and dissolved oxygen, and feed conversion efficiency on agricultural by-products (Yáñez et al., 2021; López et al., 2021).

- **Low-Energy Technology Development:** Supporting the innovation of autonomous systems powered by renewable energy. This includes R&D into solar-powered aeration systems to mitigate occasional oxygen crashes, and automated feeding systems that dispense locally sourced, non-formulated feeds, thereby reducing labor and optimizing resource use (Boyd et al., 2020).
- **Ecological Modeling:** Developing sophisticated yet user-friendly models to determine the optimal polyculture species ratios and stocking densities for different lake types and climatic zones, ensuring sustainable yields without ecological degradation (Liu et al., 2018).

Economic Incentives: Catalyzing Private and Cooperative Action

To build a decentralized and robust network of production units, state-level economic incentives are essential. These should be designed to encourage the establishment and maintenance of lake farms that adhere to pre-approved "crisis-resilient" protocols (e.g., polyculture of designated species, use of extensive/semi-intensive methods, maintenance of broodstock). Effective mechanisms could include:

- **Targeted Subsidies:** Direct grants or cost-sharing for the initial setup of resilient aquaculture systems, including the purchase of native fingerlings from certified broodstock centers and the installation of renewable energy infrastructure.
- **Tax Incentives:** Offering property tax reductions or income tax credits for lake aquaculture operations that are certified as following resilient protocols, making them financially attractive even in non-crisis times (Béné et al., 2016).
- **Support for Cooperatives:** Providing preferential loans and technical assistance to foster the creation of local fishing and aquaculture cooperatives. This model can effectively manage community lake resources, share costs and knowledge, and ensure equitable distribution of the protein harvest during a crisis (Brugère & Selbie, 2021).

Educational and Capacity Building: Cultivating Human Capital

The success of this strategy is ultimately dependent on a skilled workforce trained in the principles of resilient, low-input aquaculture. This requires a revitalization of agricultural and veterinary education to include specialized tracks in extensive lake management. Key initiatives should include:

- **Curriculum Development:** Integrating modules on crisis-resilient polyculture, low-tech health management (biosecurity, disease identification), traditional preservation

techniques, and integrated agro-aquaculture systems into university programs for ichthyologists, veterinarians, and aquaculture engineers (Stuart et al., 2023).

- Vocational Training and Extension Services: Establishing widespread vocational training and farmer field schools to disseminate these practices to existing fish farmers and new entrants. Government extension services should be equipped with the knowledge to support these decentralized hubs of production (Brugère et al., 2019).
- Knowledge Preservation: Systematically documenting and studying indigenous and local knowledge related to freshwater fish husbandry and preservation, ensuring that valuable empirical wisdom is not lost and can be integrated with modern scientific approaches (Bennett et al., 2021).

In conclusion, the integration of lake aquaculture into national security is not a single action but a sustained, multi-level endeavor. By enacting supportive legislation, driving targeted R&D, providing smart economic incentives, and investing in specialized education, nations can proactively build a distributed, resilient, and self-sufficient protein reserve. This strategic investment transforms a theoretical safeguard into a tangible, operational asset, fundamentally enhancing a nation's capacity to withstand and recover from catastrophic disruptions to its food supply.

Discussion

This analysis posits that lake-based aquaculture represents a critically undervalued component of strategic planning for food and protein security in the face of systemic catastrophes. The vulnerabilities inherent in conventional terrestrial livestock, crop production, and marine fisheries are profound and interlinked, rendering them highly susceptible to collapse under the strain of prolonged disruptions to global trade, energy supplies, and climatic stability (Godfray et al., 2010; Béné et al., 2016). In this context, the comparative advantages of lacustrine aquaculture—its superior resource efficiency, logistical simplicity, ecological buffering, and non-competitive land use—elevate it from a mere alternative to a necessary strategic reserve.

The discussion herein has moved beyond the established narrative of aquaculture as a tool for sustainable development under "business-as-usual" conditions (FAO, 2020) and reframed it as a fail-safe mechanism for societal resilience. The core argument is that the very characteristics which often relegate extensive and semi-intensive lake aquaculture to a secondary role in the modern globalized economy—lower absolute yields, slower growth rates compared to intensive systems—become its greatest strengths in a de-globalized, resource-scarce environment. Its low external input requirements and dependence on natural trophic pathways are precisely what confer its robustness (Edwards, 2015). This represents a fundamental paradigm shift: prioritizing system persistence over peak productivity.

Synthesis of Advantages and the Resilience Framework

The resilience of the proposed model is not derived from a single factor but from the synergistic integration of its components. The biological efficiency of herbivorous and omnivorous fish species, with their low Feed Conversion Ratios (FCRs), provides the foundational advantage (Boyd et al., 2020). This is amplified by the polyculture management strategy, which optimizes the use of the entire aquatic food web, from phytoplankton to benthic detritus, thereby maximizing the conversion of solar energy and natural nutrients into human-edible protein without requiring manufactured inputs (Milstein, 2019; Liu et al., 2018). This creates a production system that is largely decoupled from the fossil fuel and industrial chemical inputs upon which modern agriculture is precariously balanced.

Furthermore, the "live storage" capacity of lakes is a feature unmatched by any terrestrial system. A grain silo or a refrigerated warehouse represents a static, depleting reserve that requires energy to maintain. A stocked lake, however, is a dynamic, self-sustaining protein bank that maintains and even grows its asset (the fish biomass) using ambient solar energy captured through natural ecological processes (Brugère et al., 2019). This eliminates the need for the complex, energy-intensive logistics of harvesting, processing, and refrigerated transport that characterize both marine fisheries and terrestrial meat supply chains.

Confronting the Challenges: A Pre-emptive Strategy

Acknowledging the potential risks is essential for credible strategic planning. The proposed mitigation strategies are deliberately designed to be knowledge-intensive rather than technology- or resource-intensive, making them feasible in a post-catastrophe context. The emphasis on biosecurity and genetic selection for disease resistance (Rakus et al., 2017; Stuart et al., 2023) is a proactive approach that circumvents the need for a continuous supply of veterinary pharmaceuticals. Similarly, the establishment of decentralized broodstock networks (López et al., 2021) is a one-time strategic investment that ensures long-term autonomy in seed production, breaking a critical dependency on international supply chains.

The ecological challenge of eutrophication is addressed not with high-tech solutions, but through the prudent application of ecological principles: maintaining stocking densities within the lake's carrying capacity and using polyculture to enhance nutrient assimilation (Boyd & Tucker, 2014). This approach requires skilled management and monitoring, underscoring the importance of education and knowledge dissemination as a core element of preparedness. Likewise, the revival of low-energy preservation techniques like solar drying and salting (Ghaly et al., 2010; Doe, 2022) leverages timeless food preservation wisdom, making the system resilient to the loss of modern refrigeration.

Limitations and Research Needs

This conceptual framework has limitations that warrant further investigation. First, the model is most readily applicable to regions with abundant freshwater lakes; arid regions would need to adapt the principles to reservoir-based or water-conserving integrated systems. Second, the

potential for social conflict over access to and management of lake resources in a crisis scenario is a significant, though not insurmountable, challenge that requires parallel development of governance models (Brugère et al., 2019).

Significant research gaps remain. Future work should prioritize:

1. **Quantitative Modeling:** Developing robust models to calculate the precise carrying capacity of specific lake types for polyculture systems under low-input regimes.
2. **Genetic Resource Development:** Establishing formal breeding programs focused explicitly on enhancing the stress tolerance and disease resilience of key species like carp, tilapia, and catfish for resilient production.
3. **Optimization of Traditional Techniques:** Scientific validation and optimization of low-energy fish processing methods to maximize nutritional retention, safety, and shelf-life.
4. **Socio-economic Integration:** Research into effective governance structures and economic incentives for establishing and maintaining these systems within national food security strategies before a catastrophe occurs.

The evidence assembled strongly supports the thesis that lake aquaculture warrants inclusion as a strategic pillar in national and regional catastrophic food security planning. Its unique combination of biological efficiency, logistical independence, and ecological stability offers a pathway to secure high-quality protein in scenarios that would cripple conventional food systems. Proactive investment in the development of resilient species stocks, the dissemination of knowledge on extensive polyculture management, and the integration of these systems into local agro-ecological cycles is not merely an academic exercise. It is a pragmatic, forward-looking strategy for enhancing societal resilience against an increasingly uncertain future. Ignoring the strategic potential of inland water bodies as food reservoirs would be to overlook a critical tool for survival and recovery in the wake of a global catastrophe.

Conclusion and Implications

This analysis has systematically argued that lake-based aquaculture must be reconceptualized from a niche agricultural activity to a strategically indispensable component of 21st-century food security architecture. The escalating frequency of climate-related disasters, the persistent threat of geopolitical instability, and the ever-present risk of global pandemics reveal the profound fragility of our centralized, input-intensive food systems (Godfray et al., 2010; Béné et al., 2016; Laborde et al., 2020). In this context of systemic vulnerability, the innate resilience, resource efficiency, and functional capacity of lacustrine aquaculture in degraded infrastructural environments render it not merely an alternative, but a critical insurance policy against a wide spectrum of catastrophes.

The evidence presented demonstrates a clear comparative advantage. The superior Feed Conversion Ratio (FCR) of herbivorous and omnivorous fish species establishes a foundation of

biological efficiency that terrestrial livestock cannot match, producing more protein with less resource input (Boyd et al., 2020; Hua & Bureau, 2012). This efficiency is magnified through management models like polyculture, which optimize the use of natural trophic pathways, transforming lakes into semi-autonomous protein production systems that minimize dependence on external feed, energy, and complex logistics (Milstein, 2019; Liu et al., 2018). The "live storage" capacity of a stocked lake provides a dynamic, self-sustaining food reserve that is inherently more resilient than static grain silos or energy-dependent cold chains (Brugère et al., 2019). Furthermore, by utilizing lentic water bodies, this strategy produces high-quality protein without competing for scarce arable land, a decisive benefit in a crisis where every hectare of farmland is critical for caloric production (Bogard et al., 2017).

The strategic implementation of this model hinges on proactive measures to mitigate inherent risks. This includes the development of robust biosecurity protocols and the selective breeding of disease-resistant genetic lines to overcome veterinary supply chain collapse (Stuart et al., 2023; Rakus et al., 2017). It necessitates the creation of decentralized broodstock networks and hatcheries to ensure a self-sufficient supply of fingerlings, securing the most fundamental input for the system (López et al., 2021). Ecological sustainability must be maintained through careful stock management and the adoption of permaculture principles to prevent eutrophication, while food preservation must be guaranteed through the revitalization of low-energy techniques like solar drying and salting (Ghaly et al., 2010; Doe, 2022).

Therefore, the conclusions of this article are unequivocal. First, lake aquaculture possesses a unique combination of attributes—biological efficiency, logistical simplicity, and ecological buffering—that make it uniquely suited to provide a reliable source of protein when conventional systems fail. Second, realizing this potential requires a paradigm shift in how we value inland water bodies; they must be viewed not only as recreational or ecological assets but as strategic food reservoirs integral to national security. Third, the challenges of disease, genetics, ecology, and preservation are significant but surmountable through pre-emptive, knowledge-based strategies that do not rely on the continued functioning of a globalized technosphere.

The imperative for action is clear. Investment in the development of this sector—through research on resilient polycultures, the establishment of genetic banks, and the training of a new generation of aquaculturists in low-input methods—is not an expenditure, but a vital investment in national food sovereignty and long-term survivability (Brugère et al., 2019; Nhan et al., 2019). To ignore the strategic potential of our lacustrine resources is to leave a critical vulnerability unaddressed. By harnessing the innate productivity of lakes through the prudent, scientifically-grounded framework outlined here, we can build a more resilient food future, ensuring that even in the darkest of times, a fundamental source of nourishment and sustenance remains within reach.

References:

Allison, E. H., & Horemans, B. (2006). Putting the principles of the Sustainable Livelihoods Approach into fisheries development policy and practice. *Marine Policy*, 30(6), 757–766. <https://doi.org/10.1016/j.marpol.2006.02.001>

- Béné, C., Arthur, R., Norbury, H., Allison, E. H., Beveridge, M., Bush, S., Campling, L., Leschen, W., Little, D., Squires, D., Thilsted, S. H., Troell, M., & Williams, M. (2016). Contribution of fisheries and aquaculture to food security and poverty reduction: Assessing the current evidence. *World Development*, 79, 177–196. <https://doi.org/10.1016/j.worlddev.2015.11.007>
- Bogard, J. R., Farook, S., Marks, G. C., Waid, J., Belton, B., Ali, M., ... & Thilsted, S. H. (2017). Higher fish but lower micronutrient intakes: Temporal changes in fish consumption from capture fisheries and aquaculture in Bangladesh. *PloS One*, 12(4), e0175098. <https://doi.org/10.1371/journal.pone.0175098>
- Boyd, C. E., & Tucker, C. S. (2014). *Handbook for Aquaculture Water Quality*. Pond Aquaculture Research and Development Foundation.
- Boyd, C. E., D'Abramo, L. R., Glencross, B. D., Huyben, D. C., Juarez, L. M., Lockwood, G. S., ... & Valenti, W. C. (2020). Achieving sustainable aquaculture: Historical and current perspectives and future needs and challenges. *Journal of the World Aquaculture Society*, 51(3), 578-633. <https://doi.org/10.1111/jwas.12714>
- Brugère, C., Troell, M., & Beveridge, M. (2019). Aquaculture and the future of food. *Nature Food*, 1(6), 330-331. <https://doi.org/10.1038/s43016-020-0121-y>
- Buller, N. B. (2004). *Bacteria from fish and other aquatic animals: A practical identification manual*. CABI Publishing. <https://doi.org/10.1079/9780851997384.0000>
- Capua, I., & Alexander, D. J. (2004). Avian influenza: recent developments. *Avian Pathology*, 33(4), 393–404. <https://doi.org/10.1080/03079450410001724085>
- Cremen, M. C. M., Martinez-Goss, M. R., Corre, V. L., & Azanza, R. V. (2007). Phytoplankton and zooplankton dynamics in polyculture ponds. *Aquaculture Research*, 38(9), 945-955. <https://doi.org/10.1111/j.1365-2109.2007.01757.x>
- Doe, P. E. (2022). Fish drying. In *Seafood Processing: Technology, Quality and Safety* (pp. 61-80). Wiley-Blackwell. <https://doi.org/10.1002/9781118346174.ch4>
- Edwards, P. (2015). Aquaculture environment interactions: Past, present and likely future trends. *Aquaculture*, 447, 2-14. <https://doi.org/10.1016/j.aquaculture.2015.02.001>
- El-Sayed, A. F. M. (2019). *Tilapia culture*. Academic Press. <https://doi.org/10.1016/B978-0-12-816509-6.00001-5>
- FAO. (2020). *The State of World Fisheries and Aquaculture 2020. Sustainability in action*. Food and Agriculture Organization of the United Nations. <https://doi.org/10.4060/ca9229en>
- FAO. (2021). *The State of Food Security and Nutrition in the World 2021. Transforming food systems for food security, improved nutrition and affordable healthy diets for all*. Food and Agriculture Organization of the United Nations. <https://doi.org/10.4060/cb4474en>
- Ficke, A. D., Myrick, C. A., & Hansen, L. J. (2007). Potential impacts of global climate change on freshwater fisheries. *Reviews in Fish Biology and Fisheries*, 17(4), 581-613. <https://doi.org/10.1007/s11160-007-9059-5>
- Free, C. M., Thorson, J. T., Pinsky, M. L., Oken, K. L., Wiedenmann, J., & Jensen, O. P. (2019). Impacts of historical warming on marine fisheries production. *Science*, 363(6430), 979–983. <https://doi.org/10.1126/science.aau1758>
- Ghaly, A. E., Dave, D., Budge, S., & Brooks, M. S. (2010). Fish spoilage mechanisms and preservation techniques: A review. *American Journal of Applied Sciences*, 7(7), 859-877. <https://doi.org/10.3844/ajassp.2010.859.877>
- Gilchrist, M. J., Greko, C., Wallinga, D. B., Beran, G. W., Riley, D. G., & Thorne, P. S. (2007). The potential role of concentrated animal feeding operations in infectious disease epidemics and antibiotic resistance. *Environmental Health Perspectives*, 115(2), 313–316. <https://doi.org/10.1289/ehp.8837>

- Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., Pretty, J., Robinson, S., Thomas, S. M., & Toulmin, C. (2010). Food security: The challenge of feeding 9 billion people. *Science*, 327(5967), 812–818. <https://doi.org/10.1126/science.1185383>
- Granada, L., Sousa, N., Lopes, S., & Lemos, M. F. L. (2016). Is integrated multitrophic aquaculture the solution to the sectors' major challenges? – a review. *Reviews in Aquaculture*, 8(3), 283–300. <https://doi.org/10.1111/raq.12093>
- Herrero, M., Havlík, P., Valin, H., Notenbaert, A., Rufino, M. C., Thornton, P. K., Blümmel, M., Weiss, F., Grace, D., & Obersteiner, M. (2013). Biomass use, production, feed efficiencies, and greenhouse gas emissions from global livestock systems. *Proceedings of the National Academy of Sciences*, 110(52), 20888–20893. <https://doi.org/10.1073/pnas.1308149110>
- Hertel, T. W., Burke, M. B., & Lobell, D. B. (2010). The poverty implications of climate-induced crop yield changes by 2030. *Global Environmental Change*, 20(4), 577–585. <https://doi.org/10.1016/j.gloenvcha.2010.07.001>
- Hossain, M. A. R., Das, I., Genevier, L., Hazra, S., & Rahman, M. (2021). Biology and aquaculture of African catfish (*Clarias gariepinus*) and Asian catfish (*Pangasianodon hypophthalmus*). *Reviews in Aquaculture*, 13(4), 1973–2003. <https://doi.org/10.1111/raq.12554>
- Hua, K., & Bureau, D. P. (2012). Exploring the possibility of quantifying the effects of using alternative ingredients in fish feeds on the environmental performance of aquaculture systems. *Aquaculture*, 356, 292–299. <https://doi.org/10.1016/j.aquaculture.2012.04.045>
- IPCC. (2022). *Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- Jaba, T. (2022). Dasatinib and quercetin: short-term simultaneous administration yields senolytic effect in humans. *Issues and Developments in Medicine and Medical Research Vol. 2*, 22–31.
- Jones, B. A., Grace, D., Kock, R., Alonso, S., Rushton, J., Said, M. Y., McKeever, D., Mutua, F., Young, J., McDermott, J., & Pfeiffer, D. U. (2013). Zoonosis emergence linked to agricultural intensification and environmental change. *Proceedings of the National Academy of Sciences*, 110(21), 8399–8404. <https://doi.org/10.1073/pnas.1208059110>
- Laborde, D., Martin, W., Swinnen, J., & Vos, R. (2020). COVID-19 risks to global food security. *Science*, 369(6503), 500–502. <https://doi.org/10.1126/science.abc4765>
- Lesk, C., Rowhani, P., & Ramankutty, N. (2016). Influence of extreme weather disasters on global crop production. *Nature*, 529(7584), 84–87. <https://doi.org/10.1038/nature16467>
- Liu, X., Zhang, G., Sun, G., Wu, Y., & Chen, Y. (2018). The synergistic effects of polyculture on the ecosystem services of aquaculture ponds. *Aquaculture Research*, 49(9), 3044–3053. <https://doi.org/10.1111/are.13768>
- Lobell, D. B., Roberts, M. J., Schlenker, W., Braun, N., Little, B. B., Rejesus, R. M., & Hammer, G. L. (2014). Greater sensitivity to drought accompanies maize yield increase in the U.S. Midwest. *Science*, 344(6183), 516–519. <https://doi.org/10.1126/science.1251423>
- López, M. E., Neira, R., & Yáñez, J. M. (2021). Applications of genome-wide selection in aquaculture breeding programs. *Aquaculture Reports*, 21, 100869. <https://doi.org/10.1016/j.aqrep.2021.100869>
- Mekonnen, M. M., & Hoekstra, A. Y. (2012). A global assessment of the water footprint of farm animal products. *Ecosystems*, 15(3), 401–415. <https://doi.org/10.1007/s10021-011-9517-8>
- Milstein, A. (2019). Polyculture in aquaculture. In *Animal Agriculture* (pp. 343–355). Academic Press. <https://doi.org/10.1016/B978-0-12-817052-6.00020-7>

- Murray, A. G., & Peeler, E. J. (2015). A framework for understanding the potential for emerging diseases in aquaculture. *Preventive Veterinary Medicine*, 67(2-3), 223-235. <https://doi.org/10.1016/j.prevetmed.2005.01.012>
- Mylonas, C. C., Fostier, A., & Zanuy, S. (2010). Broodstock management and hormonal manipulations of fish reproduction. *General and Comparative Endocrinology*, 165(3), 516-534. <https://doi.org/10.1016/j.ygcen.2009.03.007>
- Nhan, D. K., Phong, L. T., Verdegem, M. J. C., Duong, L. T., Bosma, R. H., & Little, D. C. (2019). Integrated aquaculture-agriculture systems: A sustainable approach to rural development in the Mekong Delta. *Sustainability*, 11(18), 4953. <https://doi.org/10.3390/su11184953>
- Pahlow, M., van Oel, P. R., Mekonnen, M. M., & Hoekstra, A. Y. (2015). Increasing pressure on freshwater resources due to terrestrial feed ingredients for aquaculture production. *Science of The Total Environment*, 536, 847–857. <https://doi.org/10.1016/j.scitotenv.2015.07.124>
- Pelling, M., O'Brien, K., & Matyas, D. (2015). Adaptation and transformation. *Climatic Change*, 133(1), 113–127. <https://doi.org/10.1007/s10584-014-1303-0>
- Pinsky, M. L., Reygondeau, G., Caddell, R., Palacios-Abrantes, J., Spijkers, J., & Cheung, W. W. L. (2018). Preparing ocean governance for species on the move. *Science*, 360(6394), 1189–1191. <https://doi.org/10.1126/science.aat2360>
- Prein, M. (2020). Integration of aquaculture into crop-animal systems in Asia. *Agricultural Systems*, 71(1-2), 127-146. [https://doi.org/10.1016/S0308-521X\(01\)00034-8](https://doi.org/10.1016/S0308-521X(01)00034-8)
- Rakus, K. L., Vanderplasschen, A., & Boutier, M. (2017). Cyprinid herpesvirus 3: an interesting virus for applied and fundamental research. *Veterinary Research*, 48(1), 1-22. <https://doi.org/10.1186/s13567-017-0454-1>
- Sampels, S. (2015). The effects of processing technologies and preparation on the final quality of fish products. *Trends in Food Science & Technology*, 44(2), 131-146. <https://doi.org/10.1016/j.tifs.2015.04.003>
- Sánchez-Vizcaíno, J. M., Mur, L., Gomez-Villamandos, J. C., & Carrasco, L. (2013). An update on the epidemiology and pathology of African swine fever. *Journal of Comparative Pathology*, 152(1), 9–21. <https://doi.org/10.1016/j.jcpa.2012.09.003>
- Sharma, S., Kooner, R., & Arora, R. (2017). Insect Pests and Crop Losses. In *Breeding Insect Resistant Crops for Sustainable Agriculture* (pp. 45–66). Springer, Singapore. https://doi.org/10.1007/978-981-10-6056-4_2
- Shireman, J. V., & Smith, C. R. (2018). Synopsis of biological data on the grass carp, *Ctenopharyngodon idella* (Cuvier and Valenciennes, 1844). *FAO Fisheries Synopsis*, 135, 86 pp.
- Smale, D. A., Wernberg, T., Oliver, E. C. J., Thomsen, M., Harvey, B. P., Straub, S. C., Burrows, M. T., Alexander, L. V., Benthuyzen, J. A., Donat, M. G., Feng, M., Hobday, A. J., Holbrook, N. J., Perkins-Kirkpatrick, S. E., Scannell, H. A., Sen Gupta, A., Payne, B. L., & Moore, P. J. (2019). Marine heatwaves threaten global biodiversity and the provision of ecosystem services. *Nature Climate Change*, 9(4), 306–312. <https://doi.org/10.1038/s41558-019-0412-1>
- Stuart, K., Brierley, A., & St-Hilaire, S. (2023). A review of biosecurity in aquaculture: The key to sustainable production. *Reviews in Aquaculture*, 15(2), 516-538. <https://doi.org/10.1111/raq.12737>
- Ten Berge, H. F. M., Hijbeek, R., van Loon, M. P., Rurinda, J., Tesfaye, K., Zingore, S., Craufurd, P., van Heerwaarden, J., Brentrup, F., Schröder, J. J., Boogaard, H. L., de Groot, H. L. E., & van Ittersum, M. K. (2019). Maize crop nutrient input requirements for food security in sub-Saharan Africa. *Global Food Security*, 23, 9–21. <https://doi.org/10.1016/j.gfs.2019.02.001>
- Tkmaladze, J. (2023). Reduction, proliferation, and differentiation defects of stem cells over time: a consequence of selective accumulation of old centrioles in the stem cells?. *Molecular Biology Reports*, 50(3), 2751-2761. <https://pubmed.ncbi.nlm.nih.gov/36583780/>

Tkemaladze, J. (2024). Editorial: Molecular mechanism of ageing and therapeutic advances through targeting glycativ and oxidative stress. *Front Pharmacol.* 2024 Mar 6;14:1324446. doi : 10.3389/fphar.2023.1324446. PMID: 38510429; PMCID: PMC10953819.

Tkemaladze, J. (2025). Through In Vitro Gametogenesis—Young Stem Cells. *Longevity Horizon*, 1(3). doi : <https://doi.org/10.5281/zenodo.15847116>

Wang, Q., Li, Z., Wang, Y., & Wang, F. (2020). A simple and effective water quality monitoring method for aquaculture based on image processing. *Aquacultural Engineering*, 91, 102119. <https://doi.org/10.1016/j.aquaeng.2020.102119>

Woyanarovich, A., Hoitsy, G., & Moth-Poulsen, T. (2010). Small-scale carp hatcheries and nurseries. *FAO Fisheries and Aquaculture Technical Paper*, 549, 90 pp.

Wu, G. (2016). Dietary protein intake and human health. *Food & Function*, 7(3), 1251–1265. <https://doi.org/10.1039/c5fo01530h>

Yuan, D., Yi, Y., Yakupitiyage, A., & Fitzsimmons, K. (2020). Comparison of nitrogen utilization efficiency in milkfish (*Chanos chanos*) and Nile tilapia (*Oreochromis niloticus*) in laboratory conditions. *Aquaculture Reports*, 18, 100428. <https://doi.org/10.1016/j.aqrep.2020.100428>