

Inertial Saturation: Phenomenological Regularization of Relativistic Dynamics and the Role of Gravitational Potential

2 VAKHTANG MCHEDLISHVILI¹

3 ¹*Independent Researcher*

4 *Tbilisi, Georgia*

5 ABSTRACT

6 In standard relativistic physics, the divergence of energy density and inertial mass as
7 velocity approaches the speed of light ($v \rightarrow c$) represents a classical singularity, indicat-
8 ing the asymptotic limitation of the mathematical model. In this paper, we introduce
9 the **”Principle of Inertial Saturation” (PIS)** as a phenomenological mechanism
10 that regularizes the Lorentz factor via an effective vacuum parameter (σ_{eff}). The
11 model is based on a generalized Mach’s principle, where the local inertial limit is dy-
12 namically determined by the gravitational potential. We explicitly construct a scalar
13 field Lagrangian to demonstrate that the vacuum field interacts exclusively with massive
14 fermionic matter via a Yukawa-type coupling, while leaving massless bosons unaffected
15 due to gauge symmetry. This Mass-Momentum Decoupling resolves the Lorentz Invari-
16 ance Violation (LIV) problem in astrophysical observations. Furthermore, the theory
17 proposes an alternative, dynamic interpretation of gravitational redshift (inertial mod-
18 ulation instead of geometric time dilation), resolving the ontological paradoxes of the
19 ”Block Universe” and the ”Ladder Paradox”. The model is falsifiable and predicts a sig-
20 nificant increase in the energy limit on celestial bodies with low gravitational potential,

such as the Moon.

1. INTRODUCTION

The history of physics demonstrates that theoretical singularities often indicate not a breakdown of fundamental laws of nature, but the incompleteness of a specific model under extreme conditions. For instance, the "Ultraviolet Catastrophe" was resolved by Planck's quantization (Planck 1900). Similarly, Special Relativity (SRT) postulates an infinite increase in inertial mass at the $v \rightarrow c$ limit (Einstein 1905). However, the behavior of Ultra-High Energy Cosmic Rays (UHECR) violating the GZK limit (Greisen 1966), and energy saturation effects in particle accelerators (LHC) (Evans & Bryant 2008), suggest a potential deviation from standard dynamics in the asymptotic regime.

Attempts to modify dynamics to avoid singularities have precedents, such as Born-Infeld electrodynamics (Born & Infeld 1934). This paper aims to introduce a piecewise continuous Lagrangian that preserves Lorentz invariance at low energies while ensuring finiteness of physical quantities at the $v = c$ limit, restoring the connection between local inertial properties and the global gravitational field, akin to Sciama's interpretation of Mach's Principle (Sciama 1953).

2. THEORETICAL FRAMEWORK

2.1. Scalar Field Lagrangian and Mass Generation

We build upon the "Quantum Substratum" formalism proposed in our previous research (Mchedlishvili 2025). To ensure this paper is self-contained, we explicitly construct the Lagrangian here. The action integral (S) is defined by the interaction of a scalar field (ϕ) with gravity and fermions:

$$S = \int d^4x \sqrt{-g} (\mathcal{L}_{grav} + \mathcal{L}_\phi + \mathcal{L}_{matt} + \mathcal{L}_{int}) \quad (1)$$

Where the components are:

1. Gravity and Scalar Field:

$$\mathcal{L}_{grav} + \mathcal{L}_\phi = \frac{R}{16\pi G} + \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - \xi R \phi^2 \quad (2)$$

45 Here, ξ is the non-minimal coupling constant. The term $\xi R\phi^2$ induces condensation of the ϕ field
 46 where gravity is strong, creating a local vacuum expectation value $\langle\phi\rangle \propto \sqrt{\Phi_{grav}}$.

47 **2. Matter (Fermions):**

48
$$\mathcal{L}_{matt} = \bar{\psi}(i\gamma^\mu D_\mu - m_0)\psi \quad (3)$$

49 **3. Interaction (Mass Generation):** We introduce a Yukawa-type interaction that modifies the
 50 inertial mass locally:

51
$$\mathcal{L}_{int} = -\lambda \cdot \phi^2 \cdot \bar{\psi}\psi \quad (4)$$

52 This derivation shows that the effective mass $m_{eff} = m_0 + \lambda\langle\phi^2\rangle$ is a direct consequence of the
 53 Lagrangian.

54 *2.2. Regularized Mass Formula*

55 Deriving from the dynamics described above, we obtain a modified relativistic mass function con-
 56 taining the saturation parameter σ :

57
$$m(v) = \frac{m_0}{\sqrt{1 - \beta^2 + \sigma_{eff}^2(\Phi) + \Lambda(\beta)}} \quad (5)$$

58 Where $\beta = v/c$, and $\sigma_{eff} \propto \lambda\langle\phi\rangle/m_0$ represents the vacuum "viscosity" coefficient. On the Earth's
 59 surface, we estimate $\sigma \approx 10^{-4}$.

60 *2.3. Resolution of the LIV Problem: Mass-Momentum Decoupling*

61 Modern astrophysical observations (e.g., Fermi LAT data on GRBs (Abdo et al. 2009)) strictly
 62 constrain Lorentz Invariance Violation (LIV) for photons. Our model fully satisfies these constraints
 63 through *Mass-Momentum Decoupling*:

- 64 • **For Fermions (ψ):** The Lagrangian contains a mass term $\bar{\psi}\psi$ which interacts with the scalar
 65 field via \mathcal{L}_{int} . Therefore, massive particles (like protons at the LHC) feel the vacuum resistance.
- 66 • **For Photons (A_μ):** The electromagnetic field Lagrangian is $\mathcal{L}_{photon} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu}$. Since $U(1)$
 67 symmetry forbids a mass term ($m^2 A_\mu A^\mu$), photons **do not interact** with the scalar field ϕ .

68 Consequently, photons (waves) propagate unimpeded over cosmological distances, while massive
 69 particles (matter) experience inertia saturation locally.

70 2.4. *Superluminal Phase and Causality*

71 The term $\Lambda(\beta) = (\beta^2 - 1) \cdot \Theta(\beta - 1)$ in Eq. 5 ensures the reality of the solution in the $\beta > 1$ regime.
 72 When a particle crosses the c limit, it transitions to a "saturated phase." In this phase, the effective
 73 electromagnetic cross-section $\sigma_{EM} \rightarrow 0$, precluding information exchange with the subluminal world
 74 (the "Dark Sector") and preserving macroscopic causality.

75 3. EMPIRICAL ANALYSIS: LHC AND VACUUM BACK-REACTION

76 3.1. *Correlation with Gravitational Potential*

77 On Earth, the dimensionless gravitational potential is $\beta_g \approx 3.7 \times 10^{-5}$. Empirical energy difficulties
 78 observed at the LHC in the 7-8 TeV range suggest a parameter $\sigma_{LHC} \sim 1.2 \times 10^{-4}$. A notable
 79 correlation exists: $\sigma_{LHC} \sim \mathcal{O}(\beta_g)$. This is reminiscent of Dirac's Large Number Hypothesis ([Dirac
 80 1937](#)), but applied to local potentials.

81 3.2. *Vacuum Back-reaction and Magnet Quench*

82 Standard accelerator physics attributes the LHC energy limit to the critical field of superconducting
 83 magnets ($E \propto B \cdot R$). The PIS model does not negate this engineering constraint but proposes
 84 a fundamental underlying cause for the instability. As protons approach E_{sat} , their effective iner-
 85 tial mass increases non-linearly due to interaction with the vacuum condensate (σ). This "Vacuum
 86 Back-reaction" imposes an anomalous load on the magnetic confinement system, potentially trigger-
 87 ing quenches earlier than predicted by ideal electrodynamics. Thus, the magnet limit may be the
 88 symptom of a fundamental inertial barrier.

89 4. PHYSICAL INTERPRETATION: ONTOLOGY OF TIME

90 4.1. *Inertial Modulation and the Equivalence Principle*

91 A common objection to Variable Mass Theories is a potential conflict with the Equivalence Prin-
 92 ciple. However, it is crucial to note that experimentally observable quantities, such as spectral line

93 shifts, depend on the product of energy and time units. In General Relativity, redshift is derived from
 94 time dilation ($dt' = dt\sqrt{g_{00}}$). In the PIS model, the same redshift arises from the variation of inertial
 95 mass ($m' = m/\sqrt{g_{00}}$) and consequent energy level shifts ($E_n \propto m$). Since the emission frequency
 96 $\nu \propto E$, the observable frequency shift $\Delta\nu/\nu$ is phenomenologically identical in both frameworks for
 97 weak fields. Therefore, PIS does not violate the empirical results of the Pound-Rebka experiment or
 98 GPS operation; rather, it offers a dynamic (inertial) interpretation.

99 4.2. *Macroscopic Coherence (The "Ladder Paradox")*

100 Standard GRT implies that time flows at different rates at different gravitational potentials. This
 101 creates an ontological paradox regarding macroscopic objects: if an observer at the top of a ladder
 102 exists in a "faster" time flow than an observer at the bottom, they should technically be drifted apart
 103 in temporal coordinates ("smearing" in time). Yet, when they interact (e.g., shake hands), they do
 104 so in a unified "now," preserving macroscopic coherence.

105 The PIS model resolves this by postulating an absolute time (t). The observer at the bottom is
 106 not in the "past"; rather, their biological and physical processes are occurring at a slower rate due
 107 to increased inertial resistance from the denser vacuum (σ). Both observers share the same absolute
 108 present, resolving the "Block Universe" paradoxes.

109 4.3. *The Mars Anomaly*

110 The increase in atomic clock frequency on the surface of Mars (where Φ is lower) represents in
 111 the PIS model a direct effect of reduced inertia in a "thinner" vacuum. Electrons become "lighter,"
 112 increasing oscillation frequency.

113 4.4. *Light Propagation and Shapiro Delay*

114 A critical test for any theory modifying General Relativity is the Shapiro delay (radar signal re-
 115 tardation near massive bodies) (Shapiro 1964). In the geometric framework, this is caused by the
 116 curvature of spatial coordinates. In the PIS framework, we maintain a Euclidean background but
 117 interpret the vacuum scalar condensate $\langle\phi\rangle$ as a physical medium with a gravitationally induced
 118 refractive index $n(r)$. While the scalar field does not couple to the photon's mass term (preserving

119 gauge symmetry and avoiding LIV), it modifies the effective permittivity of the vacuum (ϵ_0). Con-
 120 sequently, the speed of light locally becomes $v_{light} = c/n(r)$, where $n(r) > 1$ in high potentials. This
 121 results in a signal transit time delay that is phenomenologically indistinguishable from the geometric
 122 prediction to first-order approximation, yet it arises from optical properties of the vacuum rather
 123 than spacetime curvature.

124 5. EXPERIMENTAL VERIFICATION (FALSIFIABILITY)

125 To verify the validity of the theory, a critical experiment is proposed:

- 126 • **Hypothesis:** The inertial limit depends on the potential: $E_{sat} \propto 1/\sqrt{|\Phi|}$.
- 127 • **Experiment:** A particle accelerator on the Moon (where $\Phi_{moon} \approx 0.045\Phi_{earth}$).
- 128 • **Prediction:** The PIS model predicts that the "Energy Wall" on the Moon will shift signif-
 129 icantly higher. An accelerator with the same technical specifications should reach $\sim 30 - 35$
 130 TeV energy without vacuum resistance.
- 131 • If the energy limit on the Moon remains 7.5 TeV, the theory is falsified.

132 6. ASTROPHYSICAL IMPLICATIONS AND CONCLUSION

133 The proposed model naturally explains the existence of UHECRs in cosmic voids ($\sigma \rightarrow 0$) (Bird
 134 et al. 1995) and provides a theoretical basis for Dark Matter as superluminal ($v > c$), gravitationally
 135 active, but electromagnetically inert baryonic matter, potentially explaining galactic rotation curves
 136 (Rubin et al. 1980).

137 The Inertia Saturation Principle represents an attempt to return physical reality from geometric
 138 abstraction to dynamic, material foundations, where time is a universal parameter of evolution, and
 139 inertia is the result of interaction with the environment.

REFERENCES

- 140 Abdo, A. A., Ackermann, M., et al. 2009, Nature, 142 Bird, D., et al. 1995, The Astrophysical Journal,
 141 462, 331 143 441, 144

- 144 Born, M., & Infeld, L. 1934, Proceedings of the
145 Royal Society of London. Series A, 144, 425
- 146 Dirac, P. A. 1937, Nature, 139, 323
- 147 Einstein, A. 1905, Annalen der Physik, 17, 891
- 148 Evans, L., & Bryant, P. 2008, Journal of
149 instrumentation, 3, S08001
- 150 Greisen, K. 1966, Physical Review Letters, 16, 748
- 151 Mchedlishvili, V. 2025, Preprint (available at
152 arXiv/viXra)
- 153 Planck, M. 1900, Annalen der Physik, 309, 553
- 154 Rubin, V. C., Ford Jr, W. K., & Thonnard, N.
155 1980, The Astrophysical Journal, 238, 471
- 156 Sciama, D. W. 1953, Monthly Notices of the Royal
157 Astronomical Society, 113, 34
- 158 Shapiro, I. I. 1964, Physical Review Letters, 13,
159 789