

# Mathematical Analysis of a Great Rhombicosidodecahedron

Harish Chandra Rajpoot

M.M.M. University of Technology, Gorakhpur-273010 (UP), India

10 March, 2015

## 1. Introduction

A great rhombicosidodecahedron is the largest Archimedean solid which has 30 congruent square faces, 20 congruent regular hexagonal faces & 12 congruent regular decagonal faces each having equal edge length [1]. It has 180 edges & 120 vertices lying on a spherical surface with a certain radius. It is created/generated by expanding a truncated dodecahedron [2] having 20 equilateral triangular faces & 12 regular decagonal faces. Thus by the expansion, each of 30 originally truncated edges changes into a square face, each of 20 triangular faces of the original solid changes into a regular hexagonal face & 12 regular decagonal faces of the original solid remain unchanged i.e. decagonal faces are shifted radially. Thus a solid with 30 squares, 20 hexagonal & 12 decagonal faces, is obtained which is called great rhombicosidodecahedron which is the largest Archimedean solid (see figure 1), thus we have




Figure 1: A great rhombicosidodecahedron having 30 congruent square faces, 20 congruent regular hexagonal faces & 12 congruent regular decagonal faces each of equal edge length  $a$ .

no. of square faces = 30, no. of regular hexagonal faces = 20

no. of regular decagonal faces = 12,

$$\begin{aligned} \text{no. of edges} &= 12(\text{no. of edges in a decagonal face}) + 2(\text{no. of square faces}) \\ &= 12(10) + 2(30) = 120 + 60 = 180 \end{aligned}$$

no. of vertices = 12(no. of vertices in a decagonal face) = 12(10) = 120

We will apply Theory of Polygon [3,4] to derive a mathematical relationship between radius  $R$  of the spherical surface passing through all 120 vertices & the edge length  $a$  of a great rhombicosidodecahedron for calculating its important parameters such as normal distance of each face, surface area, volume etc.

## 2. Derivation of outer (circumscribed) radius ( $R_o$ ) of great rhombicosidodecahedron

Let  $R_o$  be the radius of the spherical surface passing through all 120 vertices of a great rhombicosidodecahedron with edge length  $a$  & the centre  $O$ . Consider a square face  $ABCD$  with centre  $O_1$ , regular hexagonal face  $EFGHIJ$  with centre  $O_2$  & regular decagonal face  $KLMNPQRSTU$  with centre  $O_3$  (see the figure 2 below).

Now, **normal distance ( $H_s$ ) of square face**  $ABCD$  from the centre  $O$  of the great rhombicosidodecahedron is calculated as follows

In right  $\Delta AO_1O$  (figure 2),

$$OO_1 = \sqrt{(OA)^2 - (O_1A)^2} \quad \left( O_1A = \frac{a}{\sqrt{2}} = \text{circumscribed radius of square} \right)$$

$$\Rightarrow H_s = \sqrt{(R_o)^2 - \left(\frac{a}{\sqrt{2}}\right)^2} = \sqrt{\frac{2R_o^2 - a^2}{2}} \quad \dots \dots \dots (I)$$




Figure 2: A square face ABCD, regular hexagonal face EFGHIJ & regular decagonal face KLMNQRSTU each of edge length  $a$ .

We know that the solid angle ( $\omega$ ) subtended by any regular polygon with each side of length  $a$  at any point lying at a distance  $H$  on the vertical axis passing through the centre of plane is given by HCR's Theory of Polygon [3,4] as follows

$$\omega = 2\pi - 2n \sin^{-1} \left( \frac{2H \sin \frac{\pi}{n}}{\sqrt{4H^2 + a^2 \cot^2 \frac{\pi}{n}}} \right)$$

Hence, by substituting the corresponding values in the above expression, we get the solid angle ( $\omega_s$ ) subtended by each square face (ABCD) at the centre of the great rhombicosidodecahedron as follows

$$\begin{aligned} \omega_s &= 2\pi - 2 \times 4 \sin^{-1} \left( \frac{2 \left( \sqrt{\frac{2R_o^2 - a^2}{2}} \right) \sin \frac{\pi}{4}}{\sqrt{4 \left( \sqrt{\frac{2R_o^2 - a^2}{2}} \right)^2 + a^2 \cot^2 \frac{\pi}{4}}} \right) \\ &= 2\pi - 8 \sin^{-1} \left( \frac{\sqrt{2} \sqrt{2R_o^2 - a^2} \times \frac{1}{\sqrt{2}}}{\sqrt{4R_o^2 - 2a^2 + a^2(1)^2}} \right) = 2\pi - 8 \sin^{-1} \left( \frac{\sqrt{2R_o^2 - a^2}}{\sqrt{4R_o^2 - a^2}} \right) = 2\pi - 8 \sin^{-1} \left( \sqrt{\frac{2R_o^2 - a^2}{4R_o^2 - a^2}} \right) \end{aligned}$$

Let  $\frac{R_o}{a} = x > 1$  (any arbitrary variable)

$$\Rightarrow \omega_s = 2\pi - 8 \sin^{-1} \left( \sqrt{\frac{2 \left( \frac{R_o}{a} \right)^2 - 1}{4 \left( \frac{R_o}{a} \right)^2 - 1}} \right) = 2\pi - 8 \sin^{-1} \left( \sqrt{\frac{2x^2 - 1}{4x^2 - 1}} \right) \dots \dots \dots (II)$$

Similarly, **Normal distance** ( $H_h$ ) of **regular hexagonal face** EFGHIJ from the centre O of the great rhombicosidodecahedron is calculated as follows

In right  $\Delta EO_2O$  (figure 2 above),

$$OO_2 = \sqrt{(OE)^2 - (O_2E)^2} \quad (O_2E = a = \text{circumscribed radius of regular hexagon})$$

$$\Rightarrow H_h = \sqrt{(R_o)^2 - (a)^2} = \sqrt{R_o^2 - a^2} \quad \dots \dots \dots (III)$$

Hence, by substituting all the corresponding values, the solid angle ( $\omega_h$ ) subtended by each regular hexagonal face (EFGHIJ) at the centre of the great rhombicosidodecahedron is given as follows

$$\begin{aligned} \omega_h &= 2\pi - 2 \times 6 \sin^{-1} \left( \frac{2(\sqrt{R_o^2 - a^2}) \sin \frac{\pi}{6}}{\sqrt{4(\sqrt{R_o^2 - a^2})^2 + a^2 \cot^2 \frac{\pi}{6}}} \right) \\ &= 2\pi - 12 \sin^{-1} \left( \frac{2\sqrt{R_o^2 - a^2} \times \frac{1}{2}}{\sqrt{4R_o^2 - 4a^2 + a^2(\sqrt{3})^2}} \right) = 2\pi - 12 \sin^{-1} \left( \frac{\sqrt{R_o^2 - a^2}}{\sqrt{4R_o^2 - a^2}} \right) \\ \Rightarrow \omega_h &= 2\pi - 12 \sin^{-1} \left( \frac{\sqrt{R_o^2 - a^2}}{\sqrt{4R_o^2 - a^2}} \right) = 2\pi - 12 \sin^{-1} \left( \frac{\sqrt{x^2 - 1}}{\sqrt{4x^2 - 1}} \right) \quad \dots \dots \dots (IV) \end{aligned}$$

Similarly, **Normal distance** ( $H_D$ ) of **regular decagonal face** KLMNPQRSTU from the centre O of the great rhombicosidodecahedron is calculated as follows

In right  $\Delta KO_3O$  (figure 2 above),

$$\Rightarrow OO_3 = \sqrt{(OE)^2 - (O_3K)^2}$$

$$O_3K = \text{circumscribed radius of regular decagon} = \frac{a}{2} \operatorname{cosec} 18^\circ = \frac{a}{2} (1 + \sqrt{5})$$

$$\Rightarrow H_D = \sqrt{(R_o)^2 - \left(\frac{a}{2}(1 + \sqrt{5})\right)^2} = \sqrt{(R_o)^2 - \frac{a^2}{4} (6 + 2\sqrt{5})} = \sqrt{\frac{2R_o^2 - (3 + \sqrt{5})a^2}{2}} \quad \dots \dots \dots (V)$$

Hence, by substituting all the corresponding values, the solid angle ( $\omega_D$ ) subtended by each regular decagonal face (KLMNPQRSTU) at the centre of great rhombicosidodecahedron is given as follows

$$\omega_D = 2\pi - 2 \times 10 \sin^{-1} \left( \frac{2 \left( \sqrt{\frac{2R_o^2 - (3 + \sqrt{5})a^2}{2}} \right) \sin \frac{\pi}{10}}{\sqrt{4 \left( \sqrt{\frac{2R_o^2 - (3 + \sqrt{5})a^2}{2}} \right)^2 + a^2 \cot^2 \frac{\pi}{10}}} \right)$$

$$\begin{aligned}
 &= 2\pi - 20 \sin^{-1} \left( \frac{\sqrt{4R_o^2 - 2(3 + \sqrt{5})a^2} \left( \frac{\sqrt{5} - 1}{4} \right)}{\sqrt{4R_o^2 - 2(3 + \sqrt{5})a^2 + a^2 (\sqrt{5 + 2\sqrt{5}})^2}} \right) \\
 &= 2\pi - 20 \sin^{-1} \left( \frac{\sqrt{4 \left( \frac{6 - 2\sqrt{5}}{16} \right) R_o^2 - 2(3 + \sqrt{5}) \left( \frac{6 - 2\sqrt{5}}{16} \right) a^2}}{\sqrt{4R_o^2 - 2(3 + \sqrt{5})a^2 + (5 + 2\sqrt{5})a^2}} \right) \\
 &= 2\pi - 20 \sin^{-1} \left( \frac{\sqrt{\left( \frac{3 - \sqrt{5}}{2} \right) R_o^2 - a^2}}{\sqrt{4R_o^2 - a^2}} \right) \\
 \omega_D &= 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{\left( \frac{3 - \sqrt{5}}{2} \right) R_o^2 - a^2}{4R_o^2 - a^2}} \right) = 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{\left( \frac{3 - \sqrt{5}}{2} \right) x^2 - 1}{4x^2 - 1}} \right) \dots \dots \dots (VI)
 \end{aligned}$$

Since a great rhombicosidodecahedron is a closed surface & we know that the total solid angle, subtended by any closed surface at any point lying inside it, is  $4\pi$  sr (Ste-radian) [5] hence the sum of solid angles subtended by 30 congruent square faces, 20 congruent regular hexagonal faces & 12 congruent regular decagonal faces at the centre of the great rhombicosidodecahedron must be  $4\pi$  sr. Thus we have

$$30[\omega_s] + 20[\omega_h] + 12[\omega_D] = 4\pi$$

Now, by substituting the values of  $\omega_s$ ,  $\omega_h$  &  $\omega_D$  from eq(II), (IV) & (VI) in the above expression we get

$$\begin{aligned}
 &30 \left[ 2\pi - 8 \sin^{-1} \left( \sqrt{\frac{2x^2 - 1}{4x^2 - 1}} \right) \right] + 20 \left[ 2\pi - 12 \sin^{-1} \left( \sqrt{\frac{x^2 - 1}{4x^2 - 1}} \right) \right] \\
 &\quad + 12 \left[ 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{\left( \frac{3 - \sqrt{5}}{2} \right) x^2 - 1}{4x^2 - 1}} \right) \right] = 4\pi \\
 \Rightarrow &240 \left[ \sin^{-1} \left( \sqrt{\frac{2x^2 - 1}{4x^2 - 1}} \right) + \sin^{-1} \left( \sqrt{\frac{x^2 - 1}{4x^2 - 1}} \right) + \sin^{-1} \left( \sqrt{\frac{\left( \frac{3 - \sqrt{5}}{2} \right) x^2 - 1}{4x^2 - 1}} \right) \right] = 124\pi - 4\pi = 120\pi \\
 \Rightarrow &\sin^{-1} \left( \sqrt{\frac{2x^2 - 1}{4x^2 - 1}} \right) + \sin^{-1} \left( \sqrt{\frac{x^2 - 1}{4x^2 - 1}} \right) + \sin^{-1} \left( \sqrt{\frac{\left( \frac{3 - \sqrt{5}}{2} \right) x^2 - 1}{4x^2 - 1}} \right) = \frac{120\pi}{240} = \frac{\pi}{2}
 \end{aligned}$$

$$\begin{aligned} &\Rightarrow \sin^{-1}\left(\sqrt{\frac{2x^2-1}{4x^2-1}}\right) + \sin^{-1}\left(\sqrt{\frac{x^2-1}{4x^2-1}}\right) = \frac{\pi}{2} - \sin^{-1}\left(\sqrt{\frac{\left(\frac{3-\sqrt{5}}{2}\right)x^2-1}{4x^2-1}}\right) \\ &\Rightarrow \sin^{-1}\left(\sqrt{\frac{2x^2-1}{4x^2-1}}\sqrt{1-\left(\sqrt{\frac{x^2-1}{4x^2-1}}\right)^2} + \sqrt{\frac{x^2-1}{4x^2-1}}\sqrt{1-\left(\sqrt{\frac{2x^2-1}{4x^2-1}}\right)^2}\right) = \cos^{-1}\left(\sqrt{\frac{\left(\frac{3-\sqrt{5}}{2}\right)x^2-1}{4x^2-1}}\right) \\ &\quad (\because \sin^{-1} X + \sin^{-1} Y = \sin^{-1}(X\sqrt{1-Y^2} + Y\sqrt{1-X^2}) \quad \& \quad \frac{\pi}{2} - \sin^{-1} X = \cos^{-1} X) \\ &\Rightarrow \sin^{-1}\left(\sqrt{\frac{2x^2-1}{4x^2-1}}\sqrt{\frac{3x^2}{4x^2-1}} + \sqrt{\frac{x^2-1}{4x^2-1}}\sqrt{\frac{2x^2}{4x^2-1}}\right) = \sin^{-1}\left(\sqrt{1-\left(\sqrt{\frac{\left(\frac{3-\sqrt{5}}{2}\right)x^2-1}{4x^2-1}}\right)^2}\right) \\ &\Rightarrow \sin^{-1}\left(\frac{x\sqrt{3(2x^2-1)}}{4x^2-1} + \frac{x\sqrt{2(x^2-1)}}{4x^2-1}\right) = \sin^{-1}\left(\sqrt{\frac{\left(\frac{5+\sqrt{5}}{2}\right)x^2}{4x^2-1}}\right) = \sin^{-1}\left(x\sqrt{\frac{(5+\sqrt{5})}{2(4x^2-1)}}\right) \\ &\quad \Rightarrow \frac{x\sqrt{3(2x^2-1)}}{4x^2-1} + \frac{x\sqrt{2(x^2-1)}}{4x^2-1} = x\sqrt{\frac{(5+\sqrt{5})}{2(4x^2-1)}} \\ &\quad \Rightarrow \sqrt{2}\sqrt{3(2x^2-1)} + \sqrt{2}\sqrt{2(x^2-1)} = \sqrt{(5+\sqrt{5})(4x^2-1)} \\ &\quad \Rightarrow \left(\sqrt{6(2x^2-1)} + 2\sqrt{(x^2-1)}\right)^2 = \left(\sqrt{(5+\sqrt{5})(4x^2-1)}\right)^2 \\ &\quad \Rightarrow 6(2x^2-1) + 4(x^2-1) + 4\sqrt{6(x^2-1)(2x^2-1)} = (5+\sqrt{5})(4x^2-1) \\ &\quad \Rightarrow 4(5+\sqrt{5})x^2 - (5+\sqrt{5}) - 16x^2 + 10 = 4\sqrt{6(x^2-1)(2x^2-1)} \\ &\quad \Rightarrow \left(4(1+\sqrt{5})x^2 + (5-\sqrt{5})\right)^2 = \left(4\sqrt{6(x^2-1)(2x^2-1)}\right)^2 \\ &\quad \Rightarrow 32(3+\sqrt{5})x^4 + 32\sqrt{5}x^2 + 10(3-\sqrt{5}) = 96(x^2-1)(2x^2-1) = 192x^4 - 288x^2 + 96 \\ &\quad \Rightarrow (96 - 32\sqrt{5})x^4 - (288 + 32\sqrt{5})x^2 + (66 + 10\sqrt{5}) = 0 \\ &\quad \Rightarrow 32(3-\sqrt{5})x^4 - 32(9+\sqrt{5})x^2 + 2(33+5\sqrt{5}) = 0 \end{aligned}$$

Now, solving the above biquadratic equation for the values of  $x > 1$  as follows

$$\begin{aligned} \Rightarrow x^2 &= \frac{-(-32(9 + \sqrt{5})) \pm \sqrt{(-32(9 + \sqrt{5}))^2 - 4(32(3 - \sqrt{5}))(2(33 + 5\sqrt{5}))}}{2 \times 32(3 - \sqrt{5})} \\ &= \frac{32(9 + \sqrt{5}) \pm \sqrt{1024(86 + 18\sqrt{5}) - 256(3 - \sqrt{5})(33 + 5\sqrt{5})}}{64(3 - \sqrt{5})} \\ &= \frac{32(9 + \sqrt{5}) \pm 16\sqrt{4(86 + 18\sqrt{5}) - (74 - 18\sqrt{5})}}{64(3 - \sqrt{5})} = \frac{2(9 + \sqrt{5}) \pm \sqrt{270 + 90\sqrt{5}}}{4(3 - \sqrt{5})} \\ &= \frac{(2(9 + \sqrt{5}) \pm \sqrt{270 + 90\sqrt{5}})(3 + \sqrt{5})}{4(3 - \sqrt{5})(3 + \sqrt{5})} = \frac{2(9 + \sqrt{5})(3 + \sqrt{5}) \pm (3 + \sqrt{5})\sqrt{9(5 + \sqrt{5})^2}}{4(4)} \\ &= \frac{2(9 + \sqrt{5})(3 + \sqrt{5}) \pm 3(3 + \sqrt{5})(5 + \sqrt{5})}{16} = \frac{8(8 + 3\sqrt{5}) \pm 12(5 + 2\sqrt{5})}{16} \\ &= \frac{2(8 + 3\sqrt{5}) \pm 3(5 + 2\sqrt{5})}{4} = \frac{(16 + 6\sqrt{5}) \pm (15 + 6\sqrt{5})}{16} \end{aligned}$$

1. Taking positive sign, we have

$$x^2 = \frac{(16 + 6\sqrt{5}) + (15 + 6\sqrt{5})}{4} = \frac{31 + 12\sqrt{5}}{4} \quad \text{or} \quad x = \sqrt{\frac{31 + 12\sqrt{5}}{4}} = \frac{1}{2}\sqrt{31 + 12\sqrt{5}}$$

Since,  $x > 1$  hence, the above value is acceptable.

2. Taking negative sign, we have

$$x^2 = \frac{(16 + 6\sqrt{5}) - (15 + 6\sqrt{5})}{4} = \frac{1}{4} \quad \text{or} \quad x = \sqrt{\frac{1}{4}} = \frac{1}{2} \Rightarrow x < 1 \text{ but } x > 1 \text{ (required condition)}$$

Hence, the above value is discarded, now we have

$$x = \frac{1}{2}\sqrt{31 + 12\sqrt{5}} \Rightarrow \frac{R_o}{a} = x = \frac{1}{2}\sqrt{31 + 12\sqrt{5}} \quad \text{or} \quad R_o = \frac{a}{2}\sqrt{31 + 12\sqrt{5}}$$

Hence, outer (circumscribed) radius ( $R_o$ ) of a great rhombicosidodecahedron with edge length  $a$  is given as

$$R_o = \frac{a}{2}\sqrt{31 + 12\sqrt{5}} \approx 3.8023945a \quad \dots \dots \dots (VII)$$

**2.1. Normal distance ( $H_s$ ) of square faces from the centre of great rhombicosidodecahedron**

The normal distance ( $H_s$ ) of each of 30 congruent square faces from the centre O of a great rhombicosidodecahedron is given from eq(I) as follows

$$\begin{aligned}
H_s = OO_1 &= \sqrt{\frac{2R_o^2 - a^2}{2}} = \sqrt{\frac{2\left(\frac{a}{2}\sqrt{31+12\sqrt{5}}\right)^2 - a^2}{2}} = a\sqrt{\frac{31+12\sqrt{5}-2}{4}} = \frac{a}{2}\sqrt{29+12\sqrt{5}} \\
&= \frac{a}{2}\sqrt{(3+2\sqrt{5})^2} = \frac{(3+2\sqrt{5})a}{2} \\
\therefore H_s &= \frac{(3+2\sqrt{5})a}{2} \approx 3.736067978a
\end{aligned}$$

It's clear that all 30 congruent square faces are at an equal normal distance  $H_s$  from the centre of any great rhombicosidodecahedron.

## 2.2. Solid angle ( $\omega_s$ ) subtended by each of the square faces at the centre of great rhombicosidodecahedron

The solid angle ( $\omega_s$ ) subtended by each square face is given from eq(II) as follows

$$\omega_s = 2\pi - 8 \sin^{-1} \left( \sqrt{\frac{2x^2 - a^2}{4x^2 - a^2}} \right) = 2\pi - 8 \sin^{-1} \left( \sqrt{\frac{2R_o^2 - a^2}{4R_o^2 - a^2}} \right) \quad \left( \because x = \frac{R_o}{a} \right)$$

Hence, by substituting the corresponding value of  $R_o$  in the above expression, we get

$$\begin{aligned}
\omega_s &= 2\pi - 8 \sin^{-1} \left( \sqrt{\frac{2\left(\frac{a}{2}\sqrt{31+12\sqrt{5}}\right)^2 - a^2}{4\left(\frac{a}{2}\sqrt{31+12\sqrt{5}}\right)^2 - a^2}} \right) = 2\pi - 8 \sin^{-1} \left( \sqrt{\frac{31+12\sqrt{5}-2}{2(31+12\sqrt{5}-1)}} \right) \\
&= 2\pi - 8 \sin^{-1} \left( \sqrt{\frac{29+12\sqrt{5}}{12(5+2\sqrt{5})}} \right) = 2\pi - 8 \sin^{-1} \left( \sqrt{\frac{(29+12\sqrt{5})(5-2\sqrt{5})}{12(5+2\sqrt{5})(5-2\sqrt{5})}} \right) \\
&= 2\pi - 8 \sin^{-1} \left( \sqrt{\frac{25+2\sqrt{5}}{60}} \right) \\
\therefore \omega_s &= 2\pi - 8 \sin^{-1} \left( \frac{1}{2} \sqrt{\frac{25+2\sqrt{5}}{15}} \right) = 4 \sin^{-1} \left( \frac{5-2\sqrt{5}}{30} \right) \approx 0.070385504 \text{ sr}
\end{aligned}$$

## 2.3. Normal distance ( $H_h$ ) of regular hexagonal faces from the centre of great rhombicosidodecahedron

The normal distance ( $H_h$ ) of each of 20 congruent regular hexagonal faces from the centre O of a great rhombicosidodecahedron is given from eq(III) as follows

$$\begin{aligned}
H_h = OO_2 &= \sqrt{R_o^2 - a^2} = \sqrt{\left(\frac{a}{2}\sqrt{31+12\sqrt{5}}\right)^2 - a^2} = a\sqrt{\frac{31+12\sqrt{5}-4}{4}} = \frac{a}{2}\sqrt{27+12\sqrt{5}} \\
&= \frac{a}{2}\sqrt{3(2+\sqrt{5})^2} = \frac{\sqrt{3}(2+\sqrt{5})a}{2} \\
\therefore H_h &= \frac{\sqrt{3}(2+\sqrt{5})a}{2} \approx 3.668542481a
\end{aligned}$$

It's clear that all 20 congruent regular hexagonal faces are at an equal normal distance  $H_h$  from the centre of any great rhombicosidodecahedron.

#### 2.4. Solid angle ( $\omega_h$ ) subtended by each of the regular hexagonal faces at the centre of great rhombicosidodecahedron

The solid angle ( $\omega_h$ ) subtended by each regular hexagonal face is given from eq(IV) as follows

$$\omega_h = 2\pi - 12 \sin^{-1} \left( \sqrt{\frac{x^2 - a^2}{4x^2 - a^2}} \right) = 2\pi - 12 \sin^{-1} \left( \sqrt{\frac{R_o^2 - a^2}{4R_o^2 - a^2}} \right) \quad \left( \because x = \frac{R_o}{a} \right)$$

Hence, by substituting the corresponding value of  $R_o$  in the above expression, we get

$$\begin{aligned} \omega_h &= 2\pi - 12 \sin^{-1} \left( \sqrt{\frac{\left(\frac{a}{2}\sqrt{31+12\sqrt{5}}\right)^2 - a^2}{4\left(\frac{a}{2}\sqrt{31+12\sqrt{5}}\right)^2 - a^2}} \right) = 2\pi - 12 \sin^{-1} \left( \sqrt{\frac{31+12\sqrt{5}-4}{4(31+12\sqrt{5}-1)}} \right) \\ &= 2\pi - 12 \sin^{-1} \left( \sqrt{\frac{9+4\sqrt{5}}{8(5+2\sqrt{5})}} \right) = 2\pi - 12 \sin^{-1} \left( \sqrt{\frac{(9+4\sqrt{2})(5-2\sqrt{5})}{8(5+2\sqrt{5})(5-2\sqrt{5})}} \right) \\ &= 2\pi - 12 \sin^{-1} \left( \sqrt{\frac{5+2\sqrt{5}}{40}} \right) = 2\pi - 12 \sin^{-1} \left( \frac{1}{2} \sqrt{\frac{5+2\sqrt{5}}{10}} \right) \\ \therefore \omega_h &= 2\pi - 12 \sin^{-1} \left( \frac{1}{2} \sqrt{\frac{5+2\sqrt{5}}{10}} \right) \approx 0.184524626 \text{ sr} \end{aligned}$$

#### 2.5. Normal distance ( $H_D$ ) of regular decagonal faces from the centre of great rhombicosidodecahedron

The normal distance ( $H_D$ ) of each of 12 congruent regular decagonal faces from the centre O of a great rhombicosidodecahedron is given from eq(V) as follows

$$\begin{aligned} H_D = OO_3 &= \sqrt{\frac{2R_o^2 - (3+\sqrt{5})a^2}{2}} = \sqrt{\frac{2\left(\frac{a}{2}\sqrt{31+12\sqrt{5}}\right)^2 - (3+\sqrt{5})a^2}{2}} = \frac{a}{2} \sqrt{31+12\sqrt{5} - 2(3+\sqrt{5})} \\ &= \frac{a}{2} \sqrt{25+10\sqrt{5}} \end{aligned}$$

$$\therefore H_D = \frac{a}{2} \sqrt{25+10\sqrt{5}} \approx 3.440954801a$$

It's clear that all 12 congruent regular decagonal faces are at an equal normal distance  $H_D$  from the centre of any great rhombicosidodecahedron.

#### 2.6. Solid angle ( $\omega_D$ ) subtended by each of the regular decagonal faces at the centre of great rhombicosidodecahedron

The solid angle ( $\omega_D$ ) subtended by each regular decagonal face is given from eq(VI) as follows

$$\omega_D = 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{\left(\frac{3-\sqrt{5}}{2}\right)x^2 - 1}{4x^2 - 1}} \right) = 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{\left(\frac{3-\sqrt{5}}{2}\right)R_o^2 - a^2}{4R_o^2 - a^2}} \right) \quad \left( \because x = \frac{R_o}{a} \right)$$

Hence, by substituting the corresponding value of  $R_o$  in the above expression, we get

$$\begin{aligned}
 \omega_D &= 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{\left(\frac{3-\sqrt{5}}{2}\right) \left(\frac{a}{2} \sqrt{31+12\sqrt{5}}\right)^2 - a^2}{4 \left(\frac{a}{2} \sqrt{31+12\sqrt{5}}\right)^2 - a^2}} \right) \\
 &= 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{(3-\sqrt{5})(31+12\sqrt{5})-8}{8(31+12\sqrt{5}-1)}} \right) \\
 &= 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{25+5\sqrt{5}}{48(5+2\sqrt{5})}} \right) = 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{(25+5\sqrt{5})(5-2\sqrt{5})}{48(5+2\sqrt{5})(5-2\sqrt{5})}} \right) \\
 &= 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{75-25\sqrt{5}}{48(5)}} \right) = 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{15-5\sqrt{5}}{48}} \right) = 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{30-10\sqrt{5}}{96}} \right) \\
 &= 2\pi - 20 \sin^{-1} \left( \sqrt{\frac{(5-\sqrt{5})^2}{96}} \right) = 2\pi - 20 \sin^{-1} \left( \frac{5-\sqrt{5}}{4\sqrt{6}} \right) \\
 \therefore \omega_D &= 2\pi - 20 \sin^{-1} \left( \frac{5-\sqrt{5}}{4\sqrt{6}} \right) \approx 0.563692745 \text{ sr}
 \end{aligned}$$

It's clear from the above results that the solid angle subtended by each of 12 regular decagonal faces is greater than the solid angle subtended by each of 30 square faces & each of 20 regular hexagonal faces at the centre of any great rhombicosidodecahedron.

It's also clear from the above results that  $H_s > H_h > H_D$  i.e. the normal distance ( $H_s$ ) of square faces is greater than the normal distance  $H_h$  of the regular hexagonal faces & the normal distance  $H_D$  of the regular decagonal faces from the centre of a great rhombicosidodecahedron i.e. regular decagonal faces are closer to the centre as compared to the square & regular hexagonal faces in any great rhombicosidodecahedron.

### 3. Important parameters of a great rhombicosidodecahedron

- 1. Inner (inscribed) radius ( $R_i$ ):** It is the radius of the largest sphere inscribed (trapped inside) by a great rhombicosidodecahedron. The largest inscribed sphere always touches all 12 congruent regular decagonal faces but does not touch any of 30 congruent square & any of 20 congruent regular hexagonal faces at all since all 12 decagonal faces are closest to the centre in all the faces. Thus, the inner radius is always equal to the normal distance ( $H_D$ ) of regular decagonal faces from the centre of a great rhombicosidodecahedron & is given as follows

$$R_i = H_D = \frac{a}{2} \sqrt{25 + 10\sqrt{5}} \approx 3.440954801a$$

Hence, the **volume of inscribed sphere** is given as

$$V_{inscribed} = \frac{4}{3} \pi (R_i)^3 = \frac{4}{3} \pi \left( \frac{a}{2} \sqrt{25 + 10\sqrt{5}} \right)^3 \approx 170.6575526a^3$$

- 2. Outer (circumscribed) radius ( $R_o$ ):** It is the radius of the smallest sphere circumscribing a great rhombicosidodecahedron or it's the radius of a spherical surface passing through all 120 vertices of a great rhombicosidodecahedron. It is from the eq(VII) as follows

$$R_o = \frac{a}{2} \sqrt{31 + 12\sqrt{5}} \approx 3.8023945a$$

Hence, the **volume of circumscribed sphere** is given as

$$V_{\text{circumscribed}} = \frac{4}{3} \pi (R_o)^3 = \frac{4}{3} \pi \left( \frac{a}{2} \sqrt{31 + 12\sqrt{5}} \right)^3 = 230.2820721a^3$$

- 3. Surface area ( $A_s$ ):** We know that a great rhombicosidodecahedron has 30 congruent square faces, 20 congruent regular hexagonal faces & 12 congruent regular decagonal faces each of edge length  $a$ . Hence, its surface area is given as follows

$$A_s = 30(\text{area of square}) + 20(\text{area of regular hexagon}) + 12(\text{area of regular decagon})$$

We know that **area of any regular n-polygon** with each side of length  $a$  is given as

$$A = \frac{1}{4} n a^2 \cot \frac{\pi}{n}$$

Hence, by substituting all the corresponding values in the above expression, we get

$$\begin{aligned} A_s &= 30 \times \left( \frac{1}{4} \times 4a^2 \cot \frac{\pi}{4} \right) + 20 \times \left( \frac{1}{4} \times 6a^2 \cot \frac{\pi}{6} \right) + 12 \times \left( \frac{1}{4} \times 10a^2 \cot \frac{\pi}{10} \right) \\ &= 30a^2 + 30\sqrt{3}a^2 + 30a^2 \sqrt{5 + 2\sqrt{5}} = 30 \left( 1 + \sqrt{3} + \sqrt{5 + 2\sqrt{5}} \right) a^2 \\ A_s &= 30 \left( 1 + \sqrt{3} + \sqrt{5 + 2\sqrt{5}} \right) a^2 \approx 174.2920303a^2 \end{aligned}$$

- 4. Volume ( $V$ ):** We know that a great rhombicosidodecahedron with edge length  $a$  has 30 congruent square faces, 20 congruent regular hexagonal faces & 12 congruent regular decagonal faces. Hence, the volume ( $V$ ) of the great rhombicosidodecahedron is the sum of volumes of all its elementary right pyramids with square base, regular hexagonal base & regular decagonal base (face) (see figure 2 above) and is given as follows

$$\begin{aligned} V &= 30(\text{volume of right pyramid with square base}) \\ &\quad + 20(\text{volume of right pyramid with regular hexagonal base}) \\ &\quad + 12(\text{volume of right pyramid with regular decagonal base}) \\ &= 30 \left( \frac{1}{3} (\text{area of square}) \times H_s \right) + 20 \left( \frac{1}{3} (\text{area of regular hexagon}) \times H_h \right) \\ &\quad + 12 \left( \frac{1}{3} (\text{area of regular decagon}) \times H_D \right) \\ &= 30 \left( \frac{1}{3} \left( \frac{1}{4} \times 4a^2 \cot \frac{\pi}{4} \right) \times \frac{(3 + 2\sqrt{5})a}{2} \right) + 20 \left( \frac{1}{3} \left( \frac{1}{4} \times 6a^2 \cot \frac{\pi}{6} \right) \times \frac{\sqrt{3}(2 + \sqrt{5})a}{2} \right) \\ &\quad + 12 \left( \frac{1}{3} \left( \frac{1}{4} \times 10a^2 \cot \frac{\pi}{10} \right) \times \frac{a}{2} \sqrt{25 + 10\sqrt{5}} \right) \end{aligned}$$

$$\begin{aligned}
 &= 5(3 + 2\sqrt{5})a^3 + 15(2 + \sqrt{5})a^3 + 5a^3\sqrt{(5 + 2\sqrt{5})}\sqrt{(25 + 10\sqrt{5})} \\
 &= 5(9 + 5\sqrt{5})a^3 + 5a^3\sqrt{(10 + 5\sqrt{5})^2} = 5(9 + 5\sqrt{5})a^3 + 5(10 + 5\sqrt{5})a^3 = (95 + 50\sqrt{5})a^3 \\
 \mathbf{V} &= \mathbf{(95 + 50\sqrt{5})a^3 \approx 206.8033989a^3}
 \end{aligned}$$

**5. Mean radius ( $R_m$ ):** It is the radius of the sphere having a volume equal to that of a great rhombicosidodecahedron. It is calculated as follows

volume of sphere with mean radius  $R_m$  = volume of the great rhombicosidodecahedron

$$\frac{4}{3}\pi(R_m)^3 = (95 + 50\sqrt{5})a^3 \Rightarrow (R_m)^3 = \frac{3(95 + 50\sqrt{5})a^3}{4\pi} \text{ or } R_m = a\left(\frac{15(19 + 10\sqrt{5})}{4\pi}\right)^{\frac{1}{3}}$$

$$\mathbf{R_m = a\left(\frac{15(19 + 10\sqrt{5})}{4\pi}\right)^{\frac{1}{3}} \approx 3.668509834a}$$

It's clear from above results that  $R_i < R_m < R_o$

**6. Dihedral angles between the adjacent faces:** In order to calculate dihedral angles between the different adjacent faces with a common edge in a great rhombicosidodecahedron, let's consider one-by-one all three pairs of adjacent faces with a common edge as given below.

**a. Angle between square face & regular hexagonal face:** Draw the perpendiculars  $OO_1$  &  $OO_2$  from the centre O of great rhombicosidodecahedron to the square face & the regular hexagonal face which have a common edge (See figure 3). We know that the inscribed radius ( $r_i$ ) of any regular n-gon with each side  $a$  is given as follows

$$r_i = \text{inscribed radius of any regular } n - \text{gon} = \frac{a}{2} \cot \frac{\pi}{n}$$

$$\therefore O_1T = \text{inscribed radius of square} = \frac{a}{2} \cot \frac{\pi}{4} = \frac{a}{2} \text{ \&}$$

$$\therefore O_2T = \text{inscribed radius of regular hexagon} = \frac{a}{2} \cot \frac{\pi}{6} = \frac{a\sqrt{3}}{2}$$




Figure 3: Square face with centre  $O_1$  & regular hexagonal face with centre  $O_2$  with a common edge (denoted by point T) normal to the plane of paper.

In right  $\Delta OO_1T$  (Fig. 3),

$$\tan\theta_s = \frac{OO_1}{O_1T} = \frac{H_s}{\left(\frac{a}{2}\right)} = \frac{\left(\frac{(3 + 2\sqrt{5})a}{2}\right)}{\left(\frac{a}{2}\right)} = (3 + 2\sqrt{5})$$

$$\therefore \theta_s = \tan^{-1}(3 + 2\sqrt{5}) \approx 82.37736814^\circ \dots \dots \dots \text{(VIII)}$$

In right  $\Delta OO_2T$  (Fig. 3),

$$\tan\theta_h = \frac{OO_2}{O_2T} = \frac{H_h}{\left(\frac{a\sqrt{3}}{2}\right)} = \frac{\left(\frac{\sqrt{3}(2+\sqrt{5})a}{2}\right)}{\left(\frac{a\sqrt{3}}{2}\right)} = (2+\sqrt{5})$$

$$\therefore \theta_h = \tan^{-1}(2+\sqrt{5}) \approx 76.71747441^\circ \quad \dots \dots \dots (IX)$$

$$\begin{aligned} \Rightarrow \theta_s + \theta_h &= \tan^{-1}(3+2\sqrt{5}) + \tan^{-1}(2+\sqrt{5}) = \tan^{-1}\left(\frac{(3+2\sqrt{5})+(2+\sqrt{5})}{1-(3+2\sqrt{5})(2+\sqrt{5})}\right) \\ &= \tan^{-1}\left(\frac{5+3\sqrt{5}}{1-(16+7\sqrt{5})}\right) = \tan^{-1}\left(\frac{-(5+3\sqrt{5})}{15+7\sqrt{5}}\right) = \pi - \tan^{-1}\left(\frac{5+3\sqrt{5}}{15+7\sqrt{5}}\right) \\ &= \pi - \tan^{-1}\left(\frac{(5+3\sqrt{5})(7\sqrt{5}-15)}{(7\sqrt{5}+15)(7\sqrt{5}-15)}\right) = \pi - \tan^{-1}\left(\frac{30-10\sqrt{5}}{20}\right) = \pi - \tan^{-1}\left(\frac{3-\sqrt{5}}{2}\right) \end{aligned}$$

Hence, dihedral angle between the square face & the regular hexagonal face is given as

$$\theta_s + \theta_h = \pi - \tan^{-1}\left(\frac{3-\sqrt{5}}{2}\right) \approx 159.0948426^\circ$$

- b. Angle between square face & regular decagonal face:** Draw the perpendiculars  $OO_1$  &  $OO_3$  from the centre O of great rhombicosidodecahedron to the square face & the regular decagonal face which have a common edge (see figure 4).

$$O_3U = \text{inscribed radius of regular decagon} = \frac{a}{2} \cot \frac{\pi}{10} = \frac{a\sqrt{5+2\sqrt{5}}}{2}$$

In right  $\Delta OO_3U$  (Fig. 4),

$$\tan\theta_D = \frac{OO_3}{O_3U} = \frac{H_D}{\left(\frac{a\sqrt{5+2\sqrt{5}}}{2}\right)} = \frac{\left(\frac{a}{2}\sqrt{25+10\sqrt{5}}\right)}{\left(\frac{a\sqrt{5+2\sqrt{5}}}{2}\right)} = \sqrt{\frac{5(5+2\sqrt{5})}{5+2\sqrt{5}}} = \sqrt{5}$$

$$\therefore \theta_D = \tan^{-1}(\sqrt{5}) \approx 65.90515745^\circ \quad \dots \dots \dots (X)$$

$$\Rightarrow \theta_s + \theta_D = \tan^{-1}(3+2\sqrt{5}) + \tan^{-1}(\sqrt{5}) = \tan^{-1}\left(\frac{(3+2\sqrt{5})+(\sqrt{5})}{1-(3+2\sqrt{5})(\sqrt{5})}\right)$$

$$= \tan^{-1}\left(\frac{(3+3\sqrt{5})}{1-(10+3\sqrt{5})}\right) = \tan^{-1}\left(\frac{-3(1+\sqrt{5})}{9+3\sqrt{5}}\right) = \pi - \tan^{-1}\left(\frac{1+\sqrt{5}}{3+\sqrt{5}}\right)$$

$$= \pi - \tan^{-1}\left(\frac{(1+\sqrt{5})(3-\sqrt{5})}{(3+\sqrt{5})(3-\sqrt{5})}\right) = \pi - \tan^{-1}\left(\frac{2(\sqrt{5}-1)}{4}\right) = \pi - \tan^{-1}\left(\frac{\sqrt{5}-1}{2}\right)$$

Hence, dihedral angle between the square face & the regular decagonal face (see Fig. 4) is given as

$$\theta_s + \theta_D = \pi - \tan^{-1}\left(\frac{\sqrt{5}-1}{2}\right) \approx 148.2825256^\circ$$




Figure 4: Square face with centre  $O_1$  & regular decagonal face with centre  $O_3$  with a common edge (denoted by point U) normal to the plane of paper.

- c. **Angle between regular hexagonal face & regular decagonal face:** Draw the perpendiculars  $OO_2$  &  $OO_3$  from the centre  $O$  of great rhombicosidodecahedron to the regular hexagonal face & the regular decagonal face which have a common edge (see figure 5). Now from eq(IX) & (X), we get

$$\begin{aligned} \theta_h + \theta_D &= \tan^{-1}(2 + \sqrt{5}) + \tan^{-1}(\sqrt{5}) = \tan^{-1}\left(\frac{(2 + \sqrt{5}) + (\sqrt{5})}{1 - (2 + \sqrt{5})(\sqrt{5})}\right) = \tan^{-1}\left(\frac{(2 + 2\sqrt{5})}{1 - (5 + 2\sqrt{5})}\right) \\ &= \tan^{-1}\left(\frac{-2(1 + \sqrt{5})}{4 + 2\sqrt{5}}\right) \\ &= \pi - \tan^{-1}\left(\frac{1 + \sqrt{5}}{2 + \sqrt{5}}\right) \\ &= \pi - \tan^{-1}\left(\frac{(1 + \sqrt{5})(\sqrt{5} - 2)}{(\sqrt{5} + 2)(\sqrt{5} - 2)}\right) = \pi - \tan^{-1}\left(\frac{3 - \sqrt{5}}{1}\right) \\ &= \pi - \tan^{-1}(3 - \sqrt{5}) \end{aligned}$$

Hence, dihedral angle between the regular hexagonal face & the regular decagonal face (see Fig. 5) is given as

$$\theta_h + \theta_D = \pi - \tan^{-1}(3 - \sqrt{5}) \approx 142.6226319^\circ$$




Figure 5: Regular hexagonal face with centre  $O_2$  & regular decagonal face with centre  $O_3$  with a common edge (denoted by point  $V$ ) normal to the plane of paper.

#### 4. Construction of a solid great rhombicosidodecahedron

In order to construct a solid great rhombicosidodecahedron with edge length  $a$  there are two methods.

**4.1. Construction from elementary right pyramids:** In this method, first we construct all elementary right pyramids as follows

**Step 1.** Construct 30 congruent right pyramids with square base of side length  $a$  & normal height ( $H_s$ )

$$H_s = \frac{(3 + 2\sqrt{5})a}{2} \approx 3.736067978a$$

**Step 2.** Construct 20 congruent right pyramids with regular hexagonal base of side length  $a$  & normal height ( $H_h$ )

$$H_h = \frac{\sqrt{3}(2 + \sqrt{5})a}{2} \approx 3.668542481a$$

**Step 3.** Construct 12 congruent right pyramids with regular decagonal base of side length  $a$  & normal height ( $H_D$ )

$$H_D = \frac{a}{2} \sqrt{25 + 10\sqrt{5}} \approx 3.440954801a$$

**Step 4.** Now, paste/bond by joining all these elementary right pyramids by overlapping their lateral surfaces & keeping their apex points coincident with each other such that 4 edges of each square base (face) coincide with the edges of 2 regular hexagonal bases & 2 regular decagonal bases (faces). Thus a solid great rhombicosidodecahedron, with 30 congruent square faces, 20 congruent regular hexagonal faces & 12 congruent regular decagonal faces each of edge length  $a$ , is obtained.

**4.2. Facing a solid sphere:** It is a method of facing, first we select a **blank as a solid sphere** of certain material (i.e. metal, alloy, composite material etc.) & with suitable diameter in order to obtain the maximum desired edge length of a great rhombicosidodecahedron. Then, we perform the facing operations on the solid sphere to generate 30 congruent square faces, 20 congruent regular hexagonal faces & 12 congruent regular decagonal faces each of equal edge length.

**4.2.1. Edge length:** Let there be a blank as a solid sphere with a diameter  $D$ . Then the edge length  $a$ , of a great rhombicosidodecahedron of the maximum volume to be produced, can be co-related with the diameter  $D$  by **relation of outer radius ( $R_o$ ) with edge length ( $a$ ) of the great rhombicosidodecahedron** as follows

$$R_o = \frac{a}{2} \sqrt{31 + 12\sqrt{5}}$$

Now, substituting  $R_o = D/2$  in the above expression, we have

$$\frac{D}{2} = \frac{a}{2} \sqrt{31 + 12\sqrt{5}} \quad \text{or} \quad a = \frac{D}{\sqrt{31 + 12\sqrt{5}}}$$

$$a = \frac{D}{\sqrt{31 + 12\sqrt{5}}} \approx \mathbf{0.131496087D}$$

Above relation is very useful for determining the edge length  $a$  of a great rhombicosidodecahedron to be produced from a solid sphere with known diameter  $D$  for manufacturing purpose.

**4.2.2. Maximum volume of great rhombicosidodecahedron** produced from a solid sphere is given as follows

$$V_{max} = (95 + 50\sqrt{5})a^3 = (95 + 50\sqrt{5}) \left( \frac{D}{\sqrt{31 + 12\sqrt{5}}} \right)^3 = \frac{(95 + 50\sqrt{5})D^3}{(31 + 12\sqrt{5})\sqrt{31 + 12\sqrt{5}}}$$

$$= \frac{(95 + 50\sqrt{5})(31 - 12\sqrt{5})D^3}{241\sqrt{31 + 12\sqrt{5}}} = \frac{(-55 + 410\sqrt{5})D^3}{241\sqrt{31 + 12\sqrt{5}}} = \frac{5(82\sqrt{5} - 11)D^3}{241\sqrt{31 + 12\sqrt{5}}}$$

$$V_{max} = \frac{5(82\sqrt{5} - 11)D^3}{241\sqrt{31 + 12\sqrt{5}}} \approx \mathbf{0.470214661D^3}$$

**4.2.3. Minimum volume of material removed** is given as

$$(V_{removed})_{min} = (\text{volume of parent sphere with diameter } D) \\ - (\text{volume of great rhombicosidodecahedron})$$

$$= \frac{\pi}{6} D^3 - \frac{5(82\sqrt{5} - 11)D^3}{241\sqrt{31 + 12\sqrt{5}}} = \left( \frac{\pi}{6} - \frac{5(82\sqrt{5} - 11)}{241\sqrt{31 + 12\sqrt{5}}} \right) D^3$$

$$(V_{removed})_{min} = \left( \frac{\pi}{6} - \frac{5(82\sqrt{5} - 11)}{241\sqrt{31 + 12\sqrt{5}}} \right) D^3 \approx \mathbf{0.053384114D^3}$$

**4.2.4. Percentage (%) of minimum volume of material removed**

$$\% \text{ of } V_{\text{removed}} = \frac{\text{minimum volume removed}}{\text{total volume of sphere}} \times 100$$

$$= \frac{\left(\frac{\pi}{6} - \frac{5(82\sqrt{5} - 11)}{241\sqrt{31 + 12\sqrt{5}}}\right) D^3}{\frac{\pi}{6} D^3} \times 100 = \left(1 - \frac{30(82\sqrt{5} - 11)}{241\pi\sqrt{31 + 12\sqrt{5}}}\right) \times 100 \approx 10.19\%$$

It's obvious that when a great rhombicosidodecahedron of the maximum volume is produced from a solid sphere then about 10.19% volume of material is removed as scraps. Thus, we can select optimum diameter of blank as a solid sphere to produce a solid great rhombicosidodecahedron of the maximum volume (or with the maximum desired edge length).

**Conclusions:** Let there be any great rhombicosidodecahedron having 30 congruent square faces, 20 congruent regular hexagonal faces & 12 congruent regular decagonal faces each with edge length  $a$  then all its important parameters are calculated/determined as tabulated below.

Congruent polygonal faces	No. of faces	Normal distance of each face from the centre of the great rhombicosidodecahedron	Solid angle subtended by each face at the centre of the great rhombicosidodecahedron
Square	30	$\frac{(3 + 2\sqrt{5})a}{2} \approx 3.736067978a$	$4 \sin^{-1}\left(\frac{5 - 2\sqrt{5}}{30}\right) \approx 0.070385504 \text{ sr}$
Regular hexagon	20	$\frac{\sqrt{3}(2 + \sqrt{5})a}{2} \approx 3.668542481a$	$2\pi - 12 \sin^{-1}\left(\frac{1}{2}\sqrt{\frac{5 + 2\sqrt{5}}{10}}\right) \approx 0.184524626 \text{ sr}$
Regular decagon	12	$\frac{a}{2}\sqrt{25 + 10\sqrt{5}} \approx 3.440954801a$	$2\pi - 20 \sin^{-1}\left(\frac{5 - \sqrt{5}}{4\sqrt{6}}\right) \approx 0.563692745 \text{ sr}$

Inner (inscribed) radius ( $R_i$ )	$R_i = \frac{a}{2}\sqrt{25 + 10\sqrt{5}} \approx 3.440954801a$
Outer (circumscribed) radius ( $R_o$ )	$R_o = \frac{a}{2}\sqrt{31 + 12\sqrt{5}} \approx 3.8023945a$
Mean radius ( $R_m$ )	$R_m = a\left(\frac{15(19 + 10\sqrt{5})}{4\pi}\right)^{\frac{1}{3}} \approx 3.668509834a$
Surface area ( $A_s$ )	$A_s = 30\left(1 + \sqrt{3} + \sqrt{5 + 2\sqrt{5}}\right)a^2 \approx 174.2920303a^2$

<b>Volume (<math>V</math>)</b>	$V = (95 + 50\sqrt{5})a^3 \approx 206.8033989a^3$
--------------------------------	---

**Table for the dihedral angles between the adjacent faces of a great rhombicosidodecahedron**

Pair of the adjacent faces with a common edge	Square & regular hexagon	Square & regular decagon	Regular hexagon & regular decagon
Dihedral angle of the corresponding pair (of the adjacent faces)	$\theta_s + \theta_h = \pi - \tan^{-1}\left(\frac{3 - \sqrt{5}}{2}\right)$ $\approx 159.0948426^\circ$	$\theta_s + \theta_D = \pi - \tan^{-1}\left(\frac{\sqrt{5} - 1}{2}\right)$ $\approx 148.2825256^\circ$	$\theta_h + \theta_D = \pi - \tan^{-1}(3 - \sqrt{5})$ $\approx 142.6226319^\circ$

**Note:** Above articles had been developed & illustrated by Mr H.C. Rajpoot (B Tech, Mechanical Engineering)

M.M.M. University of Technology, Gorakhpur-273010 (UP) India

10 March, 2015

Email: [rajpootherishchandra@gmail.com](mailto:rajpootherishchandra@gmail.com), Author's Home Page: <https://notionpress.com/author/HarishChandraRajpoot>

### References

- [1] Thompson D. On the thirteen semi-regular solids of Archimedes, and on their development by the transformation of certain plane configurations. Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character. 1925 Feb 2;107(742):181-8.
- [2] Rajpoot HC. Mathematical analysis of truncated dodecahedron. 2014 Dec. doi:10.13140/RG.2.2.11319.12963.
- [3] Rajpoot HC. HCR's Theory of Polygon (proposed by harish chandra rajpoot) solid angle subtended by any polygonal plane at any point in the space. Int. J. Math. Phys. Sci. Res. 2014;2:28-56.
- [4] Rajpoot HC. HCR's Theory of Polygon. Solid angle subtended by any polygonal plane at any point in the space. 2019.
- [5] Rajpoot CH. Advanced geometry. 1st ed. Chennai: Notion Press Media Pvt. Ltd.; 2013 Apr 3. ISBN: 97893-83808-15-1.